STOPAIN CLINICAL- menthol gel Troy Manufacturing, Inc.

Stopain Clinical maximun Perfomance Pain Relief

Drug Facts

Active Ingredient

Menthol 10.0%

Purpose

Topical Analgesic

Uses

For the temporary relief of minor aches and pains of muscles and joints associated with:

- simple backache
- arthritis
- strains
- bruises
- sprains

Warnings

Avoid contact with eyes. For external use only.

Flammable: Keep away from fire or flame.

When using this product

- use only as directed
- do not bandage tightly or use a heating pad
- do not apply to wounds or damaged skin

Stop use and ask a doctor if

- condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days
- redness is present
- excessive irritation of the skin develops

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children over 12 years:

ask a doctor. Children 12 years or younger:

- apply directly onto affected area without the need to bandage
- repeat if necessary, but do not apply more than 4 times daily.

Other Information

Store at room temperature

Inactive Ingredients

acrylic acid/vinyl ester copolymer, dimethylsulfone (MSM), eucalyptus oil, glucosamine sulfate, hydroxypropylcellulose, PEG-8 dimethicone, pentylene glycol, peppermint oil, SD alcohol 39C, triethanolamine, water (USP).

Package Labeling:

Drug Facts	
Active Ingredient Menthol 10.0%	Purpose Topical Analgesic
Uses For the temporary relief of muscles and joints associated wit arthritis ■ strains ■ bruises ■ strains ■ bruises ■ strains ■ strains ■ bruises ■ strains □ stra	h: ■ simple backache
Warnings For external use or	ily. Avoid contact with eyes.
Flammable: Keep away from fir	e or flame.
When using this product ■ use bandage tightly or use a heating pwounds or damaged skin	
Stop use and ask a doctor if ■ of symptoms persist for more than 7 again within a few days ■ rednes irritation of the skin develops	days or clear up and occur
If pregnant or breast-feeding, a before use.	sk a hea l th professional
Keep out of reach of children. If help or contact a Poison Control C	
Directions Adults and childred directly onto affected area without repeat if necessary, but do not apple Children 12 years or younger: as	en over 12 years: apply the need to bandage only more than 4 times daily. sk a doctor.
Other Information ■ Store a	at room temperature
Inactive Ingredients acrylic dimethylsulfone (MSM), eucalyptu hydroxypropylcellulose, PEG-8 dir peppermint oil, SD alcohol 39C, tr	c acid/vinyl ester copolymer, is oil, glucosamine sulfate, nethicone, pentylene glycol,



STOPAIN CLINICAL

menthol gel

D	- #	lufaum atian
Proc	IUCT	Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:63936-9758

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	100 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
ACRYLIC ACID (UNII: J94PBK7X8S)	
DIMETHYL SULFONE (UNII: 9H4PO4Z4FT)	
EUCALYPTUS OIL (UNII: 2R040NI662)	
GLUCOSAMINE SULFATE (UNII: 1FW7WLR731)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
PEG-8 DIMETHICONE (UNII: GIA7T764OD)	
PENTYLENE GLYCOL (UNII: 50C1307PZG)	
PEPPERMINT OIL (UNII: AV092KU4JH)	
TROLAMINE (UNII: 903K93S3TK)	
WATER (UNII: 059QF0KO0R)	

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:63936- 9758-1	3 g in 1 PACKET; Type 0: Not a Combination Product	11/30/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	11/30/2022	

Labeler - Troy Manufacturing, Inc. (160075248)

Revised: 10/2023 Troy Manufacturing, Inc.