

**HEAD CONGESTION AND FLU SEVERE RELIEF- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated
Rite Aid Corporation**

Rite Aid 44-546-HCFS

Active ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 100 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves these symptoms due to the common cold:
 - headache
 - nasal congestion
 - cough
 - sore throat
 - minor aches and pains
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- diabetes
- liver disease
- heart disease
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

NDC 11822-5462-9

*Compare to the active ingredients in

SUDAFED PE® HEAD CONGESTION + FLU SEVERE

HEAD CONGESTION & FLU SEVERE RELIEF PE

ACETAMINOPHEN 325 mg

DEXTROMETHORPHAN HBr 10 mg

GUAIFENESIN 100 mg • **PHENYLEPHRINE HCl** 5 mg

PAIN RELIEVER/ FEVER REDUCER • COUGH SUPPRESSANT

EXPECTORANT • NASAL DECONGESTANT

Relieves sinus pressure and congestion, sinus

headache, sore throat, chest congestion and cough

ACTUAL SIZE

20
CAPLETS

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark SUDAFED PE® HEAD CONGESTION + FLU SEVERE. 50844 REV0722F54609

**SATISFACTION
GUARANTEE**

**If you're not satisfied, we'll
happily refund your money**

DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE
CAMP HILL, PA 17011

www.riteaid.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

No print/No varnish area
Lot no/Exp date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



B-1702-546-09-R2
REV072254609

Drug Facts (continued)
Uses ■ temporarily relieves these symptoms due to the common cold:
 ■ headache ■ nasal congestion ■ cough ■ minor aches and pains
 ■ sore throat ■ sinus congestion and pressure
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Drug Facts
Active ingredients (in each caplet)
 Acetaminophen 325 mg
 Dextromethorphan HBr 10 mg
 Guaifenesin 100 mg
 Phenylephrine HCl 5 mg
Purpose
 Pain reliever/fever reducer
 Cough suppressant
 Expectorant
 Nasal decongestant

NDC 11822-5462-9

*Compare to the active ingredients in
SUDAFED PE® HEAD CONGESTION + FLU SEVERE

HEAD CONGESTION & FLU SEVERE RELIEF PE

ACETAMINOPHEN 325 mg
DEXTROMETHORPHAN HBr 10 mg
GUAIFENESIN 100 mg • PHENYLEPHRINE HCl 5 mg

PAIN RELIEVER / FEVER REDUCER • COUGH SUPPRESSANT
 EXPECTORANT • NASAL DECONGESTANT

Relieves sinus pressure and congestion, sinus headache, sore throat, chest congestion and cough



20
CAPLETS

DISTRIBUTED BY:
 RITE AID, 30 HUNTER LANE,
 CAMP HILL, PA 17011
 www.riteaid.com

SATISFACTION GUARANTEE
 If you're not satisfied, we'll happily refund your money.

PARENTS: Learn about teen medicine abuse
 www.StopMedicineAbuse.org

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark SUDAFED PE® HEAD CONGESTION + FLU SEVERE. 50844 REV072254609

Drug Facts (continued)
Inactive ingredients com starch, crospovidone, FD & C yellow #6 aluminum lake, magnesium stearate, methocel, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)
Other information
 ■ each caplet contains: sodium 3 mg
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ see end flap for expiration date and lot number

Drug Facts (continued)
Directions
 ■ do not take more than directed
 ■ adults and children 12 years and over take 2 caplets every 4 hours
 ■ do not take more than 10 caplets in 24 hours
 ■ children under 12 years: ask a doctor

Warnings
 ■ difficulty in urination due to enlargement of the prostate gland
 ■ cough that occurs with too much phlegm (mucus)
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
 ■ When using this product do not exceed recommended dosage.
 ■ Stop use and ask a doctor if:
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
 ■ If pregnant or breast-feeding, ask a health professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)
Warnings
 ■ temporarily reduces fever
 ■ Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
 ■ 3 or more alcoholic drinks every day while using this product
 ■ with other drugs containing acetaminophen
 ■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin reddening
 ■ If a skin reaction occurs, stop use and seek medical help right away.
 ■ Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
 ■ Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ If you have ever had an allergic reaction to this product or any of its ingredients
 ■ Ask a doctor before use if you have ■ diabetes ■ high blood pressure ■ heart disease ■ thyroid disease ■ liver disease

Rite Aid 44-546

HEAD CONGESTION AND FLU SEVERE RELIEF
 acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-5462
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	44;546
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-5462-9	2 in 1 CARTON	05/05/2021	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/05/2021	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-5462) , pack(11822-5462)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-5462)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-5462)

Revised: 2/2024

Rite Aid Corporation