

BASIC CARE DAYTIME NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl

L. Perrigo Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Amazon Daytime Nighttime Cold & Flu Drug Facts

Active ingredients (in each softgel) – Nighttime Cold & Flu

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Active ingredients (in each softgel) – Daytime Cold & Flu

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose – Nighttime Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Purpose – Daytime Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses – Nighttime Cold & Flu

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Uses – Daytime Cold & Flu

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have – Nighttime Cold & Flu

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you have – Daytime Cold & Flu

- liver disease
- heart disease

- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are – Nighttime Cold & Flu

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

Ask a doctor or pharmacist before use if you are – Daytime Cold & Flu

taking the blood thinning drug warfarin

When using this product – Nighttime Cold & Flu

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

When using the product – Daytime Cold & Flu

do not use more than directed

Stop use and ask a doctor if – Nighttime Cold & Flu

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Stop use and ask a doctor if – Daytime Cold & Flu

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions – Nighttime Cold & Flu

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Directions – Daytime Cold & Flu

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients – Nighttime Cold & Flu

D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Inactive ingredients – Daytime Cold & Flu

edible ink*, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Daytime Relief

Compare to Vicks® DayQuil® Cold & Flu active ingredients

Non-Drowsy

Daytime cold & flu

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr

Pain Reliever, Fever Reducer, Cough Suppressant, Nasal Decongestant

Alcohol Free/Antihistamine Free

actual size

32 Softgels

Nighttime Relief

Compare to Vicks[®] NyQuil[®] Cold & Flu active ingredients

Nighttime cold & flu

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine

Alcohol Free

actual size

16 Softgels

basic+care[®]

Daytime Relief
Non-Drowsy

Compare to Vicks[®] DayQuil[®] Cold & Flu active ingredients**

daytime cold & flu

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr

Pain Reliever, Fever Reducer, Cough Suppressant, Nasal Decongestant

Alcohol Free/
Antihistamine Free



actual size

32 Softgels

Nighttime Relief

Compare to Vicks[®] NyQuil[®] Cold & Flu active ingredients**

nighttime cold & flu

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine

Alcohol Free



actual size

16 Softgels

Nighttime Cold & Flu

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses Temporarily relieves common cold/flu symptoms:
 ■ cough due to minor throat and bronchial irritation
 ■ minor aches and pains ■ sore throat ■ headache
 ■ runny nose and sneezing ■ fever

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ hives ■ rash
 If a skin reaction occurs, stop use and seek medical help right away. ▼

Daytime Cold & Flu

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses Temporarily relieves common cold/flu symptoms:
 ■ cough due to minor throat and bronchial irritation
 ■ minor aches and pains ■ sore throat ■ headache
 ■ nasal congestion ■ fever

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ hives ■ rash
 If a skin reaction occurs, stop use and seek medical help right away. ▼

Drug Facts (continued)

See throat warning. If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have:
 ■ heart disease ■ glaucoma
 ■ cough that occurs with bronchitis (trachea)
 ■ breathing problem such as emphysema or chronic bronchitis
 ■ peptic or duodenal ulcers or occurs with smoking, asthma, or emphysema
 ■ trouble urinating due to an enlarged prostate gland

Ask doctor or pharmacist before use if you are:
 ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product
 ■ stability may occur, especially in children
 ■ mild drowsiness may occur ■ avoid alcohol drinks
 ■ be careful when driving a motor vehicle or operating machinery
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
 ■ minor cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ▼

Drug Facts (continued)

See throat warning. If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have:
 ■ heart disease ■ diabetes
 ■ heart disease ■ high blood pressure ■ thyroid disease
 ■ trouble urinating due to an enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus)
 ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if:
 ■ you get nervous, dizzy or sleepless
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ▼

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Get medical attention if critical for adults as well as for child even if you do not notice any signs or symptoms.

Directions
 ■ take only as directed - see overdose warning
 ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients D & C yellow no. 10, redd 10, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, parabens, propylene glycol, purified water, sodium sorbitol solution. *may contain this ingredient.

Questions or comments? 1-800-789-9280

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Get medical attention if critical for adults as well as for child even if you do not notice any signs or symptoms.

Directions
 ■ take only as directed - see overdose warning
 ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4hrs
children 4 to under 12 yrs	ask a doctor
child under 4 yrs	do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients redd 10, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol, polydimethylsiloxane, purified water, sodium sorbitol solution. *may contain this ingredient.

Questions or comments? 1-800-789-9280

58072 AN 03



Made in Canada
Distributed by:
Perrigo
Allergan, MI 49010
GLUTEN FREE

DO NOT USE IF BLISTER UNITS ARE BROKEN OR TORN
**These products are not manufactured or distributed by Procter & Gamble, distributor of Vicks[®] DayQuil[®] Cold & Flu and Vicks[®] NyQuil[®] Cold & Flu.

PARENTS:
Learn about how medicine works.
www.StopTheDustBust.org

CONVENIENT RECYCLING TAB

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0 113-7580
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0 113-7580-72	1 in 1 CARTON; Type 0: Not a Combination Product	04/25/20 18	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	8 BLISTER PACK	16
Part 2	16 BLISTER PACK	32

Part 1 of 2

BASIC CARE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN (clear)	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	056
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/25/2018	

Part 2 of 2

BASIC CARE DAYTIME COLD AND FLUE

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	

SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	L994
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/25/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/25/2018	

Labeler - L. Perrigo Company (006013346)