

TIGECYCLINE- tigecycline injection, powder, lyophilized, for solution
Xellia Pharmaceuticals USA LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TIGECYCLINE FOR INJECTION, safely and effectively. See full prescribing information for TIGECYCLINE FOR INJECTION.

TIGECYCLINE for injection, for intravenous use Initial U.S. Approval: 2005

WARNING: ALL-CAUSE MORTALITY

See full prescribing information for complete boxed warning.

All-cause mortality was higher in patients treated with Tigecycline than comparators in a meta-analysis of clinical trials. The cause of this mortality risk difference of 0.6% (95% CI 0.1, 1.2) has not been established. Tigecycline should be reserved for use in situations when alternative treatments are not suitable (1.4, 5.1, 5.2, 6.1).

----- **RECENT MAJOR CHANGES** -----

Warnings and Precautions, Tooth Discoloration and Enamel Hypoplasia (5.6)

1/2020

Warnings and Precautions, Inhibition of Bone Growth (5.7)

1/2020

----- **INDICATIONS AND USAGE** -----

Tigecycline is a tetracycline class antibacterial indicated in patients 18 years of age and older for:

- Complicated skin and skin structure infections (1.1)
- Complicated intra-abdominal infections (1.2)
- Community-acquired bacterial pneumonia (1.3)

Limitations of Use: Tigecycline is not indicated for treatment of diabetic foot infection or hospital-acquired pneumonia, including ventilator-associated pneumonia (1.4).

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Tigecycline and other antibacterial drugs, Tigecycline should be used only to treat infections that are proven or strongly suspected to be caused by bacteria.

----- **DOSAGE AND ADMINISTRATION** -----

- Initial dose of 100 mg, followed by 50 mg every 12 hours administered intravenously over approximately 30 to 60 minutes. (2.1)
- Severe hepatic impairment (Child Pugh C): Initial dose of 100 mg followed by 25 mg every 12 hours. (2.2)

----- **DOSAGE FORMS AND STRENGTHS** -----

For Injection: 50 mg, lyophilized powder for reconstitution in a single-dose 10 mL vial. (3)

----- **CONTRAINDICATIONS** -----

- Known hypersensitivity to tigecycline. (4)

----- **WARNINGS AND PRECAUTIONS** -----

- **All-Cause Mortality:** A meta-analysis of Phase 3 and 4 clinical trials demonstrated an increase in all-cause mortality in Tigecycline -treated patients compared to controls with a risk difference of 0.6% (95% CI 0.1,1.2). The cause of this increase has not been established. An increase was also seen in a meta-analysis limited to the approved indications [0,6% (95% CI 0.0, 1.2)]. The greatest difference in mortality was seen in Tigecycline-treated patients with ventilator-associated pneumonia. (5.1, 5.2)
- **Anaphylactic Reactions:** have been reported with Tigecycline, and may be life-threatening. Avoid use in patients with known hypersensitivity to tetracyclines. (5.3)
- **Hepatic Adverse effects:** have been reported with Tigecycline. Patients who develop abnormal liver function tests during Tigecycline therapy should be monitored for evidence of worsening hepatic function and evaluated for risk/benefit of continuing tigecycline therapy. (5.4)
- **Pancreatitis,** including fatalities, has been reported with Tigecycline. If pancreatitis is suspected, then consider stopping Tigecycline. (5.5)
- **Tooth Discoloration and Enamel Hypoplasia:** The use of Tigecycline during tooth development (last half of pregnancy, infancy, and childhood to the age of 8 years) may cause permanent discoloration of the teeth (yellow-gray-brown) and enamel hypoplasia. (5.6)

- Inhibition of Bone Growth: The use of Tigecycline during the second and third trimester of pregnancy, infancy, and childhood up to the age of 8 years may cause reversible inhibition of bone growth. (5.7)
- Clostridium difficile associated diarrhea (CDAD): evaluate if diarrhea occurs. (5.8)

----- ADVERSE REACTIONS -----

The most common adverse reactions (incidence >5%) are nausea, vomiting, diarrhea, abdominal pain, headache, and increased SGPT. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Xellia Pharmaceuticals USA, LLC at safety@xellia.com or 1-833-295-6953, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- DRUG INTERACTIONS -----

- Suitable anticoagulation test should be monitored if Tigecycline is administered to patients receiving warfarin. (7.1)

----- USE IN SPECIFIC POPULATIONS -----

- Lactation: Avoid breastfeeding for longer than 3 weeks while taking Tigecycline. A lactating woman may also pump and discard breast milk during treatment and for 9 days after the last dose of Tigecycline (8.2)
- Pediatrics: Use in patients under 18 years of age is not recommended. Pediatric trials were not conducted because of the higher risk of mortality seen in adult trials (8.4)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: ALL-CAUSE MORTALITY

1 INDICATIONS AND USAGE

- 1.1 Complicated Skin and Skin Structure Infections
- 1.2 Complicated Intra-abdominal Infections
- 1.3 Community-Acquired Bacterial Pneumonia
- 1.4 Limitations of Use
- 1.5 Usage

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Adult Dosage
- 2.2 Dosage in Patients With Hepatic Impairment
- 2.3 Dosage in Pediatric Patients
- 2.4 Preparation and Administration
- 2.5 Drug Compatibilities
- 2.6 Drug Incompatibilities

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 All-Cause Mortality
- 5.2 Mortality Imbalance and Lower Cure Rates in Hospital-Acquired Pneumonia
- 5.3 Anaphylactic Reactions
- 5.4 Hepatic Adverse Effects
- 5.5 Pancreatitis
- 5.6 Tooth Discoloration and Enamel Hypoplasia
- 5.7 Inhibition of Bone Growth
- 5.8 *Clostridium difficile* Associated Diarrhea
- 5.9 Sepsis/Septic Shock in Patients with Intestinal Perforation
- 5.10 Tetracycline-Class Adverse Effects
- 5.11 Development of Drug-Resistant Bacteria

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

6.2 Post-Marketing Experience

7 DRUG INTERACTIONS

7.1 Warfarin

7.2 Oral Contraceptives

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

14.1 Complicated Skin and Skin Structure Infections

14.2 Complicated Intra-abdominal Infections

14.3 Community-Acquired Bacterial Pneumonia

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: ALL-CAUSE MORTALITY

An increase in all-cause mortality has been observed in a meta-analysis of Phase 3 and 4 clinical trials in Tigecycline-treated patients versus comparator. The cause of this mortality risk difference of 0.6% (95% CI 0.1, 1.2) has not been established. Tigecycline should be reserved for use in situations when alternative treatments are not suitable [see Indications and Usage (1.4), Warnings and Precautions (5.1, 5.2) and Adverse Reactions (6.1)].

1 INDICATIONS AND USAGE

1.1 Complicated Skin and Skin Structure Infections

Tigecycline for injection is indicated in patients 18 years of age and older for the treatment of complicated skin and skin structure infections caused by susceptible isolates of *Escherichia coli*, *Enterococcus faecalis* (vancomycin-susceptible isolates), *Staphylococcus aureus* (methicillin-susceptible and -resistant isolates), *Streptococcus agalactiae*, *Streptococcus anginosus* grp. (includes *S. anginosus*, *S. intermedius*, and *S. constellatus*), *Streptococcus pyogenes*, *Enterobacter cloacae*, *Klebsiella pneumoniae*, and *Bacteroides fragilis*.

1.2 Complicated Intra-abdominal Infections

Tigecycline for injection is indicated in patients 18 years of age and older for the treatment of complicated intra-abdominal infections caused by susceptible isolates of *Citrobacter freundii*, *Enterobacter cloacae*, *Escherichia coli*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Enterococcus faecalis* (vancomycin-susceptible isolates) *Staphylococcus aureus* (methicillin-susceptible and -resistant isolates), *Streptococcus anginosus* grp. (includes *S. anginosus*, *S. intermedius*, and *S. constellatus*), *Bacteroides fragilis*, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, *Bacteroides vulgatus*, *Clostridium perfringens*, and *Peptostreptococcus micros*.

1.3 Community-Acquired Bacterial Pneumonia

Tigecycline for injection is indicated in patients 18 years of age and older for the treatment of community-acquired bacterial pneumonia caused by susceptible isolates of *Streptococcus pneumoniae* (penicillin-susceptible isolates), including cases with concurrent bacteremia, *Haemophilus influenzae*, and *Legionella pneumophila*.

1.4 Limitations of Use

Tigecycline is not indicated for the treatment of diabetic foot infections. A clinical trial failed to demonstrate non-inferiority of Tigecycline for treatment of diabetic foot infections. Tigecycline is not indicated for the treatment of hospital-acquired or ventilator-associated pneumonia. In a comparative clinical trial, greater mortality and decreased efficacy were reported in Tigecycline-treated patients [see Warnings and Precautions (5.2)].

1.5 Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Tigecycline and other antibacterial drugs, Tigecycline should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Appropriate specimens for bacteriological examination should be obtained in order to isolate and identify the causative organisms and to determine their susceptibility to tigecycline.

Tigecycline may be initiated as empiric monotherapy before results of these tests are known.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Adult Dosage

The recommended dosage regimen for Tigecycline for Injection, USP, is an initial dose of 100 mg, followed by 50 mg every 12 hours. Intravenous infusions of Tigecycline for Injection, USP, should be administered over approximately 30 to 60 minutes every 12 hours.

The recommended duration of treatment with Tigecycline for Injection, USP, for complicated skin and skin structure infections or for complicated intra-abdominal infections is 5 to 14 days. The recommended duration of treatment with Tigecycline for Injection, USP, for community-acquired bacterial pneumonia is 7 to 14 days. The duration of therapy should be guided by the severity and site of the infection and the patient's clinical and bacteriological progress.

2.2 Dosage in Patients With Hepatic Impairment

No dosage adjustment is warranted in patients with mild to moderate hepatic impairment (Child Pugh A and Child Pugh B). In patients with severe hepatic impairment (Child Pugh C), the initial dose of Tigecycline should be 100 mg followed by a reduced maintenance dose of 25 mg every 12 hours. Patients with severe hepatic impairment (Child Pugh C) should be treated with caution and monitored for

treatment response [see *Clinical Pharmacology (12.3)* and *Use in Specific Populations (8.6)*].

2.3 Dosage in Pediatric Patients

The safety and efficacy of the proposed pediatric dosing regimens have not been evaluated due to the observed increase in mortality associated with tigecycline in adult patients. Avoid use of tigecycline in pediatric patients unless no alternative antibacterial drugs are available. Under these circumstances, the following doses are suggested:

- Pediatric patients aged 8 to 11 years should receive 1.2 mg/kg of tigecycline every 12 hours intravenously to a maximum dose of 50 mg of tigecycline every 12 hours.
- Pediatric patients aged 12 to 17 years should receive 50 mg of tigecycline every 12 hours.

The proposed pediatric doses of tigecycline were chosen based on exposures observed in pharmacokinetic trials, which included small numbers of pediatric patients [see *Use in Specific Populations (8.4)* and *Clinical Pharmacology (12.3)*].

There are no data to provide dosing recommendations in pediatric patients with hepatic impairment.

2.4 Preparation and Administration

Each vial of Tigecycline for Injection, USP, should be reconstituted with 5.3 mL of 0.9% Sodium Chloride Injection, USP, 5% Dextrose Injection, USP, or Lactated Ringer's Injection, USP, to achieve a concentration of 10 mg/mL of tigecycline. (Note: Each vial contains a 6% overage. Thus, 5 mL of reconstituted solution is equivalent to 50 mg of the drug.) The vial should be gently swirled until the drug dissolves. Reconstituted solution must be transferred and further diluted for intravenous infusion. Withdraw 5 mL of the reconstituted solution from the vial and add to a 100 mL intravenous bag for infusion (for a 100 mg dose, reconstitute two vials; for a 50 mg dose, reconstitute one vial). The maximum concentration in the intravenous bag should be 1 mg/mL. The reconstituted solution should be yellow to orange in color; if not, the solution should be discarded. Parenteral drug products should be inspected visually for particulate matter and discoloration (e.g., green or black) prior to administration. Once reconstituted, Tigecycline may be stored at room temperature (not to exceed 25°C/77°F) for up to 24 hours (up to 6 hours in the vial and the remaining time in the intravenous bag). If the storage conditions exceed 25°C (77°F) after reconstitution, tigecycline should be used immediately. Alternatively, Tigecycline mixed with 0.9% Sodium Chloride Injection, USP, or 5% Dextrose Injection, USP, may be stored refrigerated at 2° to 8°C (36° to 46°F) for up to 48 hours following immediate transfer of the reconstituted solution into the intravenous bag.

Tigecycline may be administered intravenously through a dedicated line or through a Y-site. If the same intravenous line is used for sequential infusion of several drugs, the line should be flushed before and after infusion of Tigecycline with 0.9% Sodium Chloride Injection, USP, 5% Dextrose Injection, USP, or Lactated Ringer's Injection, USP. Injection should be made with an infusion solution compatible with tigecycline and with any other drug(s) administered via this common line.

2.5 Drug Compatibilities

Compatible intravenous solutions include 0.9% Sodium Chloride Injection, USP, 5% Dextrose Injection, USP, and Lactated Ringer's Injection, USP. When administered through a Y-site, Tigecycline is compatible with the following drugs or diluents when used with either 0.9% Sodium Chloride Injection, USP, or 5% Dextrose Injection, USP: amikacin, dobutamine, dopamine HCl, gentamicin, haloperidol, Lactated Ringer's, lidocaine HCl, metoclopramide, morphine, norepinephrine, piperacillin/tazobactam (EDTA formulation), potassium chloride, propofol, ranitidine HCl, theophylline, and tobramycin.

2.6 Drug Incompatibilities

The following drugs should not be administered simultaneously through the same Y-site as Tigecycline: amphotericin B, amphotericin B lipid complex, diazepam, esomeprazole and omeprazole.

3 DOSAGE FORMS AND STRENGTHS

For Injection: Each single-dose 10 mL glass vial contains 50 mg of tigecycline as an orange lyophilized powder for reconstitution.

4 CONTRAINDICATIONS

Tigecycline is contraindicated for use in patients who have known hypersensitivity to tigecycline. Reactions have included anaphylactic reactions [see *Warnings and Precautions (5.3) and Adverse Reactions (6.2)*].

5 WARNINGS AND PRECAUTIONS

5.1 All-Cause Mortality

An increase in all-cause mortality has been observed in a meta-analysis of Phase 3 and 4 clinical trials in Tigecycline-treated patients versus comparator-treated patients. In all 13 Phase 3 and 4 trials that included a comparator, death occurred in 4.0% (150/3788) of patients receiving Tigecycline and 3.0% (110/3646) of patients receiving comparator drugs. In a pooled analysis of these trials, based on a random effects model by trial weight, the adjusted risk difference of all-cause mortality was 0.6% (95% CI 0.1, 1.2) between Tigecycline and comparator-treated patients. An analysis of mortality in all trials conducted for approved indications (cSSSI, cIAI, and CABP), including post-market trials showed an adjusted mortality rate of 2.5% (66/2640) for tigecycline and 1.8% (48/2628) for comparator, respectively. The adjusted risk difference for mortality stratified by trial weight was 0.6% (95% CI 0.0, 1.2).

The cause of this mortality difference has not been established. Generally, deaths were the result of worsening infection, complications of infection or underlying co-morbidities.

Tigecycline should be reserved for use in situations when alternative treatments are not suitable [see *Boxed Warning Indications and Usage (1.4), Warnings and Precautions (5.2) and Adverse Reactions (6.1)*].

5.2 Mortality Imbalance and Lower Cure Rates in Hospital-Acquired Pneumonia

A trial of patients with hospital acquired, including ventilator-associated, pneumonia failed to demonstrate the efficacy of Tigecycline. In this trial, patients were randomized to receive Tigecycline (100 mg initially, then 50 mg every 12 hours) or a comparator. In addition, patients were allowed to receive specified adjunctive therapies. The sub-group of patients with ventilator-associated pneumonia who received Tigecycline had lower cure rates (47.9% versus 70.1% for the clinically evaluable population).

In this trial, greater mortality was seen in patients with ventilator-associated pneumonia who received Tigecycline (25/131 [19.1%] versus 15/122 [12.3%] in comparator-treated patients) [see *Adverse Reactions (6.1)*]. Particularly high mortality was seen among Tigecycline - treated patients with ventilator-associated pneumonia and bacteremia at baseline (9/18 [50.0%] versus 1/13 [7.7%] in comparator-treated patients).

5.3 Anaphylactic Reactions

Anaphylactic reactions have been reported with nearly all antibacterial agents, including Tigecycline, and may be life-threatening. Tigecycline is structurally similar to tetracycline-class antibiotics and should be avoided in patients with known hypersensitivity to tetracycline-class antibiotics.

5.4 Hepatic Adverse Effects

Increases in total bilirubin concentration, prothrombin time and transaminases have been seen in patients treated with tigecycline. Isolated cases of significant hepatic dysfunction and hepatic failure have been reported in patients being treated with tigecycline. Some of these patients were receiving multiple concomitant medications. Patients who develop abnormal liver function tests during tigecycline therapy should be monitored for evidence of worsening hepatic function and evaluated for risk/benefit of continuing tigecycline therapy. Hepatic dysfunction may occur after the drug has been discontinued.

5.5 Pancreatitis

Acute pancreatitis, including fatal cases, has occurred in association with tigecycline treatment. The diagnosis of acute pancreatitis should be considered in patients taking tigecycline who develop clinical symptoms, signs, or laboratory abnormalities suggestive of acute pancreatitis. Cases have been reported in patients without known risk factors for pancreatitis. Patients usually improve after tigecycline discontinuation. Consideration should be given to the cessation of the treatment with tigecycline in cases suspected of having developed pancreatitis [see *Adverse Reactions (6.2)*].

5.6 Tooth Discoloration and Enamel Hypoplasia

The use of Tigecycline during tooth development (last half of pregnancy, infancy, and childhood to the age of 8 years) may cause permanent discoloration of the teeth (yellow-gray-brown). This adverse reaction is more common during long-term use of tetracyclines, but it has been observed following repeated short-term courses. Enamel hypoplasia has also been reported. Advise the patient of the potential risk to the fetus if Tigecycline is used during the second or third trimester of pregnancy [see *Use in Specific Populations (8.1, 8.4)*].

5.7 Inhibition of Bone Growth

The use of Tigecycline during the second and third trimester of pregnancy, infancy and childhood up to the age of 8 years may cause reversible inhibition of bone growth. All tetracyclines form a stable calcium complex in any bone-forming tissue. A decrease in fibula growth rate has been observed in premature infants given oral tetracycline in doses of 25 mg/kg every 6 hours. This reaction was shown to be reversible when the tetracycline was discontinued. Advise the patient of the potential risk to the fetus if Tigecycline is used during the second or third trimester of pregnancy [see *Use in Specific Populations (8.1, 8.4)*].

5.8 *Clostridium difficile* Associated Diarrhea

Clostridium difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including Tigecycline, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

5.9 Sepsis/Septic Shock in Patients with Intestinal Perforation

Monotherapy with tigecycline should be avoided in patients with complicated intra-abdominal infections (cIAI) secondary to clinically apparent intestinal perforation. In cIAI studies (n=1642), 6 patients treated with Tigecycline and 2 patients treated with imipenem/cilastatin presented with intestinal perforations and developed sepsis/septic shock. The 6 patients treated with Tigecycline had higher APACHE II scores (median = 13) versus the 2 patients treated with imipenem/cilastatin (APACHE II scores = 4 and

6). Due to differences in baseline APACHE II scores between treatment groups and small overall numbers, the relationship of this outcome to treatment cannot be established.

5.10 Tetracycline-Class Adverse Effects

Tigecycline is structurally similar to tetracycline-class antibiotics and may have similar adverse effects. Such effects may include: photosensitivity, pseudotumor cerebri, and anti-anabolic action (which has led to increased BUN, azotemia, acidosis, and hyperphosphatemia).

5.11 Development of Drug-Resistant Bacteria

Prescribing Tigecycline in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

6 ADVERSE REACTIONS

The following serious adverse reactions are described elsewhere in the labeling:

- All-Cause Mortality [see Boxed Warning and Warnings and Precautions (5.1)]
- Mortality Imbalance and Lower Cure Rates in Hospital-Acquired Pneumonia [see Warnings and Precautions (5.2)]
- Anaphylaxis [see Warning and Precautions (5.3)]
- Hepatic Adverse Effects [see Warnings and Precautions (5.4)]
- Pancreatitis [see Warnings and Precautions (5.5)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In clinical trials, 2514 patients were treated with Tigecycline. Tigecycline was discontinued due to adverse reactions in 7% of patients compared to 6% for all comparators. Table 1 shows the incidence of adverse reactions through test of cure reported in $\geq 2\%$ of patients in these trials.

Table 1. Incidence (%) of Adverse Reactions through Test of Cure Reported in $\geq 2\%$ of Patients Treated in Clinical Studies

Body System Adverse Reactions	Tigecycline (N=2514)	Comparators* (N=2307)
Body as a Whole		
Abdominal pain	6	4
Abscess	2	2
Asthenia	3	2
Headache	6	7
Infection	7	5
Cardiovascular System		
Phlebitis	3	4
Digestive System		
Diarrhea	12	11
Dyspepsia	2	2
Nausea	26	13
Vomiting	18	9
Hemic and Lymphatic System		
Anemia	5	6

Metabolic and Nutritional

Alkaline Phosphatase Increased	3	3
Amylase Increased	3	2
Bilirubinemia	2	1
BUN Increased	3	1
Healing Abnormal	3	2
Hyponatremia	2	1
Hypoproteinemia	5	3
SGOT Increased [†]	4	5
SGPT Increased [†]	5	5

Respiratory System

Pneumonia	2	2
-----------	---	---

Nervous System

Dizziness	3	3
-----------	---	---

Skin and Appendages

Rash	3	4
------	---	---

* Vancomycin/Aztreonam, Imipenem/Cilastatin, Levofloxacin, Linezolid.

† LFT abnormalities in Tigecycline-treated patients were reported more frequently in the post therapy period than those in comparator-treated patients, which occurred more often on therapy.

In all 13 Phase 3 and 4 trials that included a comparator, death occurred in 4.0% (150/3788) of patients receiving Tigecycline and 3.0% (110/3646) of patients receiving comparator drugs. In a pooled analysis of these trials, based on a random effects model by trial weight, an adjusted risk difference of all-cause mortality was 0.6% (95% CI 0.1, 1.2) between Tigecycline and comparator-treated patients (see Table 2). The cause of the imbalance has not been established. Generally, deaths were the result of worsening infection, complications of infection or underlying co-morbidities.

Table 2. Patients with Outcome of Death by Infection Type

Infection Type	Tigecycline		Comparator		Risk Difference*
	n/N	%	n/N	%	% (95% CI)
cSSSI	12/834	1.4	6/813	0.7	0.7 (-0.3, 1.7)
cIAI	42/1382	3.0	1/1393	2.2	0.8 (-0.4, 2.0)
CAP	12/424	2.8	11/422	2.6	0.2 (-2.0, 2.4)
HAP	66/467	14.1	57/467	12.2	1.9 (-2.4, 6.3)
Non-VAP [†]	41/336	12.2	42/345	12.2	0.0 (-4.9, 4.9)
VAP [†]	25/131	19.1	15/122	12.3	6.8 (-2.1, 15.7)
RP	11/128	8.6	2/43	4.7	3.9 (-4.0, 11.9)
DFI	7/553	1.3	3/508	0.6	0.7 (-0.5, 1.8)
Overall Adjusted	150/3788	4.0	110/3646	3.0	0.6 (0.1, 1.2) [‡]

CAP = Community-acquired pneumonia; cIAI = Complicated intra-abdominal infections; cSSSI = Complicated skin and skin structure infections; HAP = Hospital-acquired pneumonia; VAP = Ventilator-associated pneumonia; RP = Resistant pathogens; DFI = Diabetic foot infections.

* The difference between the percentage of patients who died in Tigecycline and comparator treatment groups. The 95% CI for each infection type was calculated using the normal approximation method without continuity correction.

† These are subgroups of the HAP population.

‡ Overall adjusted (random effects model by trial weight) risk difference estimate and 95% CI.

Note: The studies include 300, 305, 900 (cSSSI), 301, 306, 315, 316, 400 (cIAI), 308 and 313 (CAP), 311 (HAP), 307 [Resistant gram-positive pathogen study in patients with MRSA or Vancomycin-Resistant Enterococcus (VRE)], and 319 (DFI with and without osteomyelitis).

An analysis of mortality in all trials conducted for approved indications - cSSSI, cIAI, and CABP, including post-market trials (one in cSSSI and two in cIAI) - showed an adjusted mortality rate of 2.5% (66/2640) for tigecycline and 1.8% (48/2628) for comparator, respectively. The adjusted risk difference for mortality stratified by trial weight was 0.6% (95% CI 0.0, 1.2).

In comparative clinical studies, infection-related serious adverse reactions were more frequently reported for subjects treated with Tigecycline (7%) versus comparators (6%). Serious adverse reactions of sepsis/septic shock were more frequently reported for subjects treated with Tigecycline (2%) versus comparators (1%). Due to baseline differences between treatment groups in this subset of patients, the relationship of this outcome to treatment cannot be established [see *Warnings and Precautions* (5.9)].

The most common adverse reactions were nausea and vomiting which generally occurred during the first 1-2 days of therapy. The majority of cases of nausea and vomiting associated with Tigecycline and comparators were either mild or moderate in severity. In patients treated with Tigecycline, nausea incidence was 26% (17% mild, 8% moderate, 1% severe) and vomiting incidence was 18% (11% mild, 6% moderate, 1% severe).

In patients treated for complicated skin and skin structure infections (cSSSI), nausea incidence was 35% for Tigecycline and 9% for vancomycin/aztreonam; vomiting incidence was 20% for Tigecycline and 4% for vancomycin/aztreonam. In patients treated for complicated intra-abdominal infections (cIAI), nausea incidence was 25% for Tigecycline and 21% for imipenem/cilastatin; vomiting incidence was 20% for Tigecycline and 15% for imipenem/cilastatin. In patients treated for community-acquired bacterial pneumonia (CABP), nausea incidence was 24% for Tigecycline and 8% for levofloxacin; vomiting incidence was 16% for Tigecycline and 6% for levofloxacin.

Discontinuation from tigecycline was most frequently associated with nausea (1%) and vomiting (1%). For comparators, discontinuation was most frequently associated with nausea (<1%).

The following adverse reactions were reported (<2%) in patients receiving Tigecycline in clinical studies:

Body as a Whole: injection site inflammation, injection site pain, injection site reaction, septic shock, allergic reaction, chills, injection site edema, injection site phlebitis

Cardiovascular System: thrombophlebitis

Digestive System: anorexia, jaundice, abnormal stools

Metabolic/Nutritional System: increased creatinine, hypocalcemia, hypoglycemia

Special Senses: taste perversion

Hemic and Lymphatic System: prolonged activated partial thromboplastin time (aPTT), prolonged prothrombin time (PT), eosinophilia, increased international normalized ratio (INR), thrombocytopenia

Skin and Appendages: pruritus

Urogenital System: vaginal moniliasis, vaginitis, leukorrhea

6.2 Post-Marketing Experience

The following adverse reactions have been identified during post-approval use of Tigecycline. Because these reactions are reported voluntarily from a population of uncertain size, it is not always

possible to reliably estimate their frequency or establish causal relationship to drug exposure.

- anaphylactic reactions
- acute pancreatitis
- hepatic cholestasis, and jaundice
- severe skin reactions, including Stevens-Johnson Syndrome
- symptomatic hypoglycemia in patients with and without diabetes mellitus

7 DRUG INTERACTIONS

7.1 Warfarin

Prothrombin time or other suitable anticoagulation test should be monitored if tigecycline is administered with warfarin [see *Clinical Pharmacology* (12.3)].

7.2 Oral Contraceptives

Concurrent use of antibacterial drugs with oral contraceptives may render oral contraceptives less effective.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Tigecycline, like other tetracycline class antibacterial drugs, may cause permanent discoloration of deciduous teeth and reversible inhibition of bone growth when administered during the second and third trimesters of pregnancy [see Warnings and Precautions (5.6) (5.7), Data, and Use in Specific Populations (8.4)]. There are no available data on the risk of major birth defects or miscarriage following the use of Tigecycline during pregnancy. Administration of intravenous tigecycline in pregnant rats and rabbits during the period of organogenesis was associated with reduction in fetal weights and an increased incidence of skeletal anomalies (delays in bone ossification) at exposures of 5 and 1 times the human exposure at the recommended clinical dose in rats and rabbits, respectively. Advise the patient of the potential risk to the fetus if Tigecycline is used during the second or third trimester.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U. S. general population, the estimated background risk in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Human Data

The use of tetracycline-class antibacterial drugs, that includes Tigecycline, during tooth development (second and third trimester of pregnancy) may cause permanent discoloration of deciduous teeth. This adverse reaction is more common during long-term use of tetracyclines but has been observed following repeated short-term courses. Tigecycline may cause reversible inhibition of bone growth when administered during the second and third trimesters of pregnancy. A decrease in fibula growth rate has been observed in premature infants given oral tetracycline in doses of 25 mg/kg every 6 hours.

Animal Data

In embryo-fetal development studies, tigecycline was administered during the period of organogenesis at doses up to 12 mg/kg/day in rats and 4 mg/kg in rabbits or 5 and 1 times the systemic exposure at the recommended clinical dose, respectively. In the rat study, decreased fetal weight and fetal skeletal variations (reduced ossification of the pubic, ischial, and supraoccipital bones and increased incidences

of rudimentary 14th rib) were observed in the presence of maternal toxicity at 12 mg/kg/day (5 times the recommended clinical dose based on systemic exposure). In rabbits, decreased fetal weights were observed in the presence of maternal toxicity at 4 mg/kg (equivalent to the human exposure at the recommended clinical dose).

In preclinical safety studies, ¹⁴C-labeled tigecycline crossed the placenta and was found in fetal tissues.

8.2 Lactation

Risk Summary

There are no data on the presence of tigecycline in human milk; however, tetracycline-class antibacterial drugs are present in breast milk. It is not known whether tigecycline has an effect on the breastfed infant or on milk production. Tigecycline has low oral bioavailability; therefore, infant exposure is expected to be low. Tigecycline is present in rat milk with little or no systemic exposure to tigecycline in nursing pups as a result of exposure via maternal milk. When a drug is present in animal milk, it is likely that the drug will be present in human milk.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Tigecycline and any potential adverse effects on the breastfed child from Tigecycline or from the underlying maternal condition (*see Clinical Considerations*).

Clinical Considerations

Because of the theoretical risk of dental discoloration and inhibition of bone growth, avoid breastfeeding if taking Tigecycline for longer than three weeks. A lactating woman may also consider interrupting breastfeeding and pumping and discarding breastmilk during administration of Tigecycline and for 9 days (approximately 5 half-lives) after the last dose in order to minimize drug exposure to a breastfed infant.

8.4 Pediatric Use

Use in patients under 18 years of age is not recommended. Safety and effectiveness in pediatric patients below the age of 18 years have not been established. Because of the increased mortality observed in tigecycline-treated adult patients in clinical trials, pediatric trials of tigecycline to evaluate the safety and efficacy of tigecycline were not conducted.

In situations where there are no other alternative antibacterial drugs, pediatric dosing has been proposed for pediatric patients 8 to 17 years of age based on data from pediatric pharmacokinetic studies [*see Dosage and Administration (2.3) and Clinical Pharmacology (12.3)*].

Because of effects on tooth development, use in patients under 8 years of age is not recommended [*see Warnings and Precautions (5.7)*].

8.5 Geriatric Use

Of the total number of subjects who received Tigecycline in Phase 3 clinical studies (n=2514), 664 were 65 and over, while 288 were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, but greater sensitivity to adverse events of some older individuals cannot be ruled out.

No significant difference in tigecycline exposure was observed between healthy elderly subjects and younger subjects following a single 100 mg dose of tigecycline [*see Clinical Pharmacology (12.3)*].

8.6 Hepatic Impairment

No dosage adjustment is warranted in patients with mild to moderate hepatic impairment (Child Pugh A and Child Pugh B). In patients with severe hepatic impairment (Child Pugh C), the initial dose of tigecycline should be 100 mg followed by a reduced maintenance dose of 25 mg every 12 hours. Patients with severe hepatic impairment (Child Pugh C) should be treated with caution and monitored for

treatment response [see *Clinical Pharmacology (12.3) and Dosage and Administration (2.2)*].

10 OVERDOSAGE

No specific information is available on the treatment of overdose with tigecycline. Intravenous administration of Tigecycline at a single dose of 300 mg over 60 minutes in healthy volunteers resulted in an increased incidence of nausea and vomiting. Tigecycline is not removed in significant quantities by hemodialysis.

11 DESCRIPTION

Tigecycline is a tetracycline class antibacterial for intravenous infusion. The chemical name of tigecycline is (4*S*,4*aS*,5*aR*,12*aS*)-9-[2-(*tert*-butylamino)acetamido]-4,7-bis(dimethylamino)-1,4,4*a*,5,5*a*,6,11,12*a*-octahydro-3,10,12,12*a*-tetrahydroxy-1,11-dioxo-2-naphthacene carboxamide. The empirical formula is C₂₉H₃₉N₅O₈ and the molecular weight is 585.65.

The following represents the chemical structure of tigecycline:




Figure 1: Structure of Tigecycline

Tigecycline is an orange lyophilized powder or cake. Each Tigecycline for Injection, USP, single-dose 10 mL vial contains 50 mg tigecycline lyophilized powder for reconstitution for intravenous infusion and 100 mg of lactose monohydrate. The pH is adjusted with hydrochloric acid, and if necessary sodium hydroxide. The product does not contain preservatives.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Tigecycline is a tetracycline class antibacterial [see *Clinical Microbiology (12.4)*].

12.2 Pharmacodynamics

Cardiac Electrophysiology

No significant effect of a single intravenous dose of Tigecycline 50 mg or 200 mg on QTc interval was detected in a randomized, placebo- and active-controlled four-arm crossover thorough QTc study of 46 healthy subjects.

12.3 Pharmacokinetics

The mean pharmacokinetic parameters of tigecycline after single and multiple intravenous doses based on pooled data from clinical pharmacology studies, are summarized in Table 3. Intravenous infusions of tigecycline were administered over approximately 30 to 60 minutes.

Table 3. Mean (CV%) Pharmacokinetic Parameters of Tigecycline

	Single Dose 100 mg (N=224)	Multiple Dose* 50 mg every 12h (N=103)
C_{max} (mcg/mL) [†]	1.45 (22%)	0.87 (27%)
C_{max} (mcg/mL) [‡]	0.90 (30%)	0.63 (15%)
AUC (mcg·h/mL)	5.19 (36%)	- -
AUC _{0-24h} (mcg·h/mL)	- -	4.70 (36%)
C_{min} (mcg/mL)	- -	0.13 (59%)
$t_{1/2}$ (h)	27.1 (53%)	42.4 (83%)
CL (L/h)	21.8 (40%)	23.8 (33%)
CL _r (mL/min)	38.0 (82%)	51.0 (58%)
V _{ss} (L)	568 (43%)	639 (48%)

* 100 mg initially, followed by 50 mg every 12 hours

† 30-minute infusion

‡ 60-minute infusion

Distribution

The *in vitro* plasma protein binding of tigecycline ranges from approximately 71% to 89% at concentrations observed in clinical studies (0.1 to 1.0 mcg/mL). The steady-state volume of distribution of tigecycline averaged 500 to 700 L (7 to 9 L/kg), indicating tigecycline is extensively distributed beyond the plasma volume and into the tissues.

Following the administration of tigecycline 100 mg, followed by 50 mg every 12 hours to 33 healthy volunteers, the tigecycline AUC_{0-12h} (134 mcg·h/mL) in alveolar cells was approximately 78-fold higher than the AUC_{0-12h} in the serum, and the AUC_{0-12h} (2.28 mcg·h/mL) in epithelial lining fluid was approximately 32% higher than the AUC_{0-12h} in serum. The AUC_{0-12h} (1.61 mcg·h/mL) of tigecycline in skin blister fluid was approximately 26% lower than the AUC_{0-12h} in the serum of 10 healthy subjects.

In a single-dose study, tigecycline 100 mg was administered to subjects prior to undergoing elective surgery or medical procedure for tissue extraction. Concentrations at 4 hours after tigecycline administration were higher in gallbladder (38-fold, n=6), lung (3.7-fold, n=5), and colon (2.3-fold, n=6), and lower in synovial fluid (0.58-fold, n=5), and bone (0.35-fold, n=6) relative to serum. The concentration of tigecycline in these tissues after multiple doses has not been studied.

Elimination

Metabolism

Tigecycline is not extensively metabolized. *In vitro* studies with tigecycline using human liver microsomes, liver slices, and hepatocytes led to the formation of only trace amounts of metabolites. In healthy male volunteers receiving ¹⁴C-tigecycline, tigecycline was the primary ¹⁴C-labeled material recovered in urine and feces, but a glucuronide, an N-acetyl metabolite, and a tigecycline epimer (each at no more than 10% of the administered dose) were also present.

Tigecycline is a substrate of P-glycoprotein (P-gp) based on an *in vitro* study using a cell line overexpressing P-gp. The potential contribution of P-gp-mediated transport to the *in vivo* disposition of tigecycline is not known.

Excretion

The recovery of total radioactivity in feces and urine following administration of ¹⁴C-tigecycline indicates that 59% of the dose is eliminated by biliary/fecal excretion, and 33% is excreted in urine. Approximately 22% of the total dose is excreted as unchanged tigecycline in urine. Overall, the primary route of elimination for tigecycline is biliary excretion of unchanged tigecycline and its metabolites.

Glucuronidation and renal excretion of unchanged tigecycline are secondary routes.

Specific Populations

Hepatic Impairment

In a study comparing 10 patients with mild hepatic impairment (Child Pugh A), 10 patients with moderate hepatic impairment (Child Pugh B), and 5 patients with severe hepatic impairment (Child Pugh C) to 23 age and weight matched healthy control subjects, the single-dose pharmacokinetic disposition of tigecycline was not altered in patients with mild hepatic impairment. However, systemic clearance of tigecycline was reduced by 25% and the half-life of tigecycline was prolonged by 23% in patients with moderate hepatic impairment (Child Pugh B). Systemic clearance of tigecycline was reduced by 55%, and the half-life of tigecycline was prolonged by 43% in patients with severe hepatic impairment (Child Pugh C). Dosage adjustment is necessary in patients with severe hepatic impairment (Child Pugh C) [see *Use in Specific Populations (8.6) and Dosage and Administration (2.2)*].

Renal Impairment

A single dose study compared 6 subjects with severe renal impairment (creatinine clearance <30 mL/min), 4 end stage renal disease (ESRD) patients receiving tigecycline 2 hours before hemodialysis, 4 ESRD patients receiving tigecycline 1 hour after hemodialysis, and 6 healthy control subjects. The pharmacokinetic profile of tigecycline was not significantly altered in any of the renally impaired patient groups, nor was tigecycline removed by hemodialysis. No dosage adjustment of Tigecycline is necessary in patients with renal impairment or in patients undergoing hemodialysis.

Geriatric Patients

No significant differences in pharmacokinetics were observed between healthy elderly subjects (n=15, age 65-75; n= 13, age >75) and younger (n=18) receiving a single 100mg dose of Tigecycline. Therefore, no dosage adjustment is necessary based on age [see *Use in Specific Populations (8.5)*].

Pediatric Patients

A single-dose safety, tolerability, and pharmacokinetic study of tigecycline in pediatric patients aged 8–16 years who recently recovered from infections was conducted. The doses administered were 0.5, 1, or 2 mg/kg. The study showed that for children aged 12–16 years (n = 16) a dosage of 50 mg twice daily would likely result in exposures comparable to those observed in adults with the approved dosing regimen. Large variability observed in children aged 8 to 11 years of age (n = 8) required additional study to determine the appropriate dosage.

A subsequent tigecycline dose-finding study was conducted in 8–11 year old patients with cIAI, cSSSI, or CABP. The doses of tigecycline studied were 0.75 mg/kg (n = 17), 1 mg/kg (n = 21), and 1.25 mg/kg (n=20). This study showed that for children aged 8–11 years, a 1.2 mg/kg dose would likely result in exposures comparable to those observed in adults resulting with the approved dosing regimen [see *Dosage and Administration (2.3)*].

Gender

In a pooled analysis of 38 women and 298 men participating in clinical pharmacology studies, there was no significant difference in the mean (\pm SD) tigecycline clearance between women (20.7 \pm 6.5 L/h) and men (22.8 \pm 8.7 L/h). Therefore, no dosage adjustment is necessary based on gender.

Race

In a pooled analysis of 73 Asian subjects, 53 Black subjects, 15 Hispanic subjects, 190 White subjects, and 3 subjects classified as "other" participating in clinical pharmacology studies, there was no significant difference in the mean (\pm SD) tigecycline clearance among the Asian subjects (28.8 \pm 8.8 L/h), Black subjects (23.0 \pm 7.8 L/h), Hispanic subjects (24.3 \pm 6.5 L/h), White subjects (22.1 \pm 8.9 L/h), and "other" subjects (25.0 \pm 4.8 L/h). Therefore, no dosage adjustment is necessary based on race.

Drug Interaction Studies

Digoxin

Tigecycline (100 mg followed by 50 mg every 12 hours) and digoxin (0.5 mg followed by 0.25 mg, orally, every 24 hours) were co-administered to healthy subjects in a drug interaction study. Tigecycline slightly decreased the C_{max} of digoxin by 13%, but did not affect the AUC or clearance of digoxin. This small change in C_{max} did not affect the steady-state pharmacodynamic effects of digoxin as measured by changes in ECG intervals. In addition, digoxin did not affect the pharmacokinetic profile of tigecycline. Therefore, no dosage adjustment of either drug is necessary when Tigecycline is administered with digoxin.

Warfarin

Concomitant administration of Tigecycline (100 mg followed by 50 mg every 12 hours) and warfarin (25 mg single-dose) to healthy subjects resulted in a decrease in clearance of R- warfarin and S-warfarin by 40% and 23%, an increase in C_{max} by 38% and 43% and an increase in AUC by 68% and 29%, respectively. Tigecycline did not significantly alter the effects of warfarin on INR. In addition, warfarin did not affect the pharmacokinetic profile of tigecycline. However, prothrombin time or other suitable anticoagulation test should be monitored if tigecycline is administered with warfarin.

In vitro studies in human liver microsomes indicate that tigecycline does not inhibit metabolism mediated by any of the following 6 cytochrome P450 (CYP) isoforms: 1A2, 2C8, 2C9, 2C19, 2D6, and 3A4. Therefore, Tigecycline is not expected to alter the metabolism of drugs metabolized by these enzymes. In addition, because tigecycline is not extensively metabolized, clearance of tigecycline is not expected to be affected by drugs that inhibit or induce the activity of these CYP450 isoforms.

In vitro studies using Caco-2 cells indicate that tigecycline does not inhibit digoxin flux, suggesting that tigecycline is not a P-glycoprotein (P-gp) inhibitor. This *in vitro* information is consistent with the lack of effect of tigecycline on digoxin clearance noted in the *in vivo* drug interaction study described above.

Tigecycline is a substrate of P-gp based on an *in vitro* study using a cell line overexpressing P-gp. The potential contribution of P-gp-mediated transport to the *in vivo* disposition of tigecycline is not known. Coadministration of P-gp inhibitors (e.g., ketoconazole or cyclosporine) or P-gp inducers (e.g., rifampicin) could affect the pharmacokinetics of tigecycline.

12.4 Microbiology

Mechanism of Action

Tigecycline inhibits protein translation in bacteria by binding to the 30S ribosomal subunit and blocking entry of amino-acyl tRNA molecules into the A site of the ribosome. This prevents incorporation of amino acid residues into elongating peptide chains. In general, tigecycline is considered bacteriostatic; however, Tigecycline has demonstrated bactericidal activity against isolates of *S. pneumoniae* and *L. pneumophila*.

Resistance

To date there has been no cross-resistance observed between tigecycline and other antibacterials. Tigecycline is less affected by the two major tetracycline-resistance mechanisms, ribosomal protection and efflux. Additionally, tigecycline is not affected by resistance mechanisms such as beta-lactamases (including extended spectrum beta-lactamases), target-site modifications, macrolide efflux pumps or enzyme target changes (e.g. gyrase/topoisomerases). However, some ESBL-producing isolates may confer resistance to tigecycline via other resistance mechanisms. Tigecycline resistance in some bacteria (e.g. *Acinetobacter calcoaceticus*-*Acinetobacter baumannii* complex) is associated with multi-drug resistant (MDR) efflux pumps.

Interaction with Other Antimicrobials

In vitro studies have not demonstrated antagonism between tigecycline and other commonly used antibacterials.

Antimicrobial Activity

Tigecycline has been shown to be active against most of the following microorganisms, both *in vitro* and in clinical infections [see *Indications and Usage (1)*].

Gram-positive bacteria

Enterococcus faecalis (vancomycin-susceptible isolates)

Staphylococcus aureus (methicillin-susceptible and -resistant isolates)

Streptococcus agalactiae

Streptococcus anginosus group (includes *S. anginosus*, *S. intermedius*, and *S. constellatus*) *Streptococcus pneumoniae* (penicillin-susceptible isolates)

Streptococcus pyogenes

Gram-negative bacteria

Citrobacter freundii

Enterobacter cloacae

Escherichia coli

Haemophilus influenzae

Klebsiella oxytoca

Klebsiella pneumoniae

Legionella pneumophila

Anaerobic bacteria

Bacteroides fragilis

Bacteroides thetaiotaomicron

Bacteroides uniformis

Bacteroides vulgatus

Clostridium perfringens

Peptostreptococcus micros

The following *in vitro* data are available, but their clinical significance is unknown. At least 90 percent of the following bacteria exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for tigecycline against isolates of similar genus or organism group.

However, the efficacy of tigecycline in treating clinical infections caused by these bacteria have not been established in adequate and well-controlled clinical trials.

Gram-positive bacteria

Enterococcus avium

Enterococcus casseliflavus

Enterococcus faecalis (vancomycin-resistant isolates)

Enterococcus faecium (vancomycin-susceptible and -resistant isolates)

Enterococcus gallinarum *Listeria monocytogenes*

Staphylococcus epidermidis (methicillin-susceptible and -resistant isolates)

Staphylococcus haemolyticus

Gram-negative bacteria

*Acinetobacter baumannii*¹

Aeromonas hydrophila

Citrobacter koseri *Enterobacter aerogenes*

Haemophilus influenzae (ampicillin-resistant)

Haemophilus parainfluenzae *Pasteurella multocida* *Serratia marcescens*

Stenotrophomonas maltophilia

Anaerobic bacteria

Bacteroides distasonis
Bacteroides ovatus
Peptostreptococcus spp.
Porphyromonas spp.
Prevotella spp.

Other bacteria

Mycobacterium abscessus
Mycobacterium fortuitum

¹ There have been reports of the development of tigecycline resistance in *Acinetobacter* infections seen during the course of standard treatment. Such resistance appears to be attributable to an MDR efflux pump mechanism. While monitoring for relapse of infection is important for all infected patients, more frequent monitoring in this case is suggested. If relapse is suspected, blood and other specimens should be obtained and cultured for the presence of bacteria. All bacterial isolates should be identified and tested for susceptibility to tigecycline and other appropriate antimicrobials.

Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria, and associated test methods and quality control standards recognized by FDA for this drug, please see <https://www.fda.gov/STIC>.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Lifetime studies in animals have not been performed to evaluate the carcinogenic potential of tigecycline. No mutagenic or clastogenic potential was found in a battery of tests, including *in vitro* chromosome aberration assay in Chinese hamster ovary (CHO) cells, *in vitro* forward mutation assay in CHO cells (HGRPT locus), *in vitro* forward mutation assays in mouse lymphoma cells, and *in vivo* mouse micronucleus assay.

Tigecycline did not affect mating or fertility in rats at exposures up to 5 times the human daily dose based on AUC (28 mcg·hr/mL at 12 mg/kg/day). In female rats, there were no compound-related effects on ovaries or estrous cycles at exposures up to 5 times the human daily dose based on AUC.

13.2 Animal Toxicology and/or Pharmacology

In two week studies, decreased erythrocytes, reticulocytes, leukocytes, and platelets, in association with bone marrow hypocellularity, have been seen with tigecycline at exposures of 8 times and 10 times the human daily dose based on AUC in rats and dogs, (AUC of approximately 50 and 60 mcg·hr/mL at doses of 30 and 12 mg/kg/day) respectively. These alterations were shown to be reversible after two weeks of dosing.

14 CLINICAL STUDIES

14.1 Complicated Skin and Skin Structure Infections

Tigecycline was evaluated in adults for the treatment of complicated skin and skin structure infections (cSSSI) in two randomized, double-blind, active-controlled, multinational, multicenter studies (Studies 1 and 2). These studies compared Tigecycline (100 mg intravenous initial dose followed by 50 mg every 12 hours) with vancomycin (1 g intravenous every 12 hours)/aztreonam (2 g intravenous every 12 hours) for 5 to 14 days. Patients with complicated deep soft tissue infections including wound infections and cellulitis (≥ 10 cm, requiring surgery/drainage or with complicated underlying disease), major abscesses, infected ulcers, and burns were enrolled in the studies. The primary efficacy endpoint was the clinical response at the test of cure (TOC) visit in the co-primary populations of the clinically

evaluable (CE) and clinical modified intent-to-treat (c-mITT) patients. See Table 4. Clinical cure rates at TOC by pathogen in the microbiologically evaluable patients are presented in Table 5.

Table 4. Clinical Cure Rates from Two Studies in Complicated Skin and Skin Structure Infections after 5 to 14 Days of Therapy

	Tigecycline* n/N (%)	Vancomycin/Aztreonam† n/N (%)
Study 1		
CE	165/199 (82.9)	163/198 (82.3)
c-mITT	209/277 (75.5)	200/260 (76.9)
Study 2		
CE	200/223 (89.7)	201/213 (94.4)
c-mITT	220/261 (84.3)	225/259 (86.9)

* 100 mg initially, followed by 50 mg every 12 hours

† Vancomycin (1 g every 12 hours)/Aztreonam (2 g every 12 hours)

Table 5. Clinical Cure Rates by Infecting Pathogen in Microbiologically Evaluable Patients with Complicated Skin and Skin Structure Infections*

Pathogen	Tigecycline n/N (%)	Vancomycin/Aztreonam n/N (%)
Escherichia coli	29/36 (80.6)	26/30 (86.7)
Enterobacter cloacae	10/12 (83.3)	15/15 (100)
Enterococcus faecalis (vancomycin-susceptible only)	15/21 (71.4)	19/24 (79.2)
Klebsiella pneumoniae	12/14 (85.7)	15/16 (93.8)
Methicillin-susceptible Staphylococcus aureus (MSSA)	124/137 (90.5)	113/120 (94.2)
Methicillin-resistant Staphylococcus aureus (MRSA)	79/95 (83.2)	46/57 (80.7)
Streptococcus agalactiae	8/8 (100)	11/14 (78.6)
Streptococcus anginosus grp.†	17/21 (81.0)	9/10 (90.0)
Streptococcus pyogenes	31/32 (96.9)	24/27 (88.9)
Bacteroides fragilis	7/9 (77.8)	4/5 (80.0)

* Two cSSSI pivotal studies and two Resistant Pathogen studies

† Includes *Streptococcus anginosus*, *Streptococcus intermedius*, and *Streptococcus constellatus*

14.2 Complicated Intra-abdominal Infections

Tigecycline was evaluated in adults for the treatment of complicated intra-abdominal infections (cIAI) in two randomized, double-blind, active-controlled, multinational, multicenter studies (Studies 1 and 2). These studies compared Tigecycline (100 mg intravenous initial dose followed by 50 mg every 12 hours) with imipenem/cilastatin (500 mg intravenous every 6 hours) for 5 to 14 days. Patients with

complicated diagnoses including appendicitis, cholecystitis, diverticulitis, gastric/duodenal perforation, intra-abdominal abscess, perforation of intestine, and peritonitis were enrolled in the studies. The primary efficacy endpoint was the clinical response at the TOC visit for the co-primary populations of the microbiologically evaluable (ME) and the microbiologic modified intent-to-treat (m-mITT) patients. See Table 6. Clinical cure rates at TOC by pathogen in the microbiologically evaluable patients are presented in Table 7.

Table 6. Clinical Cure Rates from Two Studies in Complicated Intra-abdominal Infections after 5 to 14 Days of Therapy

	Tigecycline* n/N (%)	Imipenem/Cilastatin† n/N (%)
Study 1		
ME	199/247 (80.6)	210/255 (82.4)
m-mITT	227/309 (73.5)	244/312 (78.2)
Study 2		
ME	242/265 (91.3)	232/258 (89.9)
m-mITT	279/322 (86.6)	270/319 (84.6)

* 100 mg initially, followed by 50 mg every 12 hours

† Imipenem/Cilastatin (500 mg every 6 hours)

Table 7. Clinical Cure Rates by Infecting Pathogen in Microbiologically Evaluable Patients with Complicated Intra-abdominal Infections *

Pathogen	Tigecycline n/N (%)	Imipenem/Cilastatin n/N (%)
Citrobacter freundii	12/16 (75.0)	3/4 (75.0)
Enterobacter cloacae	15/17 (88.2)	16/17 (94.1)
Escherichia coli	284/336 (84.5)	297/342 (86.8)
Klebsiella oxytoca	19/20 (95.0)	17/19 (89.5)
Klebsiella pneumoniae	42/47 (89.4)	46/53 (86.8)
Enterococcus faecalis	29/38 (76.3)	35/47 (74.5)
Methicillin-susceptible Staphylococcus aureus (MSSA)	26/28 (92.9)	22/24 (91.7)
Methicillin-resistant Staphylococcus aureus (MRSA)	16/18 (88.9)	1/3 (33.3)
Streptococcus anginosus grp.†	101/119 (84.9)	60/79 (75.9)
Bacteroides fragilis	68/88 (77.3)	59/73 (80.8)
Bacteroides thetaiotaomicron	36/41 (87.8)	31/36 (86.1)
Bacteroides uniformis	12/17 (70.6)	14/16 (87.5)
Bacteroides vulgatus	14/16 (87.5)	4/6 (66.7)
Clostridium perfringens	18/19 (94.7)	20/22 (90.9)
Peptostreptococcus	12/17 (70.6)	0/11 (0.0)

* Two cIAI pivotal studies and two Resistant Pathogen studies

† Includes *Streptococcus anginosus*, *Streptococcus intermedius*, and *Streptococcus constellatus*

14.3 Community-Acquired Bacterial Pneumonia

Tigecycline was evaluated in adults for the treatment of community-acquired bacterial pneumonia (CABP) in two randomized, double-blind, active-controlled, multinational, multicenter studies (Studies 1 and 2). These studies compared Tigecycline (100 mg intravenous initial dose followed by 50 mg every 12 hours) with levofloxacin (500 mg intravenous every 12 or 24 hours). In Study 1, after at least 3 days of intravenous therapy, a switch to oral levofloxacin (500 mg daily) was permitted for both treatment arms. Total therapy was 7 to 14 days. Patients with community-acquired bacterial pneumonia who required hospitalization and intravenous therapy were enrolled in the studies. The primary efficacy endpoint was the clinical response at the test of cure (TOC) visit in the co-primary populations of the clinically evaluable (CE) and clinical modified intent-to-treat (c-mITT) patients. See Table 8. Clinical cure rates at TOC by pathogen in the microbiologically evaluable patients are presented in Table 9.

Table 8. Clinical Cure Rates from Two Studies in Community-Acquired Bacterial Pneumonia after 7 to 14 Days of Total Therapy

	Tigecycline* n/N (%)	Levofloxacin† n/N (%)	95% CI‡
Study 1§			
CE	125/138 (90.6)	136/156(87.2)	(-4.4, 11.2)
c-mITT	149/191 (78)	158/203 (77.8)	(-8.5, 8.9)
Study 2			
CE	128/144 (88.9)	116/136 (85.3)	(-5.0, 12.2)
c-mITT	170/203 (83.7)	163/200 (81.5)	(-5.6, 10.1)

* 100 mg initially, followed by 50 mg every 12 hours

† Levofloxacin (500 mg intravenous every 12 or 24 hours)

‡ 95% confidence interval for the treatment difference

§ After at least 3 days of intravenous therapy, a switch to oral levofloxacin (500 mg daily) was permitted for both treatment arms in Study 1.

Table 9. Clinical Cure Rates by Infecting Pathogen in Microbiologically Evaluable Patients with Community-Acquired Bacterial Pneumonia*

Pathogen	Tigecycline n/N (%)	Levofloxacin n/N (%)
<i>Haemophilus influenzae</i>	14/17 (82.4)	13/16 (81.3)
<i>Legionella pneumophila</i>	10/10 (100.0)	6/6 (100.0)
<i>Streptococcus pneumoniae</i> (penicillin-susceptible only)†	44/46 (95.7)	39/44 (88.6)

* Two CABP studies

† Includes cases of concurrent bacteremia [cure rates of 20/22 (90.9%) versus 13/18 (72.2%) for Tigecycline and levofloxacin respectively]

To further evaluate the treatment effect of tigecycline, a post-hoc analysis was conducted in CABP patients with a higher risk of mortality, for whom the treatment effect of antibiotics is supported by

historical evidence. The higher-risk group included CABP patients from the two studies with any of the following factors:

- Age \geq 50 years
- PSI score \geq 3
- *Streptococcus pneumoniae* bacteremia

The results of this analysis are shown in Table 10. Age \geq 50 was the most common risk factor in the higher-risk group.

Table 10. Post-hoc Analysis of Clinical Cure Rates in Patients with Community-Acquired Bacterial Pneumonia Based on Risk of Mortality*

	Tigecycline n/N (%)	Levofloxacin n/N (%)	95% CI†
Study 1‡			
CE			
Higher risk			
Yes	93/103 (90.3)	84/102 (82.4)	(-2.3, 18.2)
No	32/35 (91.4)	52/54 (96.3)	(-20.8, 7.1)
c-mITT			
Higher risk			
Yes	111/142 (78.2)	100/134 (74.6)	(-6.9, 14)
No	38/49 (77.6)	58/69 (84.1)	(-22.8, 8.7)
Study 2			
CE			
Higher risk			
Yes	95/107 (88.8)	68/85 (80)	(-2.2, 20.3)
No	33/37 (89.2)	48/51 (94.1)	(-21.1, 8.6)
c-mITT			
Higher risk			
Yes	112/134 (83.6)	93/120 (77.5)	(-4.2, 16.4)
No	58/69 (84.1)	70/80 (87.5)	(-16.2, 8.8)

* Patients at higher risk of death include patients with any one of the following: \geq 50 year of age; PSI score \geq 3; or bacteremia due to *Streptococcus pneumoniae*

† 95% confidence interval for the treatment difference

‡ After at least 3 days of intravenous therapy, a switch to oral levofloxacin (500 mg daily) was permitted for both treatment arms in Study 1.

16 HOW SUPPLIED/STORAGE AND HANDLING

Tigecycline for Injection, USP, is supplied in a single-dose 10 mL glass vial, each containing 50 mg tigecycline lyophilized powder for reconstitution.

Supplied:

10 \times 50 mg Single Use Vials in one carton. NDC 70594-035-021

Prior to reconstitution, Tigecycline for Injection, USP, should be stored at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]. The reconstituted solution of Tigecycline for Injection, USP may be stored at room temperature (not to exceed 25°C/77°F) for up to 24 hours (up to 6 hours in the vial and the remaining time in the intravenous

bag) [See Dosage and administration(2.1)].

17 PATIENT COUNSELING INFORMATION

Tooth Discoloration and Inhibition of Bone Growth

Advise pregnant women that Tigecycline may cause permanent discoloration of deciduous teeth and reversible inhibition of bone growth when administered during the second and third trimesters of pregnancy [see Warnings and Precautions (5.6, 5.7) and Use in Specific Populations (8.1, 8.4)].

Lactation

Advise a woman not to breastfeed for longer than 3 weeks while taking Tigecycline because of the lack of data on effects due to prolonged breastfeeding, and the theoretical risk of dental discoloration and inhibition of bone growth. Women may also consider reducing infant exposure through pumping and discarding breastmilk during and for 9 days after the last dose of tigecycline [see Use in Specific Populations (8.2)].

Diarrhea

Advise patients, their families, or caregivers that diarrhea is a common problem caused by antibacterial drugs, including Tigecycline. Sometimes, frequent watery or bloody diarrhea may occur and may be a sign of a more serious intestinal infection. If severe watery or bloody diarrhea develops, advise patients to contact his or her healthcare provider [see Warnings and Precautions (5.8)].

Development of Resistance

Patients should be counseled that antibacterial drugs including Tigecycline should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Tigecycline is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Tigecycline or other antibacterial drugs in the future.

Manufactured for:

Xellia Pharmaceuticals USA, LLC Buffalo Grove IL 60089

PRINCIPAL DISPLAY PANEL - 50 mg Vial Carton

NDC 70594-035-02

Rx only

Tigecycline for Injection, USP

50 mg per vial

FOR INTRAVENOUS INFUSION

RECONSTITUTED SOLUTION

MUST BE FURTHER DILUTED

xellia

PHARMACEUTICALS

10 Single Use Vials

NDC 70594-035-02

Rx only

Tigecycline for Injection, USP



50 mg per vial



FOR INTRAVENOUS INFUSION
RECONSTITUTED SOLUTION
MUST BE FURTHER DILUTED 10 Single Use Vials

Unvarnished Area for
Lot, Exp, SN Code & GTIN

122622



101100370594035024



xellia PHARMACEUTICALS
RECONSTITUTED SOLUTION
MUST BE FURTHER DILUTED
10 Single Use Vials

50 mg per vial
FOR INTRAVENOUS INFUSION



Tigecycline for Injection, USP

Rx only

NDC 70594-035-02

Each vial contains 100 mg lactose monohydrate. The pH is adjusted with hydrochloric acid, and if necessary sodium hydroxide. Contains no preservatives. Single use vial. Discard unused portion.
Usual Dosage: The recommended dosage regimen for Tigecycline is an initial dose of 100 mg, followed by 50 mg every 12 hours.
Following reconstitution with 5.3 mL of 0.9% Sodium Chloride Injection, USP or 5 % Dextrose Injection, USP, the resultant solution will contain 10 mg/mL.

Reconstituted solution should be yellow to orange in color; if not, the solution should be discarded.

See package insert for complete directions for use.

Each vial contains 50 mg tigecycline lyophilized powder for intravenous infusion.

Prior to Reconstitution: Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]

Product inside sealed with flip off cap and aluminum seal.

Do not use if cap or aluminum seal is missing.

Manufactured for: Xellia Pharmaceuticals USA, LLC, Buffalo Grove, IL 60089


NDC 70594-035-02

Tigecycline for Injection, USP

50 mg per vial

FOR INTRAVENOUS INFUSION
RECONSTITUTED SOLUTION
MUST BE FURTHER DILUTED





TIGECYCLINE

tigecycline injection, powder, lyophilized, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70594-035
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TIGECYCLINE (UNII: 70JE2N95KR) (TIGECYCLINE - UNII:70JE2N95KR)	TIGECYCLINE	50 mg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70594-035-02	10 in 1 CARTON	02/01/2020	
1	NDC:70594-035-01	1 in 1 VIAL, GLASS; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205722	02/01/2020	

Labeler - Xellia Pharmaceuticals USA LLC (116768762)

Registrant - Xellia Pharmaceuticals ApS (305814345)

Revised: 3/2020

Xellia Pharmaceuticals USA LLC