

# CYANOCOBALAMIN- cyanocobalamin injection

## Somerset Therapeutics, LLC

### Cyanocobalamin Injection USP 1000 mcg/mL


#### DESCRIPTION

Cyanocobalamin Injection, USP is a sterile solution of cyanocobalamin for intramuscular or subcutaneous use.

Each mL contains 1,000 mcg cyanocobalamin; sodium chloride 0.9%; benzyl alcohol 1.5%; Water for Injection q.s.; Hydrochloric acid and/or sodium hydroxide for pH adjustment if necessary (4.5-7.0).

Cyanocobalamin appears as dark, red crystals or as an amorphous or crystalline, red powder. It is very hygroscopic in the anhydrous form, and sparingly soluble in water (1:80). It is stable to autoclaving for short periods at 121°C. The Vitamin B<sub>12</sub> coenzymes are very unstable in light.

The chemical name is 5,6-dimethyl-benzimidazolyl cyanocobamide. The cobalt content is 4.34%. The structural formula is represented below:



1355.38

#### CLINICAL PHARMACOLOGY

Vitamin B<sub>12</sub> is essential to growth, cell reproduction, hematopoiesis, nucleoprotein and myelin synthesis.

Cyanocobalamin is quantitatively and rapidly absorbed from intramuscular and subcutaneous sites of injection; the plasma level of the compound reaches its peak within one hour after intramuscular injection. Absorbed Vitamin B<sub>12</sub> is transported via specific B<sub>12</sub> binding proteins, transcobalamin I and II to the various tissues. The liver is the main organ for Vitamin B<sub>12</sub> storage.

Within 48 hours after injection of 100 or 1,000 mcg of Vitamin B<sub>12</sub>, 50 to 98% of the injected dose may appear in the urine. The major portion is excreted within the first eight hours. Intravenous administration results in even more rapid excretion with little opportunity for liver storage.

Gastrointestinal absorption of Vitamin B<sub>12</sub> depends on the presence of sufficient intrinsic

factor and calcium ions. Intrinsic factor deficiency causes pernicious anemia, which may be associated with subacute combined degeneration of the spinal cord. Prompt parenteral administration of Vitamin B<sub>12</sub> prevents progression of neurologic damage.

The average diet supplies about 5 to 15 mcg/day of Vitamin B<sub>12</sub> in a protein-bound form that is available for absorption after normal digestion. Vitamin B<sub>12</sub> is not present in foods of plant origin, but is abundant in foods of animal origin. In people with normal absorption, deficiencies have been reported only in strict vegetarians who consume no products of animal origin (including no milk products or eggs).

Vitamin B<sub>12</sub> is bound to intrinsic factor during transit through the stomach; separation occurs in the terminal ileum in the presence of calcium, and Vitamin B<sub>12</sub> enters the mucosal cell for absorption. It is then transported by the transcobalamin binding proteins. A small amount (approximately 1% of the total amount ingested) is absorbed by simple diffusion, but this mechanism is adequate only with very large doses. Oral absorption is considered too undependable to rely on in patients with pernicious anemia or other conditions resulting in malabsorption of Vitamin B<sub>12</sub>.

Cyanocobalamin is the most widely used form of Vitamin B<sub>12</sub>, and has hematopoietic activity apparently identical to that of the antianemia factor in purified liver extract. Hydroxocobalamin is equally as effective as cyanocobalamin, and they share the cobalamin molecular structure.

## **INDICATIONS AND USAGE**

Cyanocobalamin is indicated for Vitamin B<sub>12</sub> deficiencies due to malabsorption which may be associated with the following conditions:

Addisonian (pernicious) anemia

Gastrointestinal pathology, dysfunction, or surgery, including gluten enteropathy or sprue, small bowel bacterial overgrowth, total or partial gastrectomy

Fish tapeworm infestation

Malignancy of pancreas or bowel

Folic acid deficiency

It may be possible to treat the underlying disease by surgical correction of anatomic lesions leading to small bowel bacterial overgrowth, expulsion of fish tapeworm, discontinuation of drugs leading to vitamin malabsorption (see **Drug/Laboratory Test Interactions**), use of a gluten-free diet in nontropical sprue, or administration of antibiotics in tropical sprue. Such measures remove the need for long-term administration of cyanocobalamin.

Requirements of Vitamin B<sub>12</sub> in excess of normal (due to pregnancy, thyrotoxicosis, hemolytic anemia, hemorrhage, malignancy, hepatic and renal disease) can usually be met with oral supplementation.

Cyanocobalamin injection is also suitable for the Vitamin B<sub>12</sub> absorption test (Schilling test).

## **CONTRAINDICATIONS**

Sensitivity to cobalt and/or Vitamin B<sub>12</sub> is a contraindication.

## **WARNINGS**

This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

Patients with early Leber's disease (hereditary optic nerve atrophy) who were treated with cyanocobalamin suffered severe and swift optic atrophy.

Hypokalemia and sudden death may occur in severe megaloblastic anemia which is treated intensely.

Anaphylactic shock and death have been reported after parenteral Vitamin B<sub>12</sub> administration. An intradermal test dose is recommended before cyanocobalamin injection is administered to patients suspected of being sensitive to this drug.

**This product contains benzyl alcohol. Benzyl alcohol has been reported to be associated with a fatal "Gasping Syndrome" in premature infants.**

## **PRECAUTIONS**

### **General**

Vitamin B<sub>12</sub> deficiency that is allowed to progress for longer than three months may produce permanent degenerative lesions of the spinal cord. Doses of folic acid greater than 0.1 mg/day may result in hematologic remission in patients with Vitamin B<sub>12</sub> deficiency. Neurologic manifestations will not be prevented with folic acid, and if not treated with Vitamin B<sub>12</sub>, irreversible damage will result.

Doses of cyanocobalamin exceeding 10 mcg daily may produce hematologic response in patients with folate deficiency. Indiscriminate administration may mask the true diagnosis.

### **Information for Patients**

Patients with pernicious anemia should be instructed that they will require monthly injections of Vitamin B<sub>12</sub> for the remainder of their lives. Failure to do so will result in return of the anemia and in development of incapacitating and irreversible damage to the nerves of the spinal cord. Also, patients should be warned about the danger of taking folic acid in place of Vitamin B<sub>12</sub>, because the former may prevent anemia but allow progression of subacute combined degeneration.

A vegetarian diet which contains no animal products (including milk products or eggs) does not supply any Vitamin B<sub>12</sub>. Patients following such a diet should be advised to take oral Vitamin B<sub>12</sub> regularly. The need for Vitamin B<sub>12</sub> is increased by pregnancy and

lactation. Deficiency has been recognized in infants of vegetarian mothers who were breast fed, even though the mothers had no symptoms of deficiency at the time.

Contains no more than 57 mcg/L of aluminum.

### **Laboratory Tests**

During the initial treatment of patients with pernicious anemia, serum potassium must be observed closely the first 48 hours and potassium replaced if necessary.

Hematocrit, reticulocyte count, Vitamin B<sub>12</sub>, folate and iron levels should be obtained prior to treatment. Hematocrit and reticulocyte counts should be repeated daily from the 5<sup>th</sup> to 7<sup>th</sup> days of therapy and then frequently until the hematocrit is normal. If folate levels are low, folic acid should also be administered. If reticulocytes have not increased after treatment or if reticulocyte counts do not continue at least twice normal as long as the hematocrit is less than 35%, diagnosis or treatment should be reevaluated. Repeat determinations of iron and folic acid may reveal a complicating illness that might inhibit the response of the marrow.

Patients with pernicious anemia have about three times the incidence of carcinoma of the stomach as the general population, so appropriate tests for this condition should be carried out when indicated.

### **Drug and Laboratory Test Interactions**

Persons taking most antibiotics, methotrexate and pyrimethamine invalidate folic acid and Vitamin B<sub>12</sub> diagnostic blood assays.

Colchicine, para-aminosalicylic acid and heavy alcohol intake for longer than two weeks may produce malabsorption of Vitamin B<sub>12</sub>.

### **Carcinogenesis, Mutagenesis**

Long-term studies in animals to evaluate carcinogenic potential have not been done. There is no evidence from long-term use in patients with pernicious anemia that cyanocobalamin is carcinogenic. Pernicious anemia is associated with an increased incidence of carcinoma of the stomach, but this is believed to be related to the underlying pathology and not to treatment with cyanocobalamin.

### **Pregnancy**

Adequate and well-controlled studies have not been done in pregnant women. However, Vitamin B<sub>12</sub> is an essential vitamin and requirements are increased during pregnancy. Amounts of Vitamin B<sub>12</sub> that are recommended by the Food and Nutrition Board, National Academy of Science-National Research Council for pregnant women (4 mcg daily) should be consumed during pregnancy.

### **Nursing Mothers**

Vitamin B<sub>12</sub> is known to be excreted in human milk. Amounts of Vitamin B<sub>12</sub> that are recommended by the Food and Nutrition Board, National Academy of Science-National Research Council for lactating women (4 mcg daily) should be consumed during lactation.

## **Pediatric Use**

Intake in children should be in the amount (0.5 to 3 mcg daily) recommended by the Food and Nutrition Board, National Academy of Science-National Research Council.

## **ADVERSE REACTIONS**

### **Generalized**

Anaphylactic shock and death have been reported with administration of parenteral Vitamin B<sub>12</sub> (see **WARNINGS**).

### **Cardiovascular**

Pulmonary edema and congestive heart failure early in treatment; peripheral vascular thrombosis.

### **Hematological**

Polycythemia vera.

### **Gastrointestinal**

Mild transient diarrhea.

### **Dermatological**

Itching; transitory exanthema.

### **Miscellaneous**

Feeling of swelling of entire body.

## **OVERDOSAGE**

No overdosage has been reported with this drug.

## **DOSAGE AND ADMINISTRATION**

Avoid using the intravenous route. Use of this product intravenously will result in almost all of the vitamin being lost in the urine.

### **Pernicious Anemia**

Parenteral Vitamin B<sub>12</sub> is the recommended treatment and will be required for the remainder of the patient's life. The oral form is not dependable. A dose of 100 mcg daily for six or seven days should be administered by intramuscular or deep subcutaneous injection. If there is clinical improvement and if a reticulocyte response is observed, the same amount may be given on alternate days for seven doses, then every three to four days for another two to three weeks. By this time hematologic values should have become normal. This regimen should be followed by 100 mcg monthly for life. Folic acid should be administered concomitantly if needed.

### **Patients with Normal Intestinal Absorption**

Where the oral route is not deemed adequate, initial treatment similar to that for patients with pernicious anemia may be indicated depending on the severity of the deficiency. Chronic treatment should be with an oral B<sub>12</sub> preparation. If other vitamin deficiencies are present, they should be treated.

### **Schilling Test**

The flushing dose is 1,000 mcg.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

### **HOW SUPPLIED**

<b>NDC Number</b>	<b>Cyanocobalamin mcg/mL</b>	<b>Volume</b>
70069- <b>005</b> -10	1,000	1 mL
70069- <b>172</b> -10		10 mL
70069- <b>172</b> -25		

1 mL multiple dose vial, packaged 25 vials per tray.

10 mL multiple dose vial, packaged as;

- 10 vials per tray and

- 25 vials per tray.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

### **PROTECT FROM LIGHT**

Use only if solution is clear and seal intact.

For Product Inquiry call 1-800-417-9175

### **Manufactured for:**

Somerset Therapeutics, LLC

Hollywood, FL 33024

### **Made in India**

Code No.: KR/DRUGS/KTK/28/289/97

ST-CYC/P/05

Revised: 09/2020

### **Made in India**

Made in India

Neutral Code No: TN/DRUGS/616/1996

ST-CYC-MPPL/P/00

Revised: 10/2022

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

Cyanocobalamin Injection, USP

1 mL Container Label

NDC 70069-005-01

Contains no more than 57 mcg/L of aluminum. NDC 70069-005-01 Rx only  
Store at 20° to 25°C (68° to 77°F)  
[see USP Controlled Room Temperature].  
**PROTECT FROM LIGHT**  
**Manufactured for:**  
Somerset Therapeutics, LLC  
Hollywood, FL 33024  
**Made in India**  
Neutral Code No: TN/DRUGS/616/1996

ST-CYC-MPPL/L/00

**Cyanocobalamin Injection, USP**  
**1,000 mcg/mL**  
For IM or SC Use  
1 mL Multiple Dose Vial  
Contains Benzyl Alcohol as a Preservative

PSMV0005




Lot :  
Exp :

Keep area blank and varnish free for overprinting LOT and EXP  
7 x 10mm

Contains no more than 57 mcg/L of aluminum. NDC 70069-005-01 Rx only  
Store at 20° to 25°C (68° to 77°F)  
[see USP Controlled Room Temperature].  
**PROTECT FROM LIGHT**  
**Manufactured for:**  
Somerset Therapeutics, LLC  
Hollywood, FL 33024  
**Made in India**  
Code No.:KR/DRUGS/KTK/28/289/97

ST-CYC15/L/07

**Cyanocobalamin Injection, USP**  
**1,000 mcg/mL**  
For IM or SC Use  
1 mL Multiple Dose Vial  
Contains Benzyl Alcohol as a Preservative




Keep area blank and varnish free for overprinting LOT and EXP  
5 x 14mm

**Container Label**

Cyanocobalamin Injection, USP

1 mL Carton Label

NDC 70069-005-10




**Carton Label**

Cyanocobalamin Injection, USP

10 mL Container Label

NDC 70069-172-01

<p>Contains no more than 57 mcg/L of aluminum. Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. <b>PROTECT FROM LIGHT</b></p> <p><b>Manufactured for:</b> Somerset Therapeutics, LLC Hollywood, FL 33024</p> <p><b>Made in India</b> Code No.:KR/DRUGS/KTK/28/289/97</p>	<p>NDC 70069-172-01      Rx only</p> <p><b>Cyanocobalamin Injection, USP</b></p> <p><b>10,000 mcg/10 mL</b> (1,000 mcg/mL)</p> <p><b>For IM or SC Use</b></p> <p>10 mL Multiple Dose Vial</p> <p>Contains Benzyl Alcohol as a Preservative</p>		<p>No Varnish Area 24 x 7.5mm</p>
---	--	---	---------------------------------------

**container Label**

Cyanocobalamin Injection, USP

10 mL Carton Label – Pack of 25's

NDC 70069-172-25


			
<p>No Varnish Area 65 x 122mm</p>			
			
<p>For Intramuscular or Subcutaneous Use CONTAINS BENZYL ALCOHOL AS A PRESERVATIVE (1,000 mcg/mL) <b>10,000 mcg/10 mL</b> <b>Cyanocobalamin Injection, USP</b> 25 x 10 mL Multiple Dose Vials NDC 70069-172-25 Rx only</p>			
<p><b>Each mL contains:</b> Cyanocobalamin 1,000 mcg; Benzyl alcohol 1.5%; Sodium chloride 0.9%; Water for injection q.s. pH adjusted with hydrochloric acid or sodium hydroxide if necessary. Contains no more than 57 mcg/L of Aluminum. Not made with natural rubber latex.</p> <p><b>Usual dosage:</b> See package insert. Use only if solution is clear and seal intact.</p> <p>Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. <b>PROTECT FROM LIGHT</b></p>	<p><b>Manufactured for:</b> Somerset Therapeutics, LLC Hollywood, FL 33024 Customer Care # 1-800-417-9175</p> <p><b>Made in India</b> Code No.:KR/DRUGS/KTK/28/289/97 ST-CYC12/C25/03</p>	<p>Rx only</p> <p>NDC 70069-172-25 25 x 10 mL Multiple Dose Vials</p> <p><b>Cyanocobalamin Injection, USP</b></p> <p><b>10,000 mcg/10 mL</b> (1,000 mcg/mL)</p> <p><b>For Intramuscular or Subcutaneous Use</b> CONTAINS BENZYL ALCOHOL AS A PRESERVATIVE</p> 	

**carton label**

Cyanocobalamin Injection, USP

10 mL Carton Label – Pack of 10's

NDC 70069-172-10



carton-label

# CYANOCOBALAMIN

cyanocobalamin injection

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:70069-005
<b>Route of Administration</b>	INTRAMUSCULAR, SUBCUTANEOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CYANOCOBALAMIN</b> (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	1000 ug in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	15 mg in 1 mL
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	9 mg in 1 mL
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Product Characteristics

<b>Color</b>	RED (Red Colored Clear Solution)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

		<b>Marketing Start</b>	<b>Marketing End</b>
--	--	------------------------	----------------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70069-005-10	25 in 1 CARTON	12/11/2015	
1	NDC:70069-005-01	1 mL in 1 VIAL; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206503	12/11/2015	

## CYANOCOBALAMIN

cyanocobalamin injection

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70069-172
Route of Administration	INTRAMUSCULAR, SUBCUTANEOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CYANOCOBALAMIN (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	1000 ug in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	15 mg in 1 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

### Product Characteristics

Color	RED (Red Colored Clear Solution)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:70069-172			

1	NDC:70069-172-10	10 in 1 CARTON	06/27/2017	
1	NDC:70069-172-01	10 mL in 1 VIAL; Type 0: Not a Combination Product		
2	NDC:70069-172-25	25 in 1 CARTON	06/27/2017	
2	NDC:70069-172-01	10 mL in 1 VIAL; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206503	06/27/2017	

**Labeler** - Somerset Therapeutics, LLC (079947873)

**Registrant** - Somerset Therapeutics, LLC (079947873)

## Establishment

Name	Address	ID/FEI	Business Operations
Somerset Therapeutics Limited		677236695	ANALYSIS(70069-005, 70069-172) , LABEL(70069-005, 70069-172) , MANUFACTURE(70069-005, 70069-172) , PACK(70069-005, 70069-172)

## Establishment

Name	Address	ID/FEI	Business Operations
Maiva Pharma Private Limited		725656438	ANALYSIS(70069-005) , LABEL(70069-005) , MANUFACTURE(70069-005) , PACK(70069-005)

Revised: 12/2023

Somerset Therapeutics, LLC