

SENTRY HC WORM X DS- pyrantel pamoate liquid

Sergeant's Pet Care Products, LLC

Sentry HC Worm X DS


For Dogs (pyrantel pamoate)

Canine Anthelmintic Suspension

Approved by FDA under ANADA # 200-007

WormX DS is a suspension of pyrantel pamoate in a palatable butterscotch flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below.



(E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-thienyl] vinyl] pyrimidine 4,4' methylenebis [3-hydroxy-2-naphthoate](1:1)

INDICATIONS FOR USE:

WormX DS suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxocaris leonina*), and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.

WormX DS suspension may also be used to prevent reinfestation of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

DOSAGE AND ADMINISTRATION:

Administer one full teaspoon (5 mL) for each 10 pounds of body weight (2.27 mg base per lb. of body weight). Although most dogs have been observed to find this formulation very palatable and willingly consume it undiluted, it may be necessary to mix a small quantity of formulation in the dog's normal ration to encourage consumption. Fasting prior to and after treatment is not necessary.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated

at monthly intervals to prevent *T. canis* reinfestation.

CAUTION: THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, **IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.**

RECOMMENDED STORAGE: Store below 86°F (30°C).

EFFICACY

Critical (worm count) studies in dogs demonstrated that pyrantel pamoate at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

SAFETY

One of the most outstanding and significant features of pyrantel pamoate is its wide margin of therapeutic safety in dogs. The acute oral LD₅₀ of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per pound of body weight, which indicates a therapeutic index in excess of 138 X the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40 x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches and males at stud. Additional data have demonstrated the safe use of pyrantel pamoate in (1) dogs having heartworm infections and/or receiving medication for heartworms, (2) dogs exposed to organophosphate flea collars or flea/tick dip treatments, and (3) dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.

If you have any questions or comments about this product, please write: SENTRY Consumer Response, PO Box 540399, Omaha, NE 68154-0399.

Distributed by:

Sergeant's Pet Care Products, Inc., Omaha, NE 68138

PRINCIPAL DISPLAY PANEL - 60 mL Bottle

INDICATIONS FOR USE

To prevent reinfestation of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping. For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*), and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

WARNING: Keep Out Of Reach of Children

Read Accompanying Professional Information

RECOMMENDED STORAGE: STORE BELOW 30° C (86°F)

Distributed by: Sergeant's Pet Care Products, LLC,
10077 S. 134th Street, Omaha, NE 68138

**DO NOT PRINT
UNCOATED AREA**



OBSERVE LABEL
DIRECTIONS
P0112535 AMIA

SENTRY^{HC}
WORM^XDS

(pyrantel pamoate)

FOR DOGS

**Canine Anthelmintic
Suspension**

Active Ingredient:

4.54 mg of pyrantel base
as pyrantel pamoate per mL

FOR ANIMAL USE ONLY

Approved by FDA under ANADA #200-007

NET CONTENTS: 2 fl oz (60 mL)

IMPORTANT: SHAKE WELL BEFORE USING

DIRECTIONS FOR USE: For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation. Administer one full teaspoonful (5 mL) for each 10 lb. of body weight.

For the removal of large roundworms (*ascaris*) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lb. of body weight. It is not necessary to withhold food prior to treatment. If medication is to be dispensed, client can be advised that dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is a reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.

The presence of large roundworms (*ascaris*) and hookworms should be confirmed by lab fecal examination. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

PRINCIPAL DISPLAY PANEL - 60 mL Carton

SENTRY^{HC}
WORM X^{DS}
(pyrantel pamoate)

SENTRY^{HC}
WORM X^{DS}
(pyrantel pamoate)

INDICATIONS FOR USE: To prevent reinfection of foxian cecis in puppies and adult dogs and in lactating bitches after whelping.

For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*), and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.

The presence of these parasites should be confirmed by laboratory fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

In case of an Emergency, call: 1-800-781-4738
 Non-Emergency, call: 1-800-224-PETS (7387)

If you have questions or comments about this product, please write:
 SENTRY Consumer Response, PO Box 540399, Omaha, NE 68154-0399

Distributed by: Sergeant's Pet Care Products, LLC, 10077 S. 134th Street, Omaha, NE 68138

IMPORTANT: SHAKE WELL BEFORE USE
WARNING: KEEP OUT OF THE REACH OF CHILDREN

DO NOT PRINT
 UNCOATED AREA



PO 1015 - AVMA

SENTRY^{HC}
WORM X^{DS}
(pyrantel pamoate)
FOR DOGS



Canine Anthelmintic Suspension

- Removes Large Roundworms & Hookworms in puppies & dogs
- Treats dogs up to 120 lbs.

Active Ingredient: 4.54 mg of pyrantel base as pyrantel pamoate per mL
FOR ANIMAL USE ONLY
 See back panel for complete use directions.
 Approved by FDA under ANADA #200-007
NET CONTENTS: 2 fl oz (60 mL)

SENTRY^{HC}
WORM X^{DS}
(pyrantel pamoate)

SENTRY^{HC}
WORM X^{DS}
(pyrantel pamoate)

Canine Anthelmintic Suspension

DIRECTIONS FOR USE: For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent fecal reinfection. Administer one full teaspoonful (5 mL) for each 10 lb of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one 1/8 teaspoonful (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to treatment.

If medication is to be dispensed, client can be advised that dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is a reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.

It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

Recommended Storage: Store below 30° C (86° F)
 Read Accompanying Professional Information

SENTRY HC WORM X DS

pyrantel pamoate liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:21091-500
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Pyrantel Pamoate (UNII: 81BK194Z5M) (Pyrantel - UNII:4QIH0N49E7)	Pyrantel	4.54 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN SODIUM (UNII: CR6K9C2NHK)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BUTTERSCOTCH	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21091-500-02	1 in 1 CARTON		
1		60 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200007	06/14/2023	

Labeler - Sergeant's Pet Care Products, LLC (876995171)

Registrant - Happy Jack (003189354)

Establishment

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	manufacture, label, pack

Establishment

Name	Address	ID/FEI	Business Operations
Cosma Spa		428655732	api manufacture

Revised: 11/2023

Sergeant's Pet Care Products, LLC