

DEXTROAMPHETAMINE SULFATE - dextroamphetamine sulfate capsule, extended release
Physicians Total Care, Inc.

DEXTROAMPHETAMINE SULFATE EXTENDED-RELEASE CAPSULES - CII

Revised SEPTEMBER 2007

11001275

Rx only


WARNING:

AMPHETAMINES HAVE A HIGH POTENTIAL FOR ABUSE. ADMINISTRATION OF AMPHETAMINES FOR PROLONGED PERIODS OF TIME MAY LEAD TO DRUG DEPENDENCE AND MUST BE AVOIDED. PARTICULAR ATTENTION SHOULD BE PAID TO THE POSSIBILITY OF SUBJECTS OBTAINING AMPHETAMINES FOR NON-THERAPEUTIC USE OR DISTRIBUTION TO OTHERS, AND THE DRUGS SHOULD BE PRESCRIBED OR DISPENSED SPARINGLY.

MISUSE OF AMPHETAMINES MAY CAUSE SUDDEN DEATH AND SERIOUS CARDIOVASCULAR ADVERSE EVENTS.

DESCRIPTION:

Dextroamphetamine sulfate is the dextro isomer of the compound *d,l*-amphetamine sulfate, a sympathomimetic amine of the amphetamine group. Chemically, dextroamphetamine is *d*-alpha-methylphenethylamine, and is present in all forms of dextroamphetamine as the neutral sulfate. The structural formula is as follows:



Each extended-release capsule is so prepared that an initial dose is released promptly and the remaining medication is released gradually over a prolonged period.

Each capsule contains dextroamphetamine sulfate, and has the following inactive ingredients: colloidal silicon dioxide, dibutyl sebacate, ethylcellulose aqueous dispersion, methylcellulose, povidone, propylene glycol, sugar spheres and talc.

The capsule shell ingredients in the 5 mg are D&C red no. 33, FD&C blue no. 1, FD&C yellow no. 6, gelatin, and titanium dioxide.

The capsule shell ingredients in the 10 mg are black iron oxide, gelatin, red iron oxide, titanium dioxide, and yellow iron oxide.

The capsule shell ingredients in the 15 mg are black iron oxide, gelatin, red iron oxide, titanium

dioxide, and yellow iron oxide.

The imprinting ingredients are D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, pharmaceutical glaze, propylene glycol, and synthetic black iron oxide.

CLINICAL PHARMACOLOGY:

Amphetamines are non-catecholamine, sympathomimetic amines with CNS stimulant activity. Peripheral actions include elevations of systolic and diastolic blood pressures and weak bronchodilator and respiratory stimulant action.

There is neither specific evidence which clearly establishes the mechanism whereby amphetamines produce mental and behavioral effects in children, nor conclusive evidence regarding how these effects relate to the condition of the central nervous system.

Dextroamphetamine sulfate extended-release capsules are formulated to release the active drug substance *in vivo* in a more gradual fashion than the standard formulation, as demonstrated by blood levels. The formulation has not been shown superior in effectiveness over the same dosage of the standard, non-controlled-release formulations given in divided doses.

Pharmacokinetics:

The pharmacokinetics of the tablet and extended-release capsule was compared in 12 healthy subjects. The extent of bioavailability of the extended-release capsule was similar compared to the immediate release tablet. Following administration of three 5 mg tablets, average maximal dextroamphetamine plasma concentrations (C_{max}) of 36.6 ng/mL were achieved at approximately 3 hours. Following administration of one 15 mg extended-release capsule, maximal dextroamphetamine plasma concentrations were obtained approximately 8 hours after dosing. The average C_{max} was 23.5 ng/mL. The average plasma $T_{1/2}$ was similar for both the tablet and extended-release capsule and was approximately 12 hours.

In 12 healthy subjects, the rate and extent of dextroamphetamine absorption were similar following administration of the extended-release capsule formulation in the fed (58 to 75 gm fat) and fasted state.

INDICATIONS AND USAGE:

Dextroamphetamine sulfate is indicated:

1. In Narcolepsy.

2. In Attention Deficit Disorder with Hyperactivity, as an integral part of a total treatment program which typically includes other remedial measures (psychological, educational, social) for a stabilizing effect in pediatric patients (ages 3 years to 16 years) with a behavioral syndrome characterized by the following group of developmentally inappropriate symptoms: moderate to severe distractibility, short attention span, hyperactivity, emotional lability, and impulsivity. The diagnosis of this syndrome should not be made with finality when these symptoms are only of comparatively recent origin. Nonlocalizing (soft) neurological signs, learning disability, and abnormal EEG may or may not be present, and a diagnosis of central nervous system dysfunction may or may not be warranted.

CONTRAINDICATIONS:

Advanced arteriosclerosis, symptomatic cardiovascular disease, moderate to severe hypertension, hyperthyroidism, known hypersensitivity or idiosyncrasy to the sympathomimetic amines, glaucoma.

Agitated states.

Patients with a history of drug abuse.

During or within 14 days following the administration of monoamine oxidase inhibitors (hypertensive crises may result).

WARNINGS:

Serious Cardiovascular Events

Sudden Death in Patients with Pre-existing Structural Cardiac Abnormalities or Other Serious Heart Problems:

Children and Adolescents: Sudden death has been reported in association with CNS stimulant treatment at usual doses in children and adolescents with structural cardiac abnormalities or other serious heart problems. Although some serious heart problems alone carry an increased risk of sudden death, stimulant products generally should not be used in children or adolescents with known serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, or other serious cardiac problems that may place them at increased vulnerability to the sympathomimetic effects of a stimulant drug.

Adults: Sudden deaths, stroke, and myocardial infarction have been reported in adults taking stimulant drugs at usual doses for ADHD. Although the role of stimulants in these adult cases is also unknown, adults have a greater likelihood than children of having serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, coronary artery disease, or other serious cardiac problems. Adults with such abnormalities should also generally not be treated with stimulant drugs (see CONTRAINDICATIONS).

Hypertension and Other Cardiovascular Conditions:

Stimulant medications cause a modest increase in average blood pressure (about 2-4 mmHg) and average heart rate (about 3-6 bpm), and individuals may have larger increases. While the mean changes alone would not be expected to have short-term consequences, all patients should be monitored for larger changes in heart rate and blood pressure. Caution is indicated in treating patients whose underlying medical conditions might be compromised by increases in blood pressure or heart rate, e.g., those with pre-existing hypertension, heart failure, recent myocardial infarction, or ventricular arrhythmia (see CONTRAINDICATIONS).

Assessing Cardiovascular Status in Patients Being Treated With Stimulant Medications:

Children, adolescents, or adults who are being considered for treatment with stimulant medications should have a careful history (including assessment for a family history of sudden death or ventricular arrhythmia) and physical exam to assess for the presence of cardiac disease, and should receive further cardiac evaluation if findings suggest such disease (e.g., electrocardiogram and echocardiogram). Patients who develop symptoms such as exertional chest pain, unexplained syncope, or other symptoms suggestive of cardiac disease during stimulant treatment should undergo a prompt cardiac evaluation.

Psychiatric Adverse Events

Pre-Existing Psychosis:

Administration of stimulants may exacerbate symptoms of behavior disturbance and thought disorder in patients with a pre-existing psychotic disorder.

Bipolar Illness:

Particular care should be taken in using stimulants to treat ADHD in patients with comorbid bipolar disorder because of concern for possible induction of a mixed/manic episode in such patients. Prior to initiating treatment with a stimulant, patients with comorbid depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression.

Emergence of New Psychotic or Manic Symptoms:

Treatment emergent psychotic or manic symptoms, e.g., hallucinations, delusional thinking, or mania in children and adolescents without a prior history of psychotic illness or mania can be caused by stimulants at usual doses. If such symptoms occur, consideration should be given to a possible causal role of the stimulant, and discontinuation of treatment may be appropriate. In a pooled analysis of multiple short-term, placebo-controlled studies, such symptoms occurred in about 0.1% (4 patients with events out of 3,482 exposed to methylphenidate or amphetamine for several weeks at usual doses) of stimulant-treated patients compared to 0 in placebo-treated patients.

Aggression:

Aggressive behavior or hostility is often observed in children and adolescents with ADHD, and has been reported in clinical trials and the postmarketing experience of some medications indicated for the treatment of ADHD. Although there is no systematic evidence that stimulants cause aggressive behavior or hostility, patients beginning treatment for ADHD should be monitored for the appearance of, or worsening of, aggressive behavior or hostility.

Long-Term Suppression of Growth:

Careful follow-up of weight and height in children ages 7 to 10 years who were randomized to either methylphenidate or non-medication treatment groups over 14 months, as well as in naturalistic subgroups of newly methylphenidate-treated and non-medication treated children over 36 months (to the ages of 10 to 13 years), suggests that consistently medicated children (i.e., treatment for 7 days per week throughout the year) have a temporary slowing in growth rate (on average, a total of about 2 cm less growth in height and 2.7 kg less growth in weight over 3 years), without evidence of growth rebound during this period of development. Published data are inadequate to determine whether chronic use of amphetamines may cause a similar suppression of growth, however, it is anticipated that they likely have this effect as well. Therefore, growth should be monitored during treatment with stimulants, and patients who are not growing or gaining height or weight as expected may need to have their treatment interrupted.

Seizures:

There is some clinical evidence that stimulants may lower the convulsive threshold in patients with prior history of seizures, in patients with prior EEG abnormalities in absence of seizures, and, very rarely, in patients without a history of seizures and no prior EEG evidence of seizures. In the presence of seizures, the drug should be discontinued.

Visual Disturbance:

Difficulties with accommodation and blurring of vision have been reported with stimulant treatment.

PRECAUTIONS:

General:

The least amount feasible should be prescribed or dispensed at one time in order to minimize the possibility of overdose.

Information for Patients:

Amphetamines may impair the ability of the patient to engage in potentially hazardous activities such as operating machinery or vehicles; the patient should therefore be cautioned accordingly.

Prescribers or other health professionals should inform patients, their families, and their caregivers about the benefits and risks associated with treatment with dextroamphetamine and should counsel them in its appropriate use. A patient Medication Guide is available for Dextroamphetamine Sulfate Extended-Release Capsules.

The prescriber or health professional should instruct patients, their families, and their caregivers to read the Medication Guide and should assist them in understanding its contents. Patients should be given the opportunity to discuss the contents of the Medication Guide and to obtain answers to any questions they may have. The complete text of the Medication Guide is reprinted at the end of this document.

Drug Interactions:

Acidifying agents:

Gastrointestinal acidifying agents (guanethidine, reserpine, glutamic acid HCl, ascorbic acid, fruit juices, etc.) lower absorption of amphetamines. Urinary acidifying agents (ammonium chloride, sodium acid phosphate, etc.) increase the concentration of the ionized species of the amphetamine molecule, thereby increasing urinary excretion. Both groups of agents lower blood levels and efficacy of amphetamines.

Adrenergic blockers:

Adrenergic blockers are inhibited by amphetamines.

Alkalinizing agents:

Gastrointestinal alkalinizing agents (sodium bicarbonate, etc.) increase absorption of amphetamines. Urinary alkalinizing agents (acetazolamide, some thiazides) increase the concentration of the non-ionized species of the amphetamine molecule, thereby decreasing urinary excretion. Both groups of agents increase blood levels and therefore potentiate the actions of amphetamines.

Antidepressants, tricyclic:

Amphetamines may enhance the activity of tricyclic or sympathomimetic agents; d-amphetamine with desipramine or protriptyline and possibly other tricyclics cause striking and sustained increases in the concentration of d-amphetamine in the brain; cardiovascular effects can be potentiated.

MAO inhibitors:

MAOI antidepressants, as well as a metabolite of furazolidone, slow amphetamine metabolism. This slowing potentiates amphetamines, increasing their effect on the release of norepinephrine and other monoamines from adrenergic nerve endings; this can cause headaches and other signs of hypertensive crisis. A variety of neurological toxic effects and malignant hyperpyrexia can occur, sometimes with fatal results.

Antihistamines:

Amphetamines may counteract the sedative effect of antihistamines.

Antihypertensives:

Amphetamines may antagonize the hypotensive effects of antihypertensives.

Chlorpromazine:

Chlorpromazine blocks dopamine and norepinephrine reuptake, thus inhibiting the central stimulant effects of amphetamines, and can be used to treat amphetamine poisoning.

Ethosuximide:

Amphetamines may delay intestinal absorption of ethosuximide.

Haloperidol:

Haloperidol blocks dopamine and norepinephrine reuptake, thus inhibiting the central stimulant effects

of amphetamines.

Lithium carbonate:

The stimulatory effects of amphetamines may be inhibited by lithium carbonate.

Meperidine:

Amphetamines potentiate the analgesic effect of meperidine.

Methenamine therapy:

Urinary excretion of amphetamines is increased, and efficacy is reduced, by acidifying agents used in methenamine therapy.

Norepinephrine:

Amphetamines enhance the adrenergic effect of norepinephrine.

Phenobarbital:

Amphetamines may delay intestinal absorption of phenobarbital; co-administration of phenobarbital may produce a synergistic anticonvulsant action.

Phenytoin:

Amphetamines may delay intestinal absorption of phenytoin; co-administration of phenytoin may produce a synergistic anticonvulsant action.

Propoxyphene:

In cases of propoxyphene overdose, amphetamine CNS stimulation is potentiated and fatal convulsions can occur.

Veratrum Alkaloids:

Amphetamines inhibit the hypotensive effect of veratrum alkaloids.

Drug/Laboratory Test Interactions:

- Amphetamines can cause a significant elevation in plasma corticosteroid levels. This increase is greatest in the evening.
- Amphetamines may interfere with urinary steroid determinations.

Carcinogenesis/Mutagenesis:

Mutagenicity studies and long-term studies in animals to determine the carcinogenic potential of dextroamphetamine sulfate have not been performed.

Pregnancy-Teratogenic Effects:

Pregnancy Category C:

Dextroamphetamine sulfate has been shown to have embryotoxic and teratogenic effects when administered to A/Jax mice and C57BL mice in doses approximately 41 times the maximum human dose. Embryotoxic effects were not seen in New Zealand white rabbits given the drug in doses 7 times the human dose nor in rats given 12.5 times the maximum human dose. While there are no adequate and well-controlled studies in pregnant women, there has been one report of severe congenital bony deformity, tracheoesophageal fistula, and anal atresia (Vater association) in a baby born to a woman who took dextroamphetamine sulfate with lovastatin during the first trimester of pregnancy. Dextroamphetamine sulfate should be used during pregnancy only if the potential benefit justifies the potential risk to the

fetus.

Nonteratogenic Effects:

Infants born to mothers dependent on amphetamines have an increased risk of premature delivery and low birth weight. Also, these infants may experience symptoms of withdrawal as demonstrated by dysphoria, including agitation, and significant lassitude.

Nursing Mothers:

Amphetamines are excreted in human milk. Mothers taking amphetamines should be advised to refrain from nursing.

Pediatric Use:

Long-term effects of amphetamines in pediatric patients have not been well established.

Amphetamines are not recommended for use in pediatric patients under 3 years of age with Attention Deficit Disorder with Hyperactivity described under **INDICATIONS AND USAGE**.

Clinical experience suggests that in psychotic children, administration of amphetamines may exacerbate symptoms of behavior disturbance and thought disorder.

Amphetamines have been reported to exacerbate motor and phonic tics and Tourette's syndrome. Therefore, clinical evaluation for tics and Tourette's syndrome in children and their families should precede use of stimulant medications.

Data are inadequate to determine whether chronic administration of amphetamines may be associated with growth inhibition; therefore, growth should be monitored during treatment.

Drug treatment is not indicated in all cases of Attention Deficit Disorder with Hyperactivity and should be considered only in light of the complete history and evaluation of the child. The decision to prescribe amphetamines should depend on the physician's assessment of the chronicity and severity of the child's symptoms and their appropriateness for his/her age. Prescription should not depend solely on the presence of one or more of the behavioral characteristics.

When these symptoms are associated with acute stress reactions, treatment with amphetamines is usually not indicated.

ADVERSE REACTIONS:

Cardiovascular:

Palpitations, tachycardia, elevation of blood pressure. There have been isolated reports of cardiomyopathy associated with chronic amphetamine use.

Central Nervous System:

Psychotic episodes at recommended doses (rare), overstimulation, restlessness, dizziness, insomnia, euphoria, dyskinesia, dysphoria, tremor, headache, exacerbation of motor and phonic tics and Tourette's syndrome.

Gastrointestinal:

Dryness of the mouth, unpleasant taste, diarrhea, constipation, other gastrointestinal disturbances. Anorexia and weight loss may occur as undesirable effects.

Allergic:

Urticaria.

Endocrine:

Impotence, changes in libido.

DRUG ABUSE AND DEPENDENCE:

Dextroamphetamine sulfate is a Schedule II controlled substance.

Amphetamines have been extensively abused. Tolerance, extreme psychological dependence and severe social disability have occurred. There are reports of patients who have increased the dosage to many times that recommended. Abrupt cessation following prolonged high dosage administration results in extreme fatigue and mental depression; changes are also noted on the sleep EEG.

Manifestations of chronic intoxication with amphetamines include severe dermatoses, marked insomnia, irritability, hyperactivity and personality changes. The most severe manifestation of chronic intoxication is psychosis, often clinically indistinguishable from schizophrenia. This is rare with oral amphetamines.

OVERDOSAGE:

Individual patient response to amphetamines varies widely. While toxic symptoms occasionally occur as an idiosyncrasy at doses as low as 2 mg, they are rare with doses of less than 15 mg; 30 mg can produce severe reactions, yet doses of 400 to 500 mg are not necessarily fatal.

In rats, the oral LD₅₀ of dextroamphetamine sulfate is 96.8 mg/kg.

Manifestations of acute overdose with amphetamines include restlessness, tremor, hyperreflexia, rhabdomyolysis, rapid respiration, hyperpyrexia, confusion, assaultiveness, hallucinations, panic states.

Fatigue and depression usually follow the central stimulation.

Cardiovascular effects include arrhythmias, hypertension or hypotension and circulatory collapse. Gastrointestinal symptoms include nausea, vomiting, diarrhea and abdominal cramps. Fatal poisoning is usually preceded by convulsions and coma.

Treatment:

Consult with a Certified Poison Control Center for up-to-date guidance and advice. Management of acute amphetamine intoxication is largely symptomatic and includes gastric lavage, administration of activated charcoal, administration of a cathartic, and sedation. Experience with hemodialysis or peritoneal dialysis is inadequate to permit recommendation in this regard. Acidification of the urine increases amphetamine excretion, but is believed to increase risk of acute renal failure if myoglobinuria is present. If acute, severe hypertension complicates amphetamine overdose, administration of intravenous phentolamine (Regitine[®], CIBA) has been suggested. However, a gradual drop in blood pressure will usually result when sufficient sedation has been achieved.

Chlorpromazine antagonizes the central stimulant effects of amphetamines and can be used to treat amphetamine intoxication.

Since much of the extended-release capsule medication is coated for gradual release, therapy directed at reversing the effects of the ingested drug and at supporting the patient should be continued for as long as overdose symptoms remain. Saline cathartics are useful for hastening the evacuation of pellets that have not already released medication.

DOSAGE AND ADMINISTRATION:

Amphetamines should be administered at the lowest effective dosage and dosage should be individually adjusted. Late evening doses - particularly with the extended-release capsule form - should be avoided because of the resulting insomnia.

Narcolepsy:

Usual dose 5 to 60 mg per day in divided doses, depending on the individual patient response. Narcolepsy seldom occurs in children under 12 years of age; however, when it does, dextroamphetamine sulfate may be used. The suggested initial dose for patients aged 6 to 12 is 5 mg daily; daily dose may be raised in increments of 5 mg at weekly intervals until optimal response is obtained. In patients 12 years of age and older, start with 10 mg daily; daily dosage may be raised in increments of 10 mg at weekly intervals until optimal response is obtained. If bothersome adverse reactions appear (e.g., insomnia or anorexia), dosage should be reduced. Extended-release capsules may be used for once-a-day dosage wherever appropriate.

Attention Deficit Disorder with Hyperactivity:

Not recommended for pediatric patients under 3 years of age.

In pediatric patients 6 years of age and older, start with 5 mg once or twice daily; daily dosage may be raised in increments of 5 mg at weekly intervals until optimal response is obtained. Only in rare cases will it be necessary to exceed a total of 40 mg per day.

Extended-release capsules may be used for once-a-day dosage wherever appropriate.

Where possible, drug administration should be interrupted occasionally to determine if there is a recurrence of behavioral symptoms sufficient to require continued therapy.

HOW SUPPLIED:

Dextroamphetamine Sulfate Extended-Release Capsules are available as:

10 mg: Brown opaque cap and colorless, clear body filled with white to off-white pellets. Imprinted in black ink **barr** 955. Available in bottles of:

Bottles of	NDC 54868-
20	5479-1

Bottles of	NDC 54868-
60	5479-0

15 mg: Dark brown opaque cap and colorless, clear body filled with white to off-white pellets. Imprinted in black ink **barr** 956. Available in bottles of:

Bottles of	NDC 54868-
10	5388-0

Bottles of	NDC 54868-
30	5388-1

Dispense in a tight, light-resistant container.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

MEDICATION GUIDE

DEXTROAMPHETAMINE SULFATE EXTENDED-RELEASE CAPSULES - CII

Read the Medication Guide that comes with Dextroamphetamine Sulfate Extended-Release Capsules before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about you or your child's treatment with Dextroamphetamine Sulfate Extended-Release Capsules.

What is the most important information I should know about Dextroamphetamine Sulfate Extended-Release Capsules?

The following have been reported with use of Dextroamphetamine Sulfate Extended-Release Capsules and other stimulant medicines.

1. Heart-related problems:

- **sudden death in patients who have heart problems or heart defects**
- **stroke and heart attack in adults**
- **increased blood pressure and heart rate**

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting Dextroamphetamine Sulfate Extended-Release Capsules.

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with Dextroamphetamine Sulfate Extended-Release Capsules.

Call your doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking Dextroamphetamine Sulfate Extended-Release Capsules.

2. Mental (Psychiatric) problems:

All Patients

- **new or worse behavior and thought problems**
- **new or worse bipolar illness**
- **new or worse aggressive behavior or hostility**

Children and Teenagers

- **new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms**

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

Call your doctor right away if you or your child have any new or worsening mental symptoms or problems while taking Dextroamphetamine Sulfate Extended-Release Capsules, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.

What are Dextroamphetamine Sulfate Extended-Release Capsules?

Dextroamphetamine Sulfate Extended-Release Capsules are a central nervous system stimulant prescription medicine. **It is used for the treatment of Attention Deficit Hyperactivity Disorder (ADHD).** Dextroamphetamine Sulfate Extended-Release Capsules may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD.

Dextroamphetamine Sulfate Extended-Release Capsules should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Dextroamphetamine Sulfate Extended-Release Capsules are also used in the treatment of a sleep disorder called narcolepsy.

Dextroamphetamine Sulfate is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep Dextroamphetamine Sulfate Extended-Release Capsules in a safe place to prevent misuse and abuse. Selling or giving away Dextroamphetamine Sulfate Extended-Release Capsules may harm others, and is against the law.

Tell your doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take Dextroamphetamine Sulfate Extended-Release Capsules?

Dextroamphetamine Sulfate Extended-Release Capsules should not be taken if you or your child:

- have heart disease or hardening of the arteries
- have moderate to severe high blood pressure
- have hyperthyroidism
- have an eye problem called glaucoma
- are very anxious, tense, or agitated
- have a history of drug abuse
- are taking or have taken within the past 14 days an anti-depression medicine called a monoamine oxidase inhibitor or MAOI.
- is sensitive to, allergic to, or had a reaction to other stimulant medicines

Dextroamphetamine Sulfate Extended-Release Capsules are not recommended for use in children less than 3 years old.

Dextroamphetamine Sulfate Extended-Release Capsules may not be right for you or your child. Before starting Dextroamphetamine Sulfate Extended-Release Capsules tell your or your child's doctor about all health conditions (or a family history of) including:

- heart problems, heart defects, high blood pressure
- mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome
- thyroid problems
- seizures or have had an abnormal brain wave test (EEG)

Tell your doctor if you are, or your child is pregnant, planning to become pregnant, or breastfeeding.

Can Dextroamphetamine Sulfate Extended-Release Capsules be taken with other medicines?

Tell your doctor about all of the medicines that you or your child takes including prescription and non-prescription medicines, vitamins, and herbal supplements.

Dextroamphetamine Sulfate Extended-Release Capsules and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking Dextroamphetamine Sulfate Extended-Release Capsules.

Your doctor will decide whether Dextroamphetamine Sulfate Extended-Release Capsules can be taken with other medicines.

Especially tell your doctor if you or your child take:

- anti-depression medicines including MAOIs
- blood pressure medicines
- antacids
- seizure medicines

Know the medicines that you or your child take. Keep a list of your medicines with you to show your doctor and pharmacist.

Do not start any new medicine while taking Dextroamphetamine Sulfate Extended-Release Capsules without talking to your doctor first.

How should Dextroamphetamine Sulfate Extended-Release Capsules be taken?

- **Take Dextroamphetamine Sulfate Extended-Release Capsules exactly as prescribed.** Your doctor may adjust the dose until it is right for you or your child.
- Dextroamphetamine Sulfate Extended-Release Capsules are usually taken once a day in the morning. Dextroamphetamine Sulfate Extended-Release Capsules is an extended release capsule. It releases medicine into your body throughout the day.
- From time to time, your doctor may stop Dextroamphetamine Sulfate Extended-Release Capsule treatment for a while to check ADHD symptoms.
- Your doctor may do regular checks of the blood, heart, and blood pressure while taking Dextroamphetamine Sulfate Extended-Release Capsules. Children should have their height and weight checked often while taking Dextroamphetamine Sulfate Extended-Release Capsules. Dextroamphetamine Sulfate Extended-Release Capsules treatment may be stopped if a problem is found during these check-ups.
- **If you or your child take too much Dextroamphetamine Sulfate Extended-Release Capsules or overdoses, call your doctor or poison control center right away, or get emergency treatment.**

What are possible side effects of Dextroamphetamine Sulfate Extended-Release Capsules?

See “**What is the most important information I should know about Dextroamphetamine Sulfate Extended-Release Capsules?**” for information on reported heart and mental problems.

Other serious side effects include:

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision

Common side effects include:

- fast heart beat
- headache
- stomach upset
- decreased appetite
- trouble sleeping
- weight loss
- tremors
- dizziness
- dry mouth

Dextroamphetamine Sulfate Extended-Release Capsules may affect your or your child’s ability to drive or do other dangerous activities.

Talk to your doctor if you or your child have side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your doctor or pharmacist for more information.

How should I store Dextroamphetamine Sulfate Extended-Release Capsules?

- Store Dextroamphetamine Sulfate Extended-Release Capsules in a safe place at room temperature, 20° to 25°C (68° to 77°F).
- **Keep Dextroamphetamine Sulfate Extended-Release Capsules and all medicines out of the reach of children.**

General information about Dextroamphetamine Sulfate Extended-Release Capsules

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Dextroamphetamine Sulfate Extended-Release Capsules for a condition for which they were not prescribed. Do not give Dextroamphetamine Sulfate Extended-Release Capsules to other people, even if they have the same condition. It may harm them and it is against the law. This Medication Guide summarizes the most important information about Dextroamphetamine Sulfate Extended-Release Capsules. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Dextroamphetamine Sulfate Extended-Release Capsules that was written for healthcare professionals. For more information about Dextroamphetamine Sulfate Extended-Release Capsules, please contact Barr Laboratories, Inc. at 1-800-BARRLAB (227-7522).

What are the ingredients in Dextroamphetamine Sulfate Extended-Release Capsules?

Active Ingredients: dextroamphetamine sulfate

Inactive Ingredients: colloidal silicon dioxide, dibutyl sebacate, ethylcellulose aqueous dispersion, methylcellulose, povidone, propylene glycol, sugar spheres and talc. The capsule shell ingredients in the 5 mg are D&C red no. 33, FD&C blue no. 1, FD&C yellow no. 6, gelatin, and titanium dioxide. The capsule shell ingredients in the 10 mg are black iron oxide, gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. The capsule shell ingredients in the 15 mg are black iron oxide, gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. The imprinting ingredients are D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, pharmaceutical glaze, propylene glycol, and synthetic black iron oxide.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

BARR LABORATORIES, INC.

Pomona, NY 10970

Revised SEPTEMBER 2007

BR-0954, 0955, 0956

Relabeling and Repackaging by:

Physicians Total Care, Inc.

Tulsa, OK 74146

PRINCIPAL DISPLAY PANEL



Dextroamphetamine Sulfate Extended-Release Capsules 10 mg Text CII

Dextroamphetamine Sulfate

Extended-Release


Capsules

10 mg

Pharmacist: Dispense with Medication Guide.

Rx only

PRINCIPAL DISPLAY PANEL



Dextroamphetamine Sulfate Extended-Release Capsules 15 mg Text CII

Dextroamphetamine

Sulfate

Extended-Release

Capsules

15 mg

Pharmacist: Dispense with Medication Guide.

Rx only

DEXTROAMPHETAMINE SULFATE

dextroamphetamine sulfate capsule, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:54868-5479(NDC:0555-0955)
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROAMPHETAMINE SULFATE (UNII: JJ768O327N) (DEXTROAMPHETAMINE - UNII:TZ47U051F)	DEXTROAMPHETAMINE SULFATE	10 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
METHYLCELLULOSE (15 CPS) (UNII: NPU9M2E6L8)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SUCROSE (UNII: C151H8M554)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4RIU)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
GELATIN (UNII: 2G86QN327L)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LM26O6933)	
SHELLAC (UNII: 46N107B71O)	

Product Characteristics

Color	brown	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	barr;955
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54868-5479-0	60 in 1 BOTTLE, PLASTIC		
2	NDC:54868-5479-1	20 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076137	11/30/2005	

DEXTROAMPHETAMINE SULFATE

dextroamphetamine sulfate capsule, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:54868-5388(NDC:0555-0956)
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTRO AMPHETAMINE SULFATE (UNII: JJ768O327N) (DEXTRO AMPHETAMINE - UNII:TZ47U051FJ)	DEXTRO AMPHETAMINE SULFATE	15 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
METHYLCELLULOSE (15 CPS) (UNII: NPU9M2E6L8)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SUCROSE (UNII: C151H8M554)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
GELATIN (UNII: 2G86QN327L)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LM26O6933)	
SHELLAC (UNII: 46N107B71O)	

Product Characteristics

Color	brown (dark brown)	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	barr;956
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54868-5388-0	10 in 1 BOTTLE, PLASTIC		
2	NDC:54868-5388-1	30 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076137	08/15/2005	

Labeler - Physicians Total Care, Inc. (194123980)

Establishment

Name	Address	ID/FEI	Business Operations
Physicians Total Care, Inc.		194123980	relabel, repack

Revised: 11/2010

Physicians Total Care, Inc.