

**DAYTIME COLD AND FLU NON DROWSY- acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled**  
**CVS Pharmacy**

-----  
**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCL 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

### **When using this product,**

**do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

## **Other information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

## **Inactive ingredients**

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow#6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

## **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

Compare to the active ingredients in Vick® DayQuil® Cold & Flu LiquiCaps®†  
Cold/Flu

Non-Drowsy

### **Daytime**

#### **Multi-Symptom Cold/Flu Relief**

ACETAMINOPHEN - Pain reliever; fever reducer

DEXTROMETHORPHAN HBr - Cough Suppressant

PHENYLEPHRINE HCL - Nasal Decongestant

Relieves:

Aching & fever, sore throat, coughing, and congested-stuffy head so you can get through the day

Alcohol free

Antihistamine free

SOFTGELS\*\*

(\*\*LIQUID-FILLED CAPSULES)

†This product is not manufactured or distributed by the Procter & Gamble Company. Vicks®, Dayquil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWING ANY SIGNS OF TAMPERING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Distributed by: CVS Pharmacy, Inc

One CVS Drive, Woonsocket, RI 02895

CVS.com

1-800-SHOP-CVS

## Package Label

**CVS Health** Compare to the active ingredients in Vicks® DayQuil® Cold & Flu LiquiCaps®†

**Cold/Flu** NDC 69842-848-24

**Non-Drowsy Daytime Multi-Symptom Cold/Flu Relief**

**ACETAMINOPHEN** - Pain reliever; Fever reducer  
**DEXTROMETHORPHAN HBr** - Cough suppressant  
**PHENYLEPHRINE HCl** - Nasal decongestant

**Relieves:**  
Aching & fever, sore throat, coughing,  
and congested-stuffy head so you  
can get through the day

**Alcohol free**  
**Antihistamine free**

**24 SOFTGELS\*\*** (\*\*LIQUID-FILLED CAPSULES)

**Drug Facts**

**Active ingredients (in each softgel)**  
Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Purposes**  
■ temporarily relieves common cold and flu symptoms: ■ minor aches and pains ■ headache  
■ sore throat ■ nasal congestion ■ fever  
■ cough due to minor throat and bronchial irritation

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters  
■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ diabetes ■ heart disease ■ thyroid disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**Drug Facts (continued)**

**When using this product, do not exceed recommended dosage.**  
Stop use and ask a doctor if ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleeplessness occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**  
**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ adults and children 12 years and over: take 2 softgels with water every 4 hours ■ swallow whole: do not crush, chew, or dissolve ■ children under 12 years: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information** ■ store between 15-30°C (59-86°F) ■ avoid excessive heat

**Inactive ingredients** butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**  
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

**Tamper Evident:** DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.  
**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Lot No.:  
Exp. Date:

## CVS HEALTH Non-Drowsy Daytime Cold & Flu Relief

### DAYTIME COLD AND FLU NON DROWSY

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-848
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

## Product Characteristics

<b>Color</b>	orange (Clear)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P19
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69842-848-48	48 in 1 CARTON	03/31/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69842-848-16	16 in 1 CARTON	03/31/2019	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:69842-848-24	24 in 1 CARTON	03/31/2019	

3	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2019	

**Labeler -** CVS Pharmacy (062312574)

Revised: 3/2024

CVS Pharmacy