

MILK OF MAGNESIA ORIGINAL- magnesium hydroxide liquid
Chain Drug Marketing Association

qc mom

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 14 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed

by a doctor

- follow each dosage with a full glass (8 oz) of liquid

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information


- **each 15 mL contains:** magnesium 500 mg
- store at room temperature and avoid freezing
- close cap tightly after use

Inactive ingredients

purified water, sodium hypochlorite

package label

NDC 63868-787-57



*Compare to the
Active Ingredient in
Phillips[®] Milk of Magnesia

Milk of Magnesia

Saline Laxative

Effective Overnight Relief
Very Low Sodium
Stimulant Free
Sugar Free


Original Flavor

12 FL OZ (355 mL)


REV 0618
231-19112-3

TAMPER EVIDENT: DO NOT USE IF IMPRINTED NECKBAND IS MISSING OR BROKEN.

Drug Facts	Purpose								
Active Ingredient (in each 15 mL) Magnesium hydroxide 1200 mg	Saline laxative								
Use									
<ul style="list-style-type: none"> • relieves occasional constipation (irregularity) • generally produces bowel movement in ½ to 6 hours 									
Warnings									
<p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • a magnesium-restricted diet • a sudden change in bowel habits that lasts more than 14 days • kidney disease • stomach pain, nausea or vomiting <p>Ask a doctor or pharmacist before use if you are presently taking a prescription drug</p> <p>This product may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. • you need to use a laxative for more than 1 week <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>									
Directions									
<ul style="list-style-type: none"> • shake well before use • do not exceed the maximum recommended daily dose in a 24 hour period • dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor • follow each dosage with a full glass (8 oz) of liquid <table border="1" style="width: 100%; border-collapse: collapse; margin-top: 5px;"> <thead> <tr> <th style="width: 50%;">age</th> <th style="width: 50%;">dose</th> </tr> </thead> <tbody> <tr> <td>adults and children 12 years and over</td> <td>30 mL to 60 mL</td> </tr> <tr> <td>children 6 to 11 years</td> <td>15 mL to 30 mL</td> </tr> <tr> <td>children under 6 years</td> <td>ask a doctor</td> </tr> </tbody> </table>		age	dose	adults and children 12 years and over	30 mL to 60 mL	children 6 to 11 years	15 mL to 30 mL	children under 6 years	ask a doctor
age	dose								
adults and children 12 years and over	30 mL to 60 mL								
children 6 to 11 years	15 mL to 30 mL								
children under 6 years	ask a doctor								
Other Information									
<ul style="list-style-type: none"> • each 15 mL contains: magnesium 500 mg • store at room temperature and avoid freezing • close cap tightly after use 									
Inactive Ingredients purified water, sodium hypochlorite									
<p>*This product is not manufactured or distributed by Bayer Healthcare LLC., distributors of Phillips[®] brand Milk of Magnesia.</p>									



6 35515 90124 7



Distributed by C.D.M.A., Inc.©
43157 W. Nine Mile
Novi, MI 48376-0995
www.qualitychoice.com
Questions: 248-449-9300 931-19112-5 REV GC1121

MILK OF MAGNESIA ORIGINAL			
magnesium hydroxide liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-787
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-787-57	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	04/01/2014	

Labeler - Chain Drug Marketing Association (011920774)**Registrant** - GCP Laboratories (965480861)**Establishment**

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(63868-787)

Revised: 11/2023

Chain Drug Marketing Association