

**MEMANTINE HYDROCHLORIDE- memantine hydrochloride solution**  
**Macleods Pharmaceuticals Limited**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use MEMANTINE HYDROCHLORIDE solution safely and effectively. See full prescribing information for MEMANTINE HYDROCHLORIDE solution.

**MEMANTINE HYDROCHLORIDE solution, for oral use**

**Initial U.S. Approval: 2003**

----- **RECENT MAJOR CHANGES** -----  
Dosage and Administration ( 2) 08/2014

----- **INDICATIONS AND USAGE** -----  
Memantine hydrochloride, USP is an N-methyl-D-aspartate (NMDA) receptor antagonist indicated for the treatment of moderate to severe dementia of the Alzheimer's type. (1)

----- **DOSAGE AND ADMINISTRATION** -----  
• May be taken with or without food ( 2)  
• Initial dose is 5 mg (2.5 mL) once daily. Increase dose in 5 mg increments to a maintenance dose of 10 mg (5 mL) twice daily. A minimum of 1 week of treatment with the previous dose should be observed before increasing the dose. ( 2)  
• Severe renal impairment: recommended dose is 5 mg (2.5 mL) twice daily. ( 2)

----- **DOSAGE FORMS AND STRENGTHS** -----  
Oral Solution: 2 mg/mL (3)

----- **CONTRAINDICATIONS** -----  
Memantine hydrochloride solution is contraindicated in patients with known hypersensitivity to memantine hydrochloride or to any excipients used in the formulation. (4)

----- **WARNINGS AND PRECAUTIONS** -----  
Conditions that raise urine pH may decrease the urinary elimination of memantine, resulting in increased plasma levels of memantine. ( 5.1, 7.1)

----- **ADVERSE REACTIONS** -----  
Most common adverse reactions ( $\geq 5\%$  and greater than placebo) are dizziness, headache, confusion and constipation. ( 6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Macleods Pharma USA, Inc., at 1-888-943-3210 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**  
**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 10/2015**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS & USAGE**

**2 DOSAGE & ADMINISTRATION**

**3 DOSAGE FORMS & STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Genitourinary Conditions

**6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

7.1 Drugs that Make the Urine Alkaline

7.2 Use with Other N-methyl-D-aspartate (NMDA) Antagonists

## 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

## 10 OVERDOSAGE

## 11 DESCRIPTION

## 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

## 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility
- 13.2 Animal Pharmacology & OR Toxicology

## 14 CLINICAL STUDIES

## 16 HOW SUPPLIED/STORAGE AND HANDLING

## 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS & USAGE

Memantine hydrochloride solution is indicated for the treatment of moderate to severe dementia of the Alzheimer's type.

### 2 DOSAGE & ADMINISTRATION

The recommended starting dose of memantine hydrochloride solution is 5 mg (2.5 mL) once daily. The dose should be increased in 5 mg increments to 10 mg/day (2.5 mL twice daily), 15 mg/day (2.5 mL and 5 mL as separate doses), and 20 mg/day (5 mL twice daily). The minimum recommended interval between dose increases is one week. The dosage shown to be effective in controlled clinical trials is 20 mg/day (5 mL twice daily).

#### Dosing Titration Schedule

	Total daily dose	Strength per dose (mg)
Starting Dose	5 mg	5 mg
Dose after week 1	10 mg	5 mg (first daily dose)
		5 mg (second daily dose)
Dose after week 2	15 mg	5 mg (first daily dose)
		10 mg (second daily dose)
Dose after week 3	20 mg	10 mg (first daily dose)
		10 mg (second daily dose)

Memantine hydrochloride solution can be taken with or without food. If a patient misses a single dose of memantine hydrochloride solution, that patient should not double up on the next dose. The next dose should be taken as scheduled. If a patient fails to take memantine hydrochloride solution for several days, dosing may need to be resumed at lower doses and retitrated as described above.

Do not mix memantine hydrochloride oral solution with any other liquid. The oral solution is administered with a dosing device that comes with the drug and consists of a syringe, syringe adaptor cap, tubing and other supplies a patient needs to administer the drug. The supplied syringe should be used to withdraw the correct volume of oral solution and the oral solution should be slowly squirted into the corner of the patient's mouth.

### Special Populations

#### *Renal Impairment*

A target dose of 5 mg (2.5 mL) twice daily is recommended in patients with severe renal impairment (creatinine clearance of 5 – 29 mL/min based on the Cockcroft-Gault equation).

#### *Hepatic Impairment :*

Memantine hydrochloride solution should be administered with caution to patients with severe hepatic impairment [see *Clinical Pharmacology ( 12.3)*].

## **3 DOSAGE FORMS & STRENGTHS**

Memantine hydrochloride 2 mg/mL oral solution: clear, alcohol-free, sugar-free, and peppermint flavored.

## **4 CONTRAINDICATIONS**

Memantine hydrochloride solution is contraindicated in patients with known hypersensitivity to memantine hydrochloride or to any excipients used in the formulation.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Genitourinary Conditions**

Conditions that raise urine pH may decrease the urinary elimination of memantine resulting in increased plasma levels of memantine [see *Drug Interactions ( 7.1)*].

## **6 ADVERSE REACTIONS**

### **6.1 Clinical Trials Experience**

Memantine hydrochloride was evaluated in eight double-blind placebo-controlled trials involving a total of 1862 dementia (Alzheimer's disease, vascular dementia) patients (940 patients treated with memantine hydrochloride and 922 patients treated with placebo) for a treatment period up to 28 weeks.

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed

in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Adverse Events Leading to Discontinuation

In placebo-controlled trials in which dementia patients received doses of memantine hydrochloride up to 20 mg/day, the likelihood of discontinuation because of an adverse reaction was the same in the memantine hydrochloride group (10.1%) as in the placebo group (11.5%). No individual adverse reaction was associated with the discontinuation of treatment in 1% or more of memantine hydrochloride-treated patients and at a rate greater than placebo.

Most Common Adverse Reactions

In double-blind placebo-controlled trials involving dementia patients, the most common adverse reactions (incidence  $\geq$  5% and higher than placebo) in patients treated with memantine hydrochloride were dizziness, headache, confusion and constipation. Table 1 lists all adverse reactions that occurred in at least 2% of patients treated with memantine hydrochloride and at an incidence greater than placebo.

**Table 1: Adverse Reactions Reported in Controlled Clinical Trials in at Least 2% of Patients Receiving memantine hydrochloride and at a Higher Frequency than Placebo-treated Patients**

Adverse Reaction	Placebo (N = 922) %	Memantine hydrochloride solution (N = 940) %
Body as a Whole		
Fatigue	1	2
Pain	1	3
Cardiovascular System		
Hypertension	2	4
Central and Peripheral Nervous System		
Dizziness	5	7
Headache	3	6
Gastrointestinal System		
Constipation	3	5
Vomiting	2	3
Musculoskeletal System		
Back pain	2	3
Psychiatric Disorders		
Confusion	5	6
Somnolence	2	3
Hallucination	2	3
Respiratory System		
Coughing	3	4
Dyspnea	1	2

The overall profile of adverse reactions and the incidence rates for individual adverse reactions in the subpopulation of patients with moderate to severe Alzheimer's disease were not different from the profile and incidence rates described above for the overall dementia population.

## Seizures

Memantine hydrochloride has not been systematically evaluated in patients with a seizure disorder. In clinical trials of memantine hydrochloride, seizures occurred in 0.2% of patients treated with memantine hydrochloride and 0.5% of patients treated with placebo.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of memantine. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. These reactions include:

Blood and Lymphatic System Disorders -agranulocytosis, leukopenia (including neutropenia), pancytopenia, thrombocytopenia, thrombotic thrombocytopenic purpura.

Cardiac Disorders -cardiac failure congestive.

Gastrointestinal Disorders -pancreatitis.

Hepatobiliary Disorders – hepatitis.

Psychiatric Disorders -suicidal ideation.

Renal and Urinary Disorders -acute renal failure (including increased creatinine and renal insufficiency).

Skin Disorders -Stevens Johnson syndrome.

## **7 DRUG INTERACTIONS**

### **7.1 Drugs that Make the Urine Alkaline**

The clearance of memantine was reduced by about 80% under alkaline urine conditions at pH 8. Therefore, alterations of urine pH towards the alkaline condition may lead to an accumulation of the drug with a possible increase in adverse effects. Urine pH is altered by diet, drugs (e.g. carbonic anhydrase inhibitors, sodium bicarbonate) and clinical state of the patient (e.g. renal tubular acidosis or severe infections of the urinary tract). Hence, memantine should be used with caution under these conditions.

### **7.2 Use with Other N-methyl-D-aspartate (NMDA) Antagonists**

The combined use of memantine hydrochloride with other NMDA antagonists (amantadine, ketamine, and dextromethorphan) has not been systematically evaluated and such use should be approached with caution.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

Teratogenic Effects

Pregnancy Category B

There are no adequate and well-controlled studies of memantine in pregnant women. Memantine hydrochloride should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Memantine given orally to pregnant rats and pregnant rabbits during the period of organogenesis was not teratogenic up to the highest doses tested (18 mg/kg/day in rats and 30 mg/kg/day in rabbits, which are 9 and 30 times, respectively, the maximum recommended human dose [MRHD] on a mg/m<sup>2</sup> basis).

Slight maternal toxicity, decreased pup weights and an increased incidence of non-ossified cervical vertebrae were seen at an oral dose of 18 mg/kg/day in a study in which rats were given oral memantine

beginning pre-mating and continuing through the postpartum period. Slight maternal toxicity and decreased pup weights were also seen at this dose in a study in which rats were treated from day 15 of gestation through the postpartum period. The no-effect dose for these effects was 6 mg/kg, which is 3 times the MRHD on a mg/m<sup>2</sup> basis.

### **8.3 Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when memantine hydrochloride is administered to a nursing mother.

### **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **8.5 Geriatric Use**

The majority of people with Alzheimer's disease are 65 years and older. In the clinical studies of memantine hydrochloride the mean age of patients was approximately 76; over 90% of patients were 65 years and older, 60% were 75 years and older, and 12% were at or above 85 years of age. The efficacy and safety data presented in the clinical trial sections were obtained from these patients. There were no clinically meaningful differences in most adverse events reported by patient groups ≥65 years old and <65 year old.

### **8.6 Renal Impairment**

No dosage adjustment is needed in patients with mild or moderate renal impairment. A dosage reduction is recommended in patients with severe renal impairment [*see Dosage and Administration ( 2) and Clinical Pharmacology ( 12.3)*].

### **8.7 Hepatic Impairment**

No dosage adjustment is needed in patients with mild or moderate hepatic impairment. memantine hydrochloride should be administered with caution to patients with severe hepatic impairment [*see Dosage and Administration ( 2) and Clinical Pharmacology ( 12.3)*].


## **10 OVERDOSAGE**

Signs and symptoms most often accompanying memantine overdosage in clinical trials and from worldwide marketing experience, alone or in combination with other drugs and/or alcohol, include agitation, asthenia, bradycardia, confusion, coma, dizziness, ECG changes, increased blood pressure, lethargy, loss of consciousness, psychosis, restlessness, slowed movement, somnolence, stupor, unsteady gait, visual hallucinations, vertigo, vomiting, and weakness. The largest known ingestion of memantine worldwide was 2.0 grams in a patient who took memantine in conjunction with unspecified antidiabetic medications. The patient experienced coma, diplopia, and agitation, but subsequently recovered. Fatal outcome has been very rarely reported with memantine, and the relationship to memantine was unclear.

Because strategies for the management of overdose are continually evolving, it is advisable to contact a poison control center to determine the latest recommendations for the management of an overdose of any drug. As in any cases of overdose, general supportive measures should be utilized, and treatment should be symptomatic. Elimination of memantine can be enhanced by acidification of urine.

## 11 DESCRIPTION

Memantine hydrochloride, USP is an orally active NMDA receptor antagonist. The chemical name for memantine hydrochloride is 1-amino-3,5-dimethyladamantane hydrochloride with the following structural formula:



The molecular formula is C<sub>12</sub>H<sub>21</sub>N•HCl and the molecular weight is 215.76. Memantine HCl occurs as a fine white to off-white powder and is soluble in water.

Memantine hydrochloride oral solution contains memantine hydrochloride in a strength equivalent to 2 mg of memantine hydrochloride in each mL. The oral solution also contains the following inactive ingredients: sorbitol solution (70%), methyl paraben, propylparaben, propylene glycol, glycerin, natural peppermint flavor #104, citric acid, sodium citrate, and purified water.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Persistent activation of central nervous system N-methyl-D-aspartate (NMDA) receptors by the excitatory amino acid glutamate has been hypothesized to contribute to the symptomatology of Alzheimer's disease. Memantine is postulated to exert its therapeutic effect through its action as a low to moderate affinity uncompetitive (open-channel) NMDA receptor antagonist which binds preferentially to the NMDA receptor-operated cation channels. There is no evidence that memantine prevents or slows neurodegeneration in patients with Alzheimer's disease.

### 12.2 Pharmacodynamics

Memantine showed low to negligible affinity for GABA, benzodiazepine, dopamine, adrenergic, histamine and glycine receptors and for voltage-dependent Ca<sup>2+</sup>, Na<sup>+</sup> or K<sup>+</sup> channels. Memantine also showed antagonistic effects at the 5HT<sub>3</sub> receptor with a potency similar to that for the NMDA receptor and blocked nicotinic acetylcholine receptors with one-sixth to one-tenth the potency.

*In vitro* studies have shown that memantine does not affect the reversible inhibition of acetylcholinesterase by donepezil, galantamine, or tacrine.

### 12.3 Pharmacokinetics

#### *Absorption*

Following oral administration memantine is highly absorbed with peak concentrations reached in about 3-7 hours. Memantine has linear pharmacokinetics over the therapeutic dose range. Food has no effect on the absorption of memantine.

#### *Distribution*

The mean volume of distribution of memantine is 9-11 L/kg and the plasma protein binding is low (45%).

#### *Metabolism*

Memantine undergoes partial hepatic metabolism. The hepatic microsomal CYP450 enzyme system does not play a significant role in the metabolism of memantine.

#### *Elimination*

Memantine is excreted predominantly (about 48%) unchanged in urine and has a terminal elimination half-life of about 60-80 hours.

The remainder is converted primarily to three polar metabolites which possess minimal NMDA receptor antagonistic activity: the N-glucuronide conjugate, 6-hydroxy memantine, and 1-nitrosodeaminated memantine. A total of 74% of the administered dose is excreted as the sum of the parent drug and the N-glucuronide conjugate. Renal clearance involves active tubular secretion moderated by pH dependent tubular reabsorption.

### Pharmacokinetics in Specific Populations

#### *Gender*

Following multiple dose administration of memantine hydrochloride solution 20 mg daily, females had about 45% higher exposure than males, but there was no difference in exposure when body weight was taken into account.

#### *Elderly*

The pharmacokinetics of memantine hydrochloride solution in young and elderly subjects are similar.

#### *Renal Impairment*

Memantine pharmacokinetics were evaluated following single oral administration of 20 mg memantine HCl in 8 subjects with mild renal impairment (creatinine clearance, CL<sub>cr</sub>, >50 – 80 mL/min), 8 subjects with moderate renal impairment (CL<sub>cr</sub> 30 – 49 mL/min), 7 subjects with severe renal impairment (CL<sub>cr</sub> 5 – 29 mL/min) and 8 healthy subjects (CL<sub>cr</sub> > 80 mL/min) matched as closely as possible by age, weight and gender to the subjects with renal impairment. Mean AUC<sub>0-∞</sub> increased by 4%, 60%, and 115% in subjects with mild, moderate, and severe renal impairment, respectively, compared to healthy subjects. The terminal elimination half-life increased by 18%, 41%, and 95% in subjects with mild, moderate, and severe renal impairment, respectively, compared to healthy subjects.

No dosage adjustment is recommended for patients with mild and moderate renal impairment. Dosage should be reduced in patients with severe renal impairment [*see Dosage and Administration ( 2)*].

#### *Hepatic Impairment*

Memantine pharmacokinetics were evaluated following the administration of single oral doses of 20 mg in 8 subjects with moderate hepatic impairment (Child-Pugh Class B, score 7-9) and 8 subjects who were age-, gender-, and weight-matched to the hepatically-impaired subjects. There was no change in memantine exposure (based on C<sub>max</sub> and AUC) in subjects with moderate hepatic impairment as compared with healthy subjects. However, terminal elimination half-life increased by about 16% in subjects with moderate hepatic impairment as compared with healthy subjects. No dose adjustment is recommended for patients with mild and moderate hepatic impairment. Memantine should be



administered with caution to patients with severe hepatic impairment as the pharmacokinetics of memantine have not been evaluated in that population.

### *Drug-Drug Interactions*

#### **Use with Cholinesterase Inhibitors**

Coadministration of memantine with the AChE inhibitor donepezil HCl did not affect the pharmacokinetics of either compound. Furthermore, memantine did not affect AChE inhibition by donepezil. In a 24-week controlled clinical study in patients with moderate to severe Alzheimer's disease, the adverse event profile observed with a combination of memantine hydrochloride and donepezil was similar to that of donepezil alone.

#### **Effect of Memantine Hydrochloride on the Metabolism of Other Drugs**

*In vitro* studies conducted with marker substrates of CYP450 enzymes (CYP1A2, -2A6, -2C9, 2D6, -2E1, -3A4) showed minimal inhibition of these enzymes by memantine. In addition, *in vitro* studies indicate that at concentrations exceeding those associated with efficacy, memantine does not induce the cytochrome P450 isozymes CYP1A2, -2C9, -2E1 and -3A4/5. No pharmacokinetic interactions with drugs metabolized by these enzymes are expected.

Pharmacokinetic studies evaluated the potential of memantine for interaction with warfarin, and bupropion. Memantine did not affect the pharmacokinetics of the CYP2B6 substrate bupropion or its metabolite hydroxybupropion. Furthermore, memantine did not affect the pharmacokinetics or pharmacodynamics of warfarin as assessed by the prothrombin INR.

#### **Effect of Other Drugs on Memantine Hydrochloride**

Memantine is predominantly renally eliminated, and drugs that are substrates and/or inhibitors of the CYP450 system are not expected to alter the metabolism of memantine.

#### **Drugs Eliminated via Renal Mechanisms**

Because memantine is eliminated in part by tubular secretion, coadministration of drugs that use the same renal cationic system, including hydrochlorothiazide (HCTZ), triamterene (TA), metformin, cimetidine, ranitidine, quinidine, and nicotine, could potentially result in altered plasma levels of both agents. However, coadministration of memantine hydrochloride and HCTZ/TA did not affect the bioavailability of either memantine or TA, and the bioavailability of HCTZ decreased by 20%. In addition, coadministration of memantine with the antihyperglycemic drug Glucovance<sup>®</sup> (glyburide and metformin HCl) did not affect the pharmacokinetics of memantine, metformin and glyburide. Furthermore, memantine did not modify the serum glucose lowering effect of Glucovance<sup>®</sup>, indicating the absence of a pharmacodynamic interaction.

#### **Drugs Highly Bound to Plasma Proteins**

Because the plasma protein binding of memantine is low (45%), an interaction with drugs that are highly bound to plasma proteins, such as warfarin and digoxin, is unlikely.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility**

There was no evidence of carcinogenicity in a 113-week oral study in mice at doses up to 40 mg/kg/day (10 times the maximum recommended human dose [MRHD] on a mg/m<sup>2</sup> basis). There was also no evidence of carcinogenicity in rats orally dosed at up to 40 mg/kg/day for 71 weeks followed by 20 mg/kg/day (20 and 10 times the MRHD on a mg/m<sup>2</sup> basis, respectively) through 128 weeks.

Memantine produced no evidence of genotoxic potential when evaluated in the *in vitro* *S. typhimurium* or

*E. coli* reverse mutation assay, an *in vitro* chromosomal aberration test in human lymphocytes, an *in vivo* cytogenetics assay for chromosome damage in rats, and the *in vivo* mouse micronucleus assay. The results were equivocal in an *in vitro* gene mutation assay using Chinese hamster V79 cells.

No impairment of fertility or reproductive performance was seen in rats administered up to 18 mg/kg/day (9 times the MRHD on a mg/m<sup>2</sup> basis) orally from 14 days prior to mating through gestation and lactation in females, or for 60 days prior to mating in males.

### **13.2 Animal Pharmacology & OR Toxicology**

Memantine induced neuronal lesions (vacuolation and necrosis) in the multipolar and pyramidal cells in cortical layers III and IV of the posterior cingulate and retrosplenial neocortices in rats, similar to those which are known to occur in rodents administered other NMDA receptor antagonists. Lesions were seen after a single dose of memantine. In a study in which rats were given daily oral doses of memantine for 14 days, the no-effect dose for neuronal necrosis was 6 times the maximum recommended human dose of 20 mg/day on a mg/m<sup>2</sup> basis

In acute and repeat-dose neurotoxicity studies in female rats, oral administration of memantine and donepezil in combination resulted in increased incidence, severity, and distribution of neurodegeneration compared with memantine alone. The no-effect levels of the combination were associated with clinically relevant plasma memantine and donepezil exposures.

The relevance of these findings to humans is unknown.

## **14 CLINICAL STUDIES**

The clinical efficacy studies described below were conducted with memantine hydrochloride tablets and not with memantine hydrochloride oral solution; however, bioequivalence of memantine hydrochloride oral solution with memantine hydrochloride tablets has been demonstrated.

The effectiveness of memantine hydrochloride as a treatment for patients with moderate to severe Alzheimer's disease was demonstrated in 2 randomized, double-blind, placebo-controlled clinical studies (Studies 1 and 2) conducted in the United States that assessed both cognitive function and day to day function. The mean age of patients participating in these two trials was 76 with a range of 50 to 93 years. Approximately 66% of patients were female and 91% of patients were Caucasian. A third study (Study 3), carried out in Latvia, enrolled patients with severe dementia, but did not assess cognitive function as a planned endpoint. Study Outcome Measures: In each U.S. study, the effectiveness of memantine hydrochloride was determined using both an instrument designed to evaluate overall function through caregiver-related assessment, and an instrument that measures cognition. Both studies showed that patients on memantine hydrochloride experienced significant improvement on both measures compared to placebo.

Day-to-day function was assessed in both studies using the modified Alzheimer's disease Cooperative Study -Activities of Daily Living inventory (ADCS-ADL). The ADCS-ADL consists of a comprehensive battery of ADL questions used to measure the functional capabilities of patients. Each ADL item is rated from the highest level of independent performance to complete loss. The investigator performs the inventory by interviewing a caregiver familiar with the behavior of the patient. A subset of 19 items, including ratings of the patient's ability to eat, dress, bathe, telephone, travel, shop, and

perform other household chores has been validated for the assessment of patients with moderate to severe dementia. This is the modified ADCS-ADL, which has a scoring range of 0 to 54, with the lower scores indicating greater functional impairment.

The ability of memantine hydrochloride to improve cognitive performance was assessed in both studies with the Severe Impairment Battery (SIB), a multi-item instrument that has been validated for the evaluation of cognitive function in patients with moderate to severe dementia. The SIB examines selected aspects of cognitive performance, including elements of attention, orientation, language, memory, visuospatial ability, construction, praxis, and social interaction. The SIB scoring range is from 0 to 100, with lower scores indicating greater cognitive impairment.

### Study 1 (Twenty-Eight-Week Study)

In a study of 28 weeks duration, 252 patients with moderate to severe probable Alzheimer's disease (diagnosed by DSM-IV and NINCDS-ADRDA criteria, with Mini-Mental State Examination scores  $\geq 3$  and  $\leq 14$  and Global Deterioration Scale Stages 5-6) were randomized to memantine hydrochloride or placebo. For patients randomized to memantine hydrochloride, treatment was initiated at 5 mg once daily and increased weekly by 5 mg/day in divided doses to a dose of 20 mg/day (10 mg twice a day).

### Effects on the ADCS-ADL

Figure 1 shows the time course for the change from baseline in the ADCS-ADL score for patients in the two treatment groups completing the 28 weeks of the study. At 28 weeks of treatment, the mean difference in the ADCS-ADL change scores for the memantine hydrochloride -treated patients compared to the patients on placebo was 3.4 units. Using an analysis based on all patients and carrying their last study observation forward (LOCF analysis), memantine hydrochloride treatment was statistically significantly superior to placebo.




Figure 1: Time course of the change from baseline in ADCS-ADL score for patients completing 28 weeks of treatment

Figure 2 shows the cumulative percentages of patients from each of the treatment groups who had attained at least the change in the ADCS-ADL shown on the X axis. The curves show that both patients assigned to memantine hydrochloride and placebo have a wide range of responses and generally show deterioration (a negative change in ADCS-ADL compared to baseline), but that the memantine hydrochloride group is more likely to show a smaller decline or an improvement. (In a cumulative distribution display, a curve for an effective treatment would be shifted to the left of the curve for placebo, while an ineffective or deleterious treatment would be superimposed upon or shifted to the right of the curve for placebo).




Figure 2: Cumulative percentage of patients completing 28 weeks of double-blind treatment with specified changes from baseline in ADCS-ADL scores.

### Effects on the SIB

Figure 3 shows the time course for the change from baseline in SIB score for the two treatment groups over the 28 weeks of the study. At 28 weeks of treatment, the mean difference in the SIB change scores for the memantine hydrochloride -treated patients compared to the patients on placebo was 5.7 units. Using an LOCF analysis, memantine hydrochloride treatment was statistically significantly superior to placebo.




Figure 3: Time course of the change from baseline in SIB score for patients completing 28 weeks of treatment

Figure 4 shows the cumulative percentages of patients from each treatment group who had attained at least the measure of change in SIB score shown on the X axis. The curves show that both patients assigned to memantine hydrochloride and placebo have a wide range of responses and generally show deterioration, but that the memantine hydrochloride group is more likely to show a smaller decline or an improvement.




Figure 4: Cumulative percentage of patients completing 28 weeks of double-blind treatment with specified changes from baseline in SIB scores.

#### Study 2 (Twenty-Four-Week Study)

In a study of 24 weeks duration, 404 patients with moderate to severe probable Alzheimer's disease (diagnosed by NINCDS-ADRDA criteria, with Mini-Mental State Examination scores  $\geq 5$  and  $\leq 14$ ) who had been treated with donepezil for at least 6 months and who had been on a stable dose of donepezil for the last 3 months were randomized to memantine hydrochloride or placebo while still receiving donepezil. For patients randomized to memantine hydrochloride, treatment was initiated at 5 mg once daily and increased weekly by 5 mg/day in divided doses to a dose of 20 mg/day (10 mg twice a day).

#### Effects on the ADCS-ADL

Figure 5 shows the time course for the change from baseline in the ADCS-ADL score for the two treatment groups over the 24 weeks of the study. At 24 weeks of treatment, the mean difference in the ADCS-ADL change scores for the memantine hydrochloride /donepezil treated patients (combination therapy) compared to the patients on placebo/donepezil (monotherapy) was 1.6 units. Using an LOCF analysis, memantine hydrochloride /donepezil treatment was statistically significantly superior to placebo/donepezil.




Figure 5: Time course of the change from baseline in ADCS-ADL score for patients completing 24 weeks of treatment.

Figure 6 shows the cumulative percentages of patients from each of the treatment groups who had attained at least the measure of improvement in the ADCS-ADL shown on the X axis. The curves show that both patients assigned to memantine hydrochloride/donepezil and placebo/donepezil have a wide range of responses and generally show deterioration, but that the memantine hydrochloride /donepezil group is more likely to show a smaller decline or an improvement.




Figure 6: Cumulative percentage of patients completing 24 weeks of double-blind treatment with specified changes from baseline in ADCS-ADL scores.

### Effects on the SIB

Figure 7 shows the time course for the change from baseline in SIB score for the two treatment groups over the 24 weeks of the study. At 24 weeks of treatment, the mean difference in the SIB change scores for the memantine hydrochloride /donepezil-treated patients compared to the patients on placebo/donepezil was 3.3 units. Using an LOCF analysis, memantine hydrochloride /donepezil treatment was statistically significantly superior to placebo/donepezil.






Figure 7: Time course of the change from baseline in SIB score for patients completing 24 weeks of treatment.

Figure 8 shows the cumulative percentages of patients from each treatment group who had attained at least the measure of improvement in SIB score shown on the X axis. The curves show that both patients assigned to memantine hydrochloride/donepezil and placebo/donepezil have a wide range of responses, but that the memantine hydrochloride /donepezil group is more likely to show an improvement or a smaller decline.




Figure 8: Cumulative percentage of patients completing 24 weeks of double-blind treatment with specified changes from baseline in SIB scores.

### Study 3 (Twelve-Week Study)

In a double-blind study of 12 weeks duration, conducted in nursing homes in Latvia, 166 patients with dementia according to DSM-III-R, a Mini-Mental State Examination score of < 10, and Global Deterioration Scale staging of 5 to 7 were randomized to either memantine hydrochloride or placebo. For patients randomized to memantine hydrochloride, treatment was initiated at 5 mg once daily and increased to 10 mg once daily after 1 week. The primary efficacy measures were the care dependency subscale of the Behavioral Rating Scale for Geriatric Patients (BGP), a measure of day-to-day function, and a Clinical Global Impression of Change (CGI-C), a measure of overall clinical effect. No valid measure of cognitive function was used in this study. A statistically significant treatment difference at 12 weeks that favored memantine hydrochloride over placebo was seen on both primary efficacy measures. Because the patients entered were a mixture of Alzheimer's disease and vascular dementia, an attempt was made to distinguish the two groups and all patients were later designated as having either vascular dementia or Alzheimer's disease, based on their scores on the Hachinski Ischemic Scale at study entry. Only about 50% of the patients had computerized tomography of the brain. For the subset designated as having Alzheimer's disease, a statistically significant treatment effect favoring memantine hydrochloride over placebo at 12 weeks was seen on both the BGP and CGI-C.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Oral Solution:

2 mg/mL Oral Solution (10 mg = 5 mL)

240 mL, bottle NDC 33342-066-28

Store memantine hydrochloride oral solution at 20°C to 25°C (68°F to 77°F) ; excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (*Patient Information and Instructions for Use*).

To assure safe and effective use of memantine hydrochloride solution, the following information and instructions provided in the patient information section should be discussed with patients and caregivers.

Patients/caregivers should be instructed to follow the dose titration schedule provided by their physician or healthcare professional for memantine hydrochloride solution.

If a patient misses a single dose of memantine hydrochloride solution, that patient should not double up on the next dose. The next dose should be taken as scheduled. If a patient fails to take memantine hydrochloride solution for several days, dosing should not be resumed without consulting that patient's healthcare professional.

Patients/caregivers should be instructed on how to use the memantine hydrochloride Oral Solution dosing device. They should be made aware of the patient instruction sheet that is enclosed with the product. Patients/caregivers should be instructed to address any questions on the usage of the solution to their physician or pharmacist.

Manufactured for:

**Macleods Pharma USA, Inc.**

Plainsboro, NJ 08536

Manufactured by:

Macleods Pharmaceutical Ltd.

Baddi, Himachal Pradesh, INDIA

**Revised** : October 2015

## SPL PATIENT PACKAGE INSERT SECTION

### Patient Information

#### **Memantine Hydrochloride Oral Solution (me-MAN-teen HYE-droe-KLOR-ide)**

Read this Patient Information that comes with memantine hydrochloride solution before you start taking it and each time you get a refill. There may be new information. This information does not take the place of talking to your doctor about your medical condition or your treatment.

#### **What is memantine hydrochloride solution?**

Memantine hydrochloride solution is a prescription medicine used for the treatment of moderate to severe dementia in people with Alzheimer's disease. memantine hydrochloride belongs to a class of medicines called NMDA (N-methyl-D-aspartate) inhibitors.

It is not known if memantine hydrochloride solution is safe and effective in children.

#### **Who should not take memantine hydrochloride solution?**

**Do not take memantine hydrochloride solution if you** are allergic to memantine or any of the ingredients in memantine hydrochloride solution. See the end of this leaflet for a complete list of ingredients in memantine hydrochloride solution.

#### **What should I tell my doctor before taking memantine hydrochloride solution?**

## **Before you take memantine hydrochloride solution, tell your doctor if you:**

- have or have had seizures
- have or have had problems passing urine
- have or have had bladder or kidney problems
- have liver problems
- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if memantine hydrochloride solution will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if memantine hydrochloride passes into your breast milk. You and your doctor should decide if you will take memantine hydrochloride solution or breastfeed.

**Tell your doctor about all the medicines you take**, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Taking memantine hydrochloride solution with certain other medicines may affect each other. Taking memantine hydrochloride solution with other medicines can cause serious side effects.

Especially tell your doctor if you take:

- other NMDA antagonists such as amantadine, ketamine, and dextromethorphan
- medicines that make your urine alkaline such as carbonic anhydrase inhibitors and sodium bicarbonate

Ask your doctor or pharmacist for a list of these medicines, if you are not sure.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

## **How should I take memantine hydrochloride solution?**

- **See the step-by-step instructions for taking memantine hydrochloride Oral Solution at the end of this Patient Information.**
- Your doctor will tell you how much memantine hydrochloride solution to take and when to take it.
- Your doctor may change your dose if needed.
- Memantine hydrochloride solution can be taken with food or without food.
- If you forget to take one dose of memantine hydrochloride solution, do not double up on the next dose. You should take only the next dose as scheduled.
- If you have forgotten to take memantine hydrochloride solution for several days, you should not take the next dose until you talk to your doctor.
  - If you take too much memantine hydrochloride solution, call your doctor or poison control center at 1-800222-1222 right away, or go to the nearest hospital emergency room.

## **What are the possible side effects of memantine hydrochloride solution?**

### **Memantine hydrochloride solution may cause side effects, including:**

The most common side effects of memantine hydrochloride solution include:

- 1 dizziness
- 2 confusion
- 3 headache
- 4 constipation

These are not all the possible side effects of memantine hydrochloride solution. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store memantine hydrochloride solution?**

• Store memantine hydrochloride oral solution at 20°C to 25°C (68°F to 77°F) ; excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

**What are the ingredients in memantine hydrochloride oral solution?****Memantine hydrochloride oral solution oral solution:**

Active ingredients: memantine hydrochloride, USP

Inactive ingredients: sorbitol solution (70%), methyl paraben, propylparaben, propylene glycol, glycerin, natural peppermint flavor #104, citric acid, sodium citrate, and purified water

**Keep memantine hydrochloride oral solution and all medicines out of the reach of children.****General information about the safe and effective use of memantine hydrochloride oral solution.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not take memantine hydrochloride oral solution for a condition for which it was not prescribed. Do not give memantine hydrochloride oral solution to other people, even if they have the same condition. It may harm them.

This Patient Information leaflet summarizes the most important information about memantine hydrochloride oral solution. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about memantine hydrochloride oral solution that was written for healthcare professionals.

For more information about memantine hydrochloride oral solution, call Macleods Pharma USA, Inc., at 1-888-943-3210.

**INSTRUCTIONS FOR USE****Memantine Hydrochloride Oral Solution**



(me-MAN-teen HYE-droe-KLOR-ide)

**Directions for Using your Memantine Hydrochloride Oral Solution**


Read these instructions before taking Memantine Hydrochloride Oral Solution and each time you get a refill. There may be new information. This information does not take the place of talking to your doctor about your medical condition or your treatment.

Preparing your dose of Memantine Hydrochloride Oral Solution. You will need the following supplies:

1. Memantine Hydrochloride Oral Solution bottle with Child-resistant cap
2. Blue syringe adaptor cap with lid
3. Oral dosing syringe
4. Plastic tube (in plastic bag)
5. Prescribing Information



1. Remove the oral dosing syringe, the blue syringe adaptor cap, and remove the plastic tube from its protective plastic bag. Attach the tube to the blue syringe adaptor cap if it is not already attached.



2. The bottle comes with a child-resistant cap. To remove the cap, you should push down on the cap and at the same time; turn the cap counter-clockwise (to the left).



3. Carefully remove the seal from the bottle



and throw away.



4. Insert the blue syringe adapter cap, with the attached tube, all the way into the bottle and tightly screw the cap onto the bottle by turning the cap clockwise (to the right).




5. The blue syringe adaptor cap has an opening with an attached lid. The adaptor is used to withdraw the correct dose of medicine from the bottle with the syringe. The attached adaptor lid should be closed in between doses.



6. Keep the bottle upright on a table. Open the syringe adaptor lid and insert the tip of syringe into the syringe adaptor opening.

- Make sure the syringe is pushed firmly into the adaptor opening.



7. While holding the syringe in place, gently pull the plunger of the syringe until you get to the correct mL (amount) of medicine you need.





- Do not worry about a few tiny bubbles. This will not affect your dose.



8. Remove the syringe from the syringe adaptor cap.



9. Remove the syringe from the bottle and slowly squirt the Memantine Hydrochloride Oral Solution into the corner of you or the

	<p>oral solution into the corner of your or the patient's mouth. <b>Do not mix Memantine Hydrochloride Oral Solution with any other liquid.</b></p>
	<p>10. After use, reseal the bottle by snapping the attached syringe adaptor lid closed.</p>
	<p>11. Rinse the empty syringe by inserting the open end of the syringe into a glass of water, pulling the plunger out to draw in water, and pushing the plunger in to remove the water. Repeat several times. Allow the syringe to air dry.</p>
	<p>12. Store the bottle upright.</p>

This Patient Information and Instructions for Use have been approved by the U.S. Food and Drug Administration.

Manufactured for:  
**Macleods Pharma USA, Inc.**  
 Plainsboro, NJ 08536

Manufactured by:  
 Macleods Pharmaceutical Ltd.  
 Baddi, Himachal Pradesh, INDIA

**Revised:** October 2015

**PM02007101**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

Rx Only



NDC 33342-066-28  
Memantine Hydrochloride Oral Solution  
2 mg/mL  
Bottle of 240 mL

**MACLEODS**  
Pharma USA  
NDC 33342-066-28

Each mL of oral solution contains :  
Memantine hydrochloride, USP 2 mg

USUAL DOSAGE :  
See prescribing information for Dosage and administration.

Batch No. :  
Exp. Date :

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

Keep out of reach of children

Dispense in tight container

Manufactured for :  
**Macleods Pharma USA, Inc.**  
Plainsboro, NJ 08536

Manufactured by :  
Macleods Pharmaceuticals Ltd.  
Baddi, Himachal Pradesh, INDIA

Code No.: HP/152/07  
PM02005001

**Memantine Hydrochloride Oral Solution**  
**2 mg/mL**

**Rx Only**      **240 mL**


Pharmacist: Dispense the Patient Information Leaflet with the Drug Product.

OPZ AREA  
23 x 23 mm

Rx Only  
NDC 33342-066-28  
Memantine Hydrochloride Oral Solution  
2 mg/mL  
Carton for Bottle of 240 mL

Sr. No.:  
 GTIN:  
 Exp. Date:  
 Batch No.:

OPZ AREA  
 62 X 36 mm



PM02013201

 <p>NDC 33342-066-28</p> <p><b>Memantine Hydrochloride Oral Solution</b>  <b>2 mg/mL</b></p> <p><b>Pharmacist:</b> Dispense the Patient Information Leaflet with the Drug Product.</p> <p><b>R<sub>x</sub> Only</b> <b>240 mL</b></p>	<p><b>Memantine Hydrochloride Oral Solution</b>  <b>2 mg/mL</b></p>  <p><b>Macleods Pharma USA</b>      Manufactured for :  <b>Macleods Pharma USA, Inc.</b>      Plainsboro, NJ 08536</p> <p>Manufactured by:      Macleods Pharmaceuticals Ltd.      Baddi, Himachal Pradesh, INDIA      Code No.: HP/152/07</p>	 <p>NDC 33342-066-28</p> <p><b>Memantine Hydrochloride Oral Solution</b>  <b>2 mg/mL</b></p> <p><b>Pharmacist:</b> Dispense the Patient Information Leaflet with the Drug Product.</p> <p><b>R<sub>x</sub> Only</b> <b>240 mL</b></p>	<p><b>Memantine Hydrochloride Oral Solution</b>  <b>2 mg/mL</b></p> <p>Each mL of oral solution contains :      Memantine hydrochloride, USP 2 mg</p> <p><b>USUAL DOSAGE :</b>      See prescribing information for Dosage and administration.</p> <p>Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].</p> <p>Keep out of reach of children</p> <p>Dispense in tight container</p>
--	---	---	--

**MEMANTINE HYDROCHLORIDE**

memantine hydrochloride solution

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:33342-066
<b>Route of Administration</b>	ORAL		

<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
MEMANTINE HYDROCHLORIDE (UNII: JY0WD0UA60) (MEMANTINE - UNII:W8O17SJF3T)		MEMANTINE HYDROCHLORIDE	2 mg in 1 mL	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
SORBITOL (UNII: 506T60A25R)				
METHYL PARABEN (UNII: A2I8C7HI9T)				
PROPYL PARABEN (UNII: Z8IX2SC1OH)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
GLYCERIN (UNII: PDC6A3C0OX)				
PEPPERMINT (UNII: V95R5KMY2B)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
SODIUM CITRATE (UNII: 1Q73Q2JULR)				
WATER (UNII: 059QF0KO0R)				
<b>Product Characteristics</b>				
<b>Color</b>		<b>Score</b>		
<b>Shape</b>		<b>Size</b>		
<b>Flavor</b>	PEPPERMINT (PEPPERMINT)	<b>Imprint Code</b>		
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33342-066-28	240 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/13/2015	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA202790	10/13/2015		

**Labeler** - Macleods Pharmaceuticals Limited (862128535)

### Establishment

Name	Address	ID/FEI	Business Operations
Macleods Pharmaceuticals Limited		676369519	analysis(33342-066) , label(33342-066) , manufacture(33342-066) , pack(33342-066)