# OXY-GARD M SANITIZING TEAT DIP- hydrogen peroxide, dodecylbenzenesulfonic acid, lactic acid solution Ecolab Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

# **KEEP OUT OF REACH OF CHILDREN**

# CAUTION!

MAY CAUSE EYE IRRITATION. Use suitable protective equipment. Wash thoroughly after handling.

# FOR INDUSTRIAL USE ONLY - NOT FOR HOUSEHOLD USE

# NOT FOR HUMAN USE

# Read other label for directions for use, first aid and precautionary statements

## **DIRECTIONS:**

**IMPORTANT:** Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container.

**Udder Prep:** When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

## **Directions for Teat Dipping**

**Pre-Milk Dipping:** Before each cow is milked, and using fresh Oxy-Gard M, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.

**Post-Milk Dipping:** Using fresh Oxy-Gard M, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

# **Directions for Teat Spraying**

**Pre-Milk Spraying:** Before each cow is milked, and using fresh Oxy-Gard M, spray entire teat. Wipe teats after dipping using single-service towels to avoid contamination of milk.

**Post-Milk Spraying:** Using fresh Oxy-Gard M, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

**Expanded Usage:** When freshening cows, begin dipping teats twice daily for about 10 days before calving.

**PRECAUTION:** Oxy-Gard M is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.

# FIRST AID

**Eye Contact:** In case of contact, immediately flush eyes with plenty of water. Remove contact lenses and flush again. Get medical attention if irritation persists.

Skin Contact: No known effect after skin contact. Rinse with water for a few minutes.

Inhalation: No special measures required. Treat symptomatically.

**Ingestion:** Get medical attention if symptoms occur.

# READ MATERIAL SAFETY DATA SHEET (MSDS) BEFORE USING THIS PRODUCT

**EMERGENCY HEALTH INFORMATION:** 1 800 328 0026. If located outside the United States and Canada, call collect 1 651 222 5352 (number is in the US).

Principal display panel and representative label

ECOLAB
Oxy-Gard™ M
Sanitizing Teat Dip
An aid in reducing the spread of organisms which may cause mastitis
Contains an emollient system (glycerin and sorbitol) 11%
ACTIVE INGREDIENTS:
Hydrogen peroxide0.5%
Lactic acid 1.7%
Dodecyl benzene sulfonic acid0.5%
INERT INGREDIENTS:
TOTAL:100.0%
208 L (55 US gal)
6300644
753774/5301/0713
Ecolab $\cdot$ 370 Wabasha Street N $\cdot$ St Paul MN 55102-1390 USA $\cdot$ tel: 1 800 392 3392
$^{\odot}$ 2012 Ecolab USA Inc. $\cdot$ All rights reserved
Made in United States
www.ecolab.com · 753777/5300/1112

# $\textbf{Oxy-Gard}^{{}^{\scriptscriptstyle{\mathsf{M}}}} \textbf{M}$

# Sanitizing Teat Dip Desinfectante de pezones por inmersión

- An aid in reducing the spread of organisms which may cause mastitis | Ayuda a evitar la proliferación de organismos que pueden causar mastitis
- Contains an emollient system (glycerin and sorbitol) 11% | Contiene un sistema emoliente (glicerina y sorbitol) 11%

ACTIVE INGREDIENTS:

Hydrogen peroxide	0.5%
Lactic acid	
Dodecyl benzene sulfonic acid	0.5%
INERT INGREDIENTS:	97.3%
TOTAL:	100.0%

CAUTION | PRECAUCIÓN

KEEP OUT OF REACH OF CHILDREN | MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS FOR INDUSTRIAL USE ONLY - NOT FOR HOUSEHOLD USE | PARA USO INDUSTRIAL SOLAMENTE - NO ES PARA USO DOMESTICO NOT FOR HUMAN USE | NO ES PARA USO HUMANO

Read other label for directions for use, first aid and precautionary statements Lea la otra etiqueta para direcciones de uso, primeros auxilios y medidas preventivas 753774/5301/0713






Teat Dip

Baño de tetina









# Oxy-Gard<sup>™</sup> M

#### Sanitizing Teat Dip

#### FOR INDUSTRIAL USE ONLY

DIRECTIONS:

IMPORTANT: Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container.

Udder Prep: When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

Directions for Teat Dipping

Pre-Milk Dipping: Before each cow is milked, and using fresh Oxy-Gard M, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Using fresh Oxy-Gard M, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

#### **Directions for Teat Spraying**

Pre-Milk Spraying: Before each cow is milked, and using fresh Oxy-Gard M, spray entire teat. Wipe teats after dipping using singleservice towels to avoid contamination of milk.

Post-Milk Spraying: Using fresh Oxy-Gard M, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving

PRECAUTION: Oxy-Gard M is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.

#### KEEP OUT OF REACH OF CHILDREN

CAUTION!

MAY CAUSE EYE IRRITATION. Use suitable protective equipment. Wash thoroughly after handling.

#### FIRST AID

Eye Contact: In case of contact, immediately flush eyes with plenty of water. Remove contact lenses and flush again. Get medical attention if irritation persists.

Skin Contact: No known effect after skin contact. Rinse with water for a few minutes.

Inhalation: No special measures required. Treat symptomatically.

#### Ingestion: Get medical attention if symptoms occur.

READ MATERIAL SAFETY DATA SHEET (MSDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: 1 800 328 0026. If located outside the United States and Canada, call collect 1 65i 222 5352 (number is in the US).

#### Desinfectante de pezones por inmersión PARA USO INDUSTRIAL SOLAMENTE

#### INSTRUCCIONES:

IMPORTANTE: No lo vuelva a diluir con agua ni lo mezcle con otros productos para baño de pezones. Si el producto del recipiente para realizar el baño está visiblemente sucio, deseche el conterido y rellene el recipiente con producto fresco. No vuelva a utilizar el producto ni vuelva a colocar producto sin usar en el contenedor original.

Preparado para ubres: Si se realiza un paso de limpieza de ubres antes del ordeño, asegúrese de limpiar los pezones con una solución para limpieza de ubres adecuada y procedimientos de limpieza apropiados. Los pezones deben secarse con toallas de uso único.

#### Instrucciones para el sellado de pezones

Sellado del pezón previo al ordeño: Antes de ordeñar cada vaca, bañe toda la superficie de cada pezón en un recipiente que contengo (xxy-Gard M fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche. Sellado del pezón posterior al ordeño: Bañe toda la superficie de cada pezón en un recipiente que contenga Oxy-Gard M fresco. No utilice paños. Deje secara al air co de No deje las vacas en un clima helado hasta que el producto se haya secado por completo. Instrucciones para el rociado de pezones

Rociado previo al ordeño: Antes de ordeñar cada vaca, rocíe toda la superficie de cada pezón con Oxy-Gard M fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche.

Rociado de pezones posterior al ordeño: Inmediatamente después del ordeño, rocíe toda la superficie de cada pezón con Oxy-Gard M fresco. No utilice paños. Deje secar al aire. No deje las vacas en un clima helado hasta que el producto se haya secado por completo. Uso ampliado: Si la vaca va a parir, comience a realizar el baño dos veces al día unos 10 días antes del parto.

PRECAÚCIÓN: Oxy-Gard M no está pensado para curar ni ayudar a sanar los pezones agrietados o irritados. Como cualquier germicida, el producto puede causar irritación o sensibilización en los animales sensibles. Si los pezones están agrietados o irritados, encarque a un veterinario que examine la condición y la trate si es necesario.

Consulte con su Representante de ECOLAB para obtener instrucciones específicas sobre el uso y el equipo dispensador recomendado. MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

#### PRECAUCIÓN!

PUEDE PROVOCAR IRRITACIÓN OCULAR. Use equipo protector adecuado. Lávese completamente después del manejo. PRIMEROS AUXILIOS

Contacto con los ojos: En caso de contacto, lávese de inmediato los ojos con abundante agua. Dultese los lentes de contacto y lávese nuevamente. Obtenga atención médica si persiste la irritación.

Contacto con la piel: No se conocen efectos sobre el contacto con la piel, enjuague con agua durante unos minutos. Inhalación: No se requieren medidas especiales. Tratar sintomáticamente.

Ingestión: Obtenga atención médica si se presentan síntomas.

#### LEA LA HOJA DE DATOS DE SEGURIDAD ANTES DE USAR ESTE PRODUCTO

INFORMACIÓN DE EMERGENCIA SOBRE SALUD: 1 800 328 0026. Si está fuera de los Estados Unidos y Canadá, llame a cobro revertido al 1 651 222 5352 (número en los EE.UU.).

Ecolab - 370 Wabasha Street N - St Paul MN 55102-1390 USA - tel: 1 800 392 3392 © 2012 Ecolab USA Inc. - All rights reserved | Todos los derechos reservados Made in United States | Hecho en Estados Unidos www.ecolab.com - 753771/5300/112

# **OXY-GARD M SANITIZING TEAT DIP**

hydrogen peroxide, dodecylbenzenesulfonic acid, lactic acid solution

Product Information							
Product Information							
Product T ype	OTC ANIMAL DRUG	Item Code (Source)		NDC:47593-121			
Route of Administration	TOPICAL						
Active Ingredient/Active Moiety							
Ingr	edient Name		Basis of Sti	rength	Strength		
HYDROGEN PEROXIDE (UNII: BBX06 UNII:BBX060AN9V)	0 AN9 V) (HYDROGEN PEROXIDE	2 -	HYDROGEN PERO	XIDE	5 mg in 1 mL		
<b>DODECYLBENZENESULFONIC ACID</b> (DODECYLBENZENESULFONIC ACID			DODECYLBENZEN ACID	NESULFONIC	5 mg in 1 mL		
					17 mg		

LACTIC ACID

<b>Inactive Ingredient</b>	s					
	Ingredient Name			Strength		
SORBITOL (UNII: 506T6	0A25R)					
GLYCERIN (UNII: PDC6A	3C0OX)					
Packaging						
# Item Code	Package Description	Marketing Start Date		Marketing End Date		
<b>1</b> NDC:47593-121-17	56780 mL in 1 DRUM					
<b>2</b> NDC:47593-121-21	208200 mL in 1 DRUM					
<b>3</b> NDC:47593-121-88	1040900 mL in 1 CONTAINER					
Marketing Info	mation					
Marketing Info Marketing Category	<b>'mation</b> Application Number or Monograph	Citation	Marketing Start Da	te Marketing End Date		

Labeler - Ecolab Inc. (006154611)

Revised: 10/2019

Ecolab Inc.