

GUAIFENESIN 400MG- guaifenesin tablet
Marlex Pharmaceuticals Inc

Guaifenesin 400mg

GUAIFENESIN400MG TABLETS

MarlexPharmaceuticals, Inc.

Drug Fact

Active ingredient (in each tablet)

Guaifenesin 400mg

Purpose

Expectorant

Guaifenesin 400 mg Tablets

Uses

helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageway of bothersome mucus and make coughs more productive

Warnings

Ask a doctor before use if you have

- Persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- Cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if

cough last more than 7 days, come back or is accompanied by fever, rash, or persistent headache. There could be signs of a serious illness.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

If pregnant or breast- feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- Do not take more than 6 doses in any 24-hour period.
- This product is not intended for use in children under 12 years of age
- Adults & children 12 years and over: 1 to 2 tablets every 4 hours
- Children under 12 years: do not use

Other Information:

- Store at 15°C-30°C (59°F-86°F)

Inactive Ingredients:

Colloidal silicon dioxide, FD&C RED #40(AI-lake), magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, stearic acid

Manufactured for & Distributed by:

Marlex Pharmaceuticals, Inc.

New Castle, DE 19720

Rev: 02/19TCL

PRINCIPAL DISPLAY PANEL

NDC 10135-0682-30

Guaifenesin

400 mg

30 CAPLETS



PRINCIPAL DISPLAY PANEL

NDC 10135-0682-60
Guaifenesin
400 mg
60 CAPLETS

NDC 10135-0682-60

MARLEX
PHARMACEUTICALS, INC.

Guaifenesin

400 mg

60 CAPLETS

Drug Facts

Active ingredient (in each caplet) Purpose
Guaifenesin 400 mgExpectorant

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both some mucus and make coughs more productive

Warnings
Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious illness.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Manufactured for/Distributed by:
Marlex Pharmaceuticals, Inc.
New Castle, DE 19720

Rev. 03/19 HP


N 3 10135 68260 7

Peel Here
For More Drug Facts

Drug Facts (continued on back)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- this product is not intended for use in children under 12 years of age
- adults & children 12 years and over: 1 caplet every 4 hours
- children under 12 years: do not use

Other information

- store at 15°C - 30°C (59°F - 86°F)

Inactive ingredients

colloidal silicon dioxide, magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, stearic acid

<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>
<p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than 6 doses in any 24-hour period ■ this product is not intended for use in children under 12 years of age ■ adults & children 12 years and over: 1 caplet every 4 hours ■ children under 12 years: do not use
<p>Other information</p> <ul style="list-style-type: none"> ■ store at 15°C - 30°C (59°F - 86°F)
<p>Inactive ingredients</p> <p>colloidal silicon dioxide, magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, stearic acid</p>

GUAIFENESIN 400MG

guaifenesin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10135-682
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Caplet)	Size	10mm
Flavor		Imprint Code	G400
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10135-682-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2019	
2	NDC:10135-682-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2019	

Labeler - Marlex Pharmaceuticals Inc (782540215)

Revised: 11/2023

Marlex Pharmaceuticals Inc