


**TIZANIDINE HYDROCHLORIDE - tizanidine hydrochloride tablet**  
**STAT Rx USA LLC**

-----  
**Tizanidine Hydrochloride Tablets 2 mg and 4 mg**

**Rx Only**

**DESCRIPTION**

Tizanidine hydrochloride is a centrally acting  $\alpha_2$ -adrenergic agonist. Tizanidine HCl (tizanidine) is a white to off-white, fine crystalline powder, odorless or with a faint characteristic odor. Tizanidine is slightly soluble in water and methanol; solubility in water decreases as the pH increases. Its chemical name is 5-chloro-4-(2-imidazolin-2-ylamino)-2,1,3-benzothiazole hydrochloride. Tizanidine's molecular formula is  $C_9H_8ClN_5S \cdot HCl$ , its molecular weight is 290.2 and its structural formula is:



Tizanidine is supplied as 2 and 4 mg tablets for oral administration. Tizanidine Hydrochloride Tablets are composed of the active ingredient, tizanidine hydrochloride (2.288 mg equivalent to 2 mg tizanidine base and 4.576 mg equivalent to 4 mg tizanidine base) and the inactive ingredients, colloidal silicon dioxide, lactose monohydrate, microcrystalline cellulose and stearic acid.

**CLINICAL PHARMACOLOGY**

**Mechanism of Action**

Tizanidine is an agonist at  $\alpha_2$ -adrenergic receptor sites and presumably reduces spasticity by increasing presynaptic inhibition of motor neurons. In animal models, tizanidine has no direct effect on skeletal muscle fibers or the neuromuscular junction, and no major effect on monosynaptic spinal reflexes. The effects of tizanidine are greatest on polysynaptic pathways. The overall effect of these actions is thought to reduce facilitation of spinal motor neurons.

The imidazoline chemical structure of tizanidine is related to that of the anti-hypertensive drug clonidine and other  $\alpha_2$ -adrenergic agonists. Pharmacological studies in animals show similarities between the two compounds, but tizanidine was found to have one-tenth to one-fiftieth (1/50) of the potency of clonidine in lowering blood pressure.

**Pharmacokinetics**


A single dose of two 4 mg tablets was administered under fed and fasting conditions in an open label, four period, randomized crossover study in 96 human volunteers, of whom 81 were eligible for the

statistical analysis.

Following oral administration of the tablet (in the fasted state), tizanidine has peak plasma concentrations occurring 1 hour after dosing with a half-life of approximately 2 hours.

When two 4 mg tablets are administered with food the mean maximal plasma concentration is increased by approximately 30% and the median time to peak plasma concentration is increased by 25 minutes, to 1 hour and 25 minutes.

Food also increases the extent of absorption of the tablet. (see Figures 1 and 2).



**Figure 1: Mean Tizanidine Concentration vs. Time Profiles for Tizanidine Tablets (2 x 4 mg) Under Fasted and Fed Conditions. (Linear Scale)**

### Special Populations

**Age Effects:** No specific pharmacokinetic study was conducted to investigate age effects. Cross study comparison of pharmacokinetic data following single dose administration of 6 mg tizanidine showed that younger subjects cleared the drug four times faster than the elderly subjects. Tizanidine has not been evaluated in children (see **PRECAUTIONS**).

**Hepatic Impairment:** Pharmacokinetic differences due to hepatic impairment have not been studied (see **WARNINGS**).

**Renal Impairment:** Tizanidine clearance is reduced by more than 50% in elderly patients with renal insufficiency (creatinine clearance < 25 mL/min) compared to healthy elderly subjects; this would be expected to lead to a longer duration of clinical effect. Tizanidine should be used with caution in renally impaired patients (see **PRECAUTIONS**).

**Gender Effects:** No specific pharmacokinetic study was conducted to investigate gender effects. Retrospective analysis of pharmacokinetic data, however, following single and multiple dose administration of 4 mg tizanidine showed that gender had no effect on the pharmacokinetics of tizanidine.

*Race Effects:* Pharmacokinetic differences due to race have not been studied.

*Drug Interactions-Oral Contraceptives:* No specific pharmacokinetic study was conducted to investigate interaction between oral contraceptives and tizanidine. Retrospective analysis of population pharmacokinetic data following single and multiple dose administration of 4 mg tizanidine, however, showed that women concurrently taking oral contraceptives had 50% lower clearance of tizanidine compared to women not on oral contraceptives (see PRECAUTIONS).


## Clinical Studies

Tizanidine's capacity to reduce increased muscle tone associated with spasticity was demonstrated in two adequate and well controlled studies in patients with multiple sclerosis or spinal cord injury.

In one study, patients with multiple sclerosis were randomized to receive single oral doses of drug or placebo. Patients and assessors were blind to treatment assignment and efforts were made to reduce the likelihood that assessors would become aware indirectly of treatment assignment (e.g., they did not provide direct care to patients and were prohibited from asking questions about side effects). In all, 140 patients received either placebo, 8 mg or 16 mg of tizanidine.

Response was assessed by physical examination; muscle tone was rated on a 5 point scale (Ashworth score), with a score of 0 used to describe normal muscle tone. A score of 1 indicated a slight spastic catch while a score of 2 indicated more marked muscle resistance. A score of 3 was used to describe considerable increase in tone, making passive movement difficult. A muscle immobilized by spasticity was given a score of 4. Spasm counts were also collected.

Assessments were made at 1, 2, 3 and 6 hours after treatment. A statistically significant reduction of the Ashworth score for tizanidine compared to placebo was detected at 1, 2 and 3 hours after treatment. Figure 2 below shows a comparison of the mean change in muscle tone from baseline as measured by the Ashworth scale. The greatest reduction in muscle tone was 1 to 2 hours after treatment. By 6 hours after treatment, muscle tone in the 8 and 16 mg tizanidine groups was indistinguishable from muscle tone in placebo treated patients. Within a given patient, improvement in muscle tone was correlated with plasma concentration. Plasma concentrations were variable from patient to patient at a given dose. Although 16 mg produced a larger effect, adverse events including hypotension were more common and more severe than in the 8 mg group. There were no differences in the number of spasms occurring in each group.




**Figure 2: Single Dose Study - Mean Change in Muscle Tone from Baseline as Measured by the Ashworth Scale  $\pm$  95% Confidence Interval (A Negative Ashworth Score Signifies an Improvement in Muscle Tone**

from Baseline)

In a multiple dose study, 118 patients with spasticity secondary to spinal cord injury were randomized to either placebo or tizanidine. Steps similar to those taken in the first study were employed to ensure the integrity of blinding.

Patients were titrated over 3 weeks up to a maximum tolerated dose or 36 mg daily given in three unequal doses (e.g., 10 mg given in the morning and afternoon and 16 mg given at night). Patients were then maintained on their maximally tolerated dose for 4 additional weeks (i.e., maintenance phase). Throughout the maintenance phase, muscle tone was assessed on the Ashworth scale within a period of 2.5 hours following either the morning or afternoon dose. The number of daytime spasms was recorded daily by patients.

At endpoint (the protocol-specified time of outcome assessment), there was a statistically significant reduction in muscle tone and frequency of spasms in the tizanidine treated group compared to placebo. The reduction in muscle tone was not associated with a reduction in muscle strength (a desirable outcome) but also did not lead to any consistent advantage of tizanidine treated patients on measures of activities of daily living. Figure 3 below shows a comparison of the mean change in muscle tone from baseline as measured by the Ashworth scale.



**Figure 3: Multiple Dose Study - Mean Change in Muscle Tone 0.5-2.5 Hours after Dosing as Measured by the Ashworth Scale ± 95% Confidence Interval (A Negative Ashworth Score Signifies an Improvement in Muscle Tone from Baseline)**

## INDICATIONS AND USAGE

Tizanidine is a short-acting drug for the management of spasticity. Because of the short duration of effect, treatment with tizanidine should be reserved for those daily activities and times when relief of spasticity is most important (see **DOSAGE AND ADMINISTRATION**).

## CONTRAINDICATIONS

Tizanidine is contraindicated in patients with known hypersensitivity to tizanidine or its ingredients.

## WARNINGS

### LIMITED DATA BASE FOR CHRONIC USE OF SINGLE DOSES ABOVE 8 MG AND MULTIPLE DOSES ABOVE 24 mg PER DAY

Clinical experience with long-term use of tizanidine at doses of 8 to 16 mg single doses or total daily doses of 24 to 36 mg (see DOSAGE AND ADMINISTRATION) is limited. In safety studies, approximately 75 patients have been exposed to individual doses of 12 mg or more for at least one year or more and approximately 80 patients have been exposed to total daily doses of 30 to 36 mg/day for at least one year or more. There is essentially no long-term experience with single, daytime doses of 16 mg. Because long-term clinical study experience at high doses is limited, only those adverse events with a relatively high incidence are likely to have been identified (see **WARNINGS, PRECAUTIONS** and **ADVERSE REACTIONS**).

### Hypotension

Tizanidine is an  $\alpha_2$ -adrenergic agonist (like clonidine) and can produce hypotension. In a single dose study where blood pressure was monitored closely after dosing, two-thirds of patients treated with 8 mg of tizanidine had a 20% reduction in either the diastolic or systolic BP. The reduction was seen within 1 hour after dosing, peaked 2 to 3 hours after dosing and was associated, at times, with bradycardia, orthostatic hypotension, lightheadedness/dizziness and rarely syncope. The hypotensive effect is dose related and has been measured following single doses of  $\geq 2$  mg.

The chance of significant hypotension may possibly be minimized by titration of the dose and by focusing attention on signs and symptoms of hypotension prior to dose advancement. In addition, patients moving from a supine to a fixed upright position may be at increased risk for hypotension and orthostatic effects.

Caution is advised when tizanidine is to be used in patients receiving concurrent antihypertensive therapy and should not be used with other  $\alpha_2$ -adrenergic agonists.

### Risk of Liver Injury

Tizanidine occasionally causes liver injury, most often hepatocellular in type. In controlled clinical studies, approximately 5% of patients treated with tizanidine had elevations of liver function tests (ALT/SGPT, AST/SGOT) to greater than 3 times the upper limit of normal (or 2 times if baseline levels were elevated) compared to 0.4% in the control patients. Most cases resolved rapidly upon drug withdrawal with no reported residual problems. In occasional symptomatic cases, nausea, vomiting, anorexia and jaundice have been reported. In postmarketing experience, three deaths associated with liver failure have been reported in patients treated with tizanidine. In one case, a 49 year-old male developed jaundice and liver enlargement following 2 months of tizanidine treatment, primarily at 6 mg tid. A liver biopsy showed multilobular necrosis without eosinophilic infiltration. Treatment was discontinued and the patient died in hepatic coma 10 days later. There was no evidence of hepatitis B and C in this patient and other therapy included only oxazepam and ranitidine. There was thus no explanation, other than a reaction to tizanidine, to explain the liver injury. In the two other cases, patients were taking other drugs with known potential for liver toxicity. One patient, treated with tizanidine at a dose of 4 mg/day, was also on carbamazepine when he developed cholestatic jaundice after 2 months of treatment; this patient died with pneumonia about 20 days later. Another patient, treated with tizanidine for 11 days, was also treated with dantrolene for about 2 weeks prior to developing fatal fulminant hepatic failure.

Monitoring of aminotransferase levels is recommended during the first 6 months of treatment (e.g., baseline, 1, 3 and 6 months) and periodically thereafter, based on clinical status. Because of the

potential toxic hepatic effect of tizanidine, the drug should be used only with extreme caution in patients with impaired hepatic function.

## **Sedation**

In the multiple dose, controlled clinical studies, 48% of patients receiving any dose of tizanidine reported sedation as an adverse event. In 10% of these cases, the sedation was rated as severe compared to < 1% in the placebo treated patients. Sedation may interfere with everyday activity.

The effect appears to be dose related. In a single dose study, 92% of the patients receiving 16 mg, when asked, reported that they were drowsy during the 6 hour study. This compares to 76% of the patients on 8 mg and 35% of the patients on placebo. Patients began noting this effect 30 minutes following dosing. The effect peaked 1.5 hours following dosing. Of the patients who received a single dose of 16 mg, 51% continued to report drowsiness 6 hours following dosing compared to 13% in the patients receiving placebo or 8 mg of tizanidine.

In the multiple dose studies, the prevalence of patients with sedation peaked following the first week of titration and then remained stable for the duration of the maintenance phase of the study.

## **Hallucinosi s/Psychotic-Like Symptoms**

Tizanidine use has been associated with hallucinations. Formed, visual hallucinations or delusions have been reported in 5 of 170 patients (3%) in two North American controlled clinical studies. These 5 cases occurred within the first 6 weeks. Most of the patients were aware that the events were unreal. One patient developed psychoses in association with the hallucinations. One patient among these 5 continued to have problems for at least 2 weeks following discontinuation of tizanidine.

## **PRECAUTIONS**

### **Cardiovascular**

Prolongation of the QT interval and bradycardia were noted in chronic toxicity studies in dogs at doses equal to the maximum human dose on a mg/m<sup>2</sup> basis. ECG evaluation was not performed in the controlled clinical studies. Reduction in pulse rate has been noted in association with decreases in blood pressure in the single dose controlled study (see **WARNINGS**).

### **Ophthalmic**

Dose-related retinal degeneration and corneal opacities have been found in animal studies at doses equivalent to approximately the maximum recommended dose on a mg/m<sup>2</sup> basis. There have been no reports of corneal opacities or retinal degeneration in the clinical studies.

### **Use in Renally Impaired Patients**

Tizanidine should be used with caution in patients with renal insufficiency (creatinine clearance < 25 mL/min), as clearance is reduced by more than 50%. In these patients, during titration, the individual doses should be reduced. If higher doses are required, individual doses rather than dosing frequency should be increased. These patients should be monitored closely for the onset or increase in severity of the common adverse events (dry mouth, somnolence, asthenia and dizziness) as indicators of potential overdose.

### **Use in Women Taking Oral Contraceptives**

Tizanidine should be used with caution in women taking oral contraceptives, as clearance of tizanidine is reduced by approximately 50% in such patients. In these patients, during titration, the individual doses should be reduced.

### **Discontinuing Therapy**

If therapy needs to be discontinued, especially in patients who have been receiving high doses for long periods, the dose should be decreased slowly to minimize the risk of withdrawal and rebound hypertension, tachycardia and hypertonia.

### **Information for Patients**

Patients should be advised of the limited clinical experience with tizanidine both in regard to duration of use and the higher doses required to reduce muscle tone (see **WARNINGS**).

Because of the possibility of tizanidine lowering blood pressure, patients should be warned about the risk of clinically significant orthostatic hypotension (see **WARNINGS**).

Because of the possibility of sedation, patients should be warned about performing activities requiring alertness, such as driving a vehicle or operating machinery (see **WARNINGS**). Patients should also be instructed that the sedation may be additive when tizanidine is taken in conjunction with drugs (baclofen, benzodiazepines) or substances (e.g., alcohol) that act as CNS depressants.

Patients should be advised of the change in the absorption profile of tizanidine if taken with food and the potential changes in efficacy and adverse effect profiles that may result (see Pharmacokinetics).

Patients should be advised not to stop tizanidine suddenly as rebound hypertension and tachycardia may occur (see **PRECAUTION, Discontinuing Therapy**).

Tizanidine should be used with caution where spasticity is utilized to sustain posture and balance in locomotion or whenever spasticity is utilized to obtain increased function.

### **Drug Interactions**

In vitro studies of cytochrome P450 isoenzymes using human liver microsomes indicate that neither tizanidine nor the major metabolites are likely to affect the metabolism of other drugs metabolized by cytochrome P450 isoenzymes.

#### **Acetaminophen**

Tizanidine delayed the  $T_{max}$  of acetaminophen by 16 minutes. Acetaminophen did not affect the pharmacokinetics of tizanidine.

#### **Alcohol**

Alcohol increased the AUC of tizanidine by approximately 20% while also increasing its  $C_{max}$  by approximately 15%. This was associated with an increase in side effects of tizanidine. The CNS depressant effects of tizanidine and alcohol are additive.

#### **Oral Contraceptives**

No specific pharmacokinetic study was conducted to investigate interaction between oral contraceptives and tizanidine, but retrospective analysis of population pharmacokinetic data following single and multiple dose administration of 4 mg tizanidine showed that women concurrently taking oral contraceptives had 50% lower clearance of tizanidine than women not on oral contraceptives.

#### **Rofecoxib**

Rofecoxib may potentiate the adverse effects of tizanidine. Eight case reports of a potential rofecoxib-tizanidine drug interaction have been identified in postmarketing safety reports. Most of the adverse events reported involved the nervous system (e.g., hallucinations, psychosis, somnolence, hypotonia, etc.) and the cardiovascular system (e.g., hypotension, tachycardia, bradycardia). In all cases, adverse events resolved following discontinuation of tizanidine, rofecoxib, or both. Rechallenges with both drugs were not performed. The possible mechanism and the potential for a drug interaction between tizanidine and rofecoxib remain unclear.

## **Carinogenesis, Mutagenesis, Impairment of Fertility**

No evidence for carcinogenicity was seen in two dietary studies in rodents. Tizanidine was administered to mice for 78 weeks at doses up to 16 mg/kg, which is equivalent to 2 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. Tizanidine was also administered to rats for 104 weeks at doses up to 9 mg/kg, which is equivalent to 2.5 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. There was no statistically significant increase in tumors in either species.

Tizanidine was not mutagenic or clastogenic in the following *in vitro* assays: the bacterial Ames test and the mammalian gene mutation test and chromosomal aberration test in Chinese hamster cells. It was also negative in the following *in vivo* assays: the bone marrow micronucleus test in mice, the bone marrow micronucleus and cytogenicity test in Chinese hamsters, the dominant lethal mutagenicity test in mice and the unscheduled DNA synthesis (UDS) test in mice.

Tizanidine did not affect fertility in male rats at doses of 10 mg/kg, approximately 2.7 times the maximum recommended human dose on a mg/m<sup>2</sup> basis and in females at doses of 3 mg/kg, approximately equal to the maximum recommended human dose on a mg/m<sup>2</sup> basis; fertility was reduced in males receiving 30 mg/kg (8 times the maximum recommended human dose on a mg/m<sup>2</sup> basis) and in females receiving 10 mg/kg (2.7 times the maximum recommended human dose on a mg/m<sup>2</sup> basis). At these doses, maternal behavioral effects and clinical signs were observed including marked sedation, weight loss and ataxia.

## **Pregnancy**

### **Pregnancy Category C**

Reproduction studies performed in rats at a dose of 3 mg/kg, equal to the maximum recommended human dose on a mg/m<sup>2</sup> basis, and in rabbits at 30 mg/kg, 16 times the maximum recommended human dose on a mg/m<sup>2</sup> basis, did not show evidence of teratogenicity. Tizanidine at doses that are equal to and up to 8 times the maximum recommended human dose on a mg/m<sup>2</sup> basis increased gestation duration in rats. Prenatal and postnatal pup loss was increased and developmental retardation occurred. Postimplantation loss was increased in rabbits at doses of 1 mg/kg or greater, equal to or greater than 0.5 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. Tizanidine has not been studied in pregnant women. Tizanidine should be given to pregnant women only if clearly needed.

## **Labor and Delivery**

The effect of tizanidine on labor and delivery in humans is unknown.

## **Nursing Mothers**

It is not known whether tizanidine is excreted in human milk, although as a lipid soluble drug, it might be expected to pass into breast milk.

## **Geriatric Use**

Tizanidine should be used with caution in elderly patients because clearance is decreased four-fold.

## **Pediatric Use**

There are no adequate and well-controlled studies to document the safety and efficacy of tizanidine in children.

## **ADVERSE REACTIONS**

In multiple dose, placebo-controlled clinical studies, 264 patients were treated with tizanidine and 261 with placebo. Adverse events, including severe adverse events, were more frequently reported with tizanidine than with placebo.

## Common Adverse Events Leading to Discontinuation

Forty-five of 264 (17%) patients receiving tizanidine and 13 of 261 (5%) patients receiving placebo in three multiple dose, placebo-controlled clinical studies discontinued treatment for adverse events. When patients withdrew from the study, they frequently had more than one reason for discontinuing. The adverse events most frequently leading to withdrawal of tizanidine treated patients in the controlled clinical studies were asthenia (weakness, fatigue and/or tiredness) (3%), somnolence (3%), dry mouth (3%), increased spasm or tone (2%) and dizziness (2%).

## Most Frequent Adverse Clinical Events Seen in Association with the Use of Tizanidine

In multiple dose, placebo-controlled clinical studies involving 264 patients with spasticity, the most frequent adverse events were dry mouth, somnolence/sedation, asthenia (weakness, fatigue and/or tiredness) and dizziness. Three-quarters of the patients rated the events as mild to moderate and one-quarter of the patients rated the events as being severe. These events appeared to be dose related.

## Adverse Events Reported in Controlled Studies

The events cited reflect experience gained under closely monitored conditions of clinical studies in a highly selected patient population. In actual clinical practice or in other clinical studies, these frequency estimates may not apply, as the conditions of use, reporting behavior and the kinds of patients treated may differ. Table 1 lists treatment emergent signs and symptoms that were reported in greater than 2% of patients in three multiple dose, placebo-controlled studies who received tizanidine where the frequency in the tizanidine group was at least as common as in the placebo group. These events are not necessarily related to tizanidine treatment. For comparison purposes, the corresponding frequency of the event (per 100 patients) among placebo treated patients is also provided.

**Table 1: Multiple Dose, Placebo-Controlled Studies - Frequent (> 2%) Adverse Events Reported for Which Tizanidine Incidence is Greater Than Placebo**

Event	Placebo N=261 %	Tizanidine Hydrochloride Tablet N=264 %
Dry Mouth	10	49
Somnolence	10	48
Asthenia	16	41
Dizziness	4	16
UTI	7	10
Infection	5	6
Constipation	1	4
Liver function tests abnormal	1	3
Vomiting	0	3
Speech disorder	0	3
Amblyopia (blurred vision)	<1	3
Urinary frequency	2	3
Flu symptom	2	3
SGPT/ALT increased	<1	3
Dyskinesia	0	3
Nervousness	<1	3
Pharyngitis	1	3
Rhinitis	2	3

In the single dose, placebo-controlled study involving 142 patients with spasticity, the patients were specifically asked if they had experienced any of the four most common adverse events: dry mouth, somnolence (drowsiness), asthenia (weakness, fatigue and/or tiredness) and dizziness. In addition, hypotension and bradycardia were observed. The occurrence of these adverse events are summarized in Table 2. Other events were, in general, reported at a rate of 2% or less.

**Table 2: Single Dose, Placebo-Controlled Study - Common Adverse Events Reported**

<b>Event</b>	<b>Placebo N=48 %</b>	<b>Tizanidine Hydrochloride Tablets 8 mg N=45 %</b>	<b>Tizanidine Hydrochloride Tablets 16 mg N=49 %</b>
Somnolence	31	78	92
Dry mouth	35	76	88
Asthenia	40	67	78
Dizziness	4	22	45
Hypotension	0	16	33
Bradycardia	0	2	10

### **Other Adverse Events Observed During the Evaluation of Tizanidine**

Tizanidine was administered to 1385 patients in additional clinical studies where adverse event information was available. The conditions and duration of exposure varied greatly, and included (in overlapping categories) double-blind and open-label studies, uncontrolled and controlled studies, inpatient and outpatient studies, and titration studies. Untoward events associated with this exposure were recorded by clinical investigators using terminology of their own choosing. Consequently, it is not possible to provide a meaningful estimate of the proportion of individuals experiencing adverse events without first grouping similar types of untoward events into a smaller number of standardized event categories.

In the tabulations that follow, reported adverse events were classified using a standard COSTART-based dictionary terminology. The frequencies presented, therefore, represent the proportion of the 1385 patients exposed to tizanidine who experienced an event of the type cited on at least one occasion while receiving tizanidine. All reported events are included except those already listed in Table 1. If the COSTART term for an event was so general as to be uninformative, it was replaced with a more informative term. It is important to emphasize that, although the events reported occurred during treatment with tizanidine, they were not necessarily caused by it.

Events are further characterized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring on one or more occasions in at least 1/100 patients (only those not already listed in the tabulated results from placebo-controlled studies appear in this listing); infrequent adverse events are those occurring in 1/100 to 1/1000 patients.

Body as a Whole: *Frequent*: fever; *Infrequent*: allergic reaction, moniliasis, malaise, abscess, neck pain, sepsis, cellulitis, death, overdose; *Rare*: carcinoma, congenital anomaly, suicide attempt.

Cardiovascular System: *Infrequent*: vasodilatation, postural hypotension, syncope, migraine, arrhythmia; *Rare*: angina pectoris, coronary artery disorder, heart failure, myocardial infarct, phlebitis, pulmonary embolus, ventricular extrasystoles, ventricular tachycardia.

Digestive System: *Frequent*: abdomen pain, diarrhea, dyspepsia; *Infrequent*: dysphagia, cholelithiasis, fecal impaction, flatulence, gastrointestinal hemorrhage, hepatitis, melena; *Rare*: gastroenteritis, hematemesis, hepatoma, intestinal obstruction, liver damage.

Hemic and Lymphatic System: *Infrequent*: ecchymosis, hypercholesteremia, anemia, hyperlipemia,

leukopenia, leukocytosis, sepsis; *Rare*: petechia, purpura, thrombocythemia, thrombocytopenia.

Metabolic and Nutritional System: *Infrequent*: edema, hypothyroidism, weight loss; *Rare*: adrenal cortex insufficiency, hyperglycemia, hypokalemia, hyponatremia, hypoproteinemia, respiratory acidosis.

Musculoskeletal System: *Frequent*: myasthenia, back pain; *Infrequent*: pathological fracture, arthralgia, arthritis, bursitis.

Nervous System: *Frequent*: depression, anxiety, paresthesia; *Infrequent*: tremor, emotional lability, convulsion, paralysis, thinking abnormal, vertigo, abnormal dreams, agitation, depersonalization, euphoria, migraine, stupor, dysautonomia, neuralgia; *Rare*: dementia, hemiplegia, neuropathy.

Respiratory System: *Infrequent*: sinusitis, pneumonia, bronchitis; *Rare*: asthma.

Skin and Appendages: *Frequent*: rash, sweating, skin ulcer; *Infrequent*: pruritus, dry skin, acne, alopecia, urticaria; *Rare*: exfoliative dermatitis, herpes simplex, herpes zoster, skin carcinoma.

Special Senses: *Infrequent*: ear pain, tinnitus, deafness, glaucoma, conjunctivitis, eye pain, optic neuritis, otitis media, retinal hemorrhage, visual field defect; *Rare*: iritis, keratitis, optic atrophy.

Urogenital System: *Infrequent*: urinary urgency, cystitis, menorrhagia, pyelonephritis, urinary retention, kidney calculus, uterine fibroids enlarged, vaginal moniliasis, vaginitis; *Rare*: albuminuria, glycosuria, hematuria, metrorrhagia.

## **DRUG ABUSE AND DEPENDENCE**

Abuse potential was not evaluated in human studies. Rats were able to distinguish tizanidine from saline in a standard discrimination paradigm, after training, but failed to generalize the effects of morphine, cocaine, diazepam or phenobarbital to tizanidine. Monkeys were shown to self-administer tizanidine in a dose-dependent manner, and abrupt cessation of tizanidine produced transient signs of withdrawal at doses > 35 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. These transient withdrawal signs (increased locomotion, body twitching, and aversive behavior toward the observer) were not reversed by naloxone administration.

Tizanidine is closely related to clonidine, which is often abused in combination with narcotics and is known to cause symptoms of rebound upon abrupt withdrawal. Three cases of rebound symptoms on sudden withdrawal of tizanidine have been reported. The case reports suggest that these patients were also misusing narcotics. Withdrawal symptoms included hypertension, tachycardia, hypertonia, tremor, and anxiety. As with clonidine, withdrawal is expected to be more likely in cases where high doses are used, especially for prolonged periods.

## **OVERDOSAGE**

A search of a safety surveillance database revealed a total of eighteen cases of tizanidine overdose. Of the fourteen intentional overdoses, five have resulted in fatality, and in at least three of these cases, other CNS depressants were involved. One fatality was secondary to pneumonia and sepsis, which were sequelae of the ingestion. The majority of cases involve depressed consciousness (somnolence, stupor or coma), depressed cardiovascular function (bradycardia, hypotension) and depressed respiratory function (respiratory depression or failure).

Should overdose occur, basic steps to ensure the adequacy of an airway and the monitoring of cardiovascular and respiratory systems should be undertaken. In general, symptoms resolve within one to three days following discontinuation of tizanidine and administration of appropriate therapy. Due to the similar mechanism of action, symptoms and management of tizanidine overdose are similar to those following clonidine overdose. For the most recent information concerning the management of overdose, contact a poison control center.

## DOSAGE AND ADMINISTRATION

A single oral dose of 8 mg of tizanidine reduces muscle tone in patients with spasticity for a period of several hours. The effect peaks at approximately 1 to 2 hours and dissipates between 3 to 6 hours. Effects are dose-related.

Although single doses of less than 8 mg have not been demonstrated to be effective in controlled clinical studies, the dose-related nature of tizanidine's common adverse events make it prudent to begin treatment with single oral doses of 4 mg. Increase the dose gradually (2 mg to 4 mg steps) to optimum effect (satisfactory reduction of muscle tone at a tolerated dose).

The dose can be repeated at 6 to 8 hour intervals, as needed, to a maximum of three doses in 24 hours. The total daily dose should not exceed 36 mg.

Experience with single doses exceeding 8 mg and daily doses exceeding 24 mg is limited. There is essentially no experience with repeated, single, daytime doses greater than 12 mg or total daily doses greater than 36 mg (see **WARNINGS**).

Food has complex effects on tizanidine pharmacokinetics. These pharmacokinetic differences may result in clinically significant differences when switching administration of the tablet between the fed and fasted state. These changes may result in increased adverse events or delayed/more rapid onset of activity, depending upon the nature of the switch. For this reason, the prescriber should be thoroughly familiar with the changes in kinetics associated with these different conditions (see Pharmacokinetics).

## HOW SUPPLIED

**Tizanidine Hydrochloride Tablets, 2 mg (base)** are supplied as white, round, flat faced, beveled edge, debossed "E" over "34" on one side and bisected on the other side and are available in:

Bottles of 28 42549-549-28  
Bottles of 30 42549-549-30  
Bottles of 60 42549-549-60  
Bottles of 90 42549-549-90  
Bottles of 120 42549-549-12  
Bottles of 140 42549-549-04

**Tizanidine Hydrochloride Tablets, 4 mg (base)** are supplied as white, round, flat faced, beveled edge, debossed "E" over "44" on one side and quadrisected on the other side and are available in:

Bottles of 28 42549-597-28  
Bottles of 30 42549-597-30  
Bottles of 60 42549-597-60  
Bottles of 90 42549-597-90  
Bottles of 120 42549-597-12  
Bottles of 140 42549-597-04

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Dispense contents in a tight, light-resistant container as defined in the USP with a child-resistant closure, as required.

**To report SUSPECTED ADVERSE REACTIONS, contact Sandoz Inc. at 1-800-525-8747 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**2 mg and 4 mg**

Manufactured by  
Epic Pharma, LLC  
Laurelton, NY 11413  
for Sandoz Inc.  
Princeton, NJ 08540

**4 mg**

Manufactured by  
Sandoz Inc.  
Princeton, NJ 08540  
OS8800  
Rev. 02/09  
MF0034REV02/09

**Relabeling and Repackaging by:**

STAT Rx USA LLC  
Gainesville, GA 30501

**PRINCIPAL DISPLAY PANEL - 2MG**

**Tizanidine 2mg tablet(s)**

**RX only**

Packaged and Distributed By:  STAT Rx USA Gainesville, GA 30501

**2mg Tizanidine 30 Tabs**

Generic For: **Zanaflex**

**NDC 42549-549-30** Prod# 549-30  
Lot# SAMPLE

\* Each Tablet Contains: 2.288mg Tizanidine Hcl.  
equivalent to 2mg tizanidine base.

Mfg By: Sandoz Inc. NDC 0185-0034-51  
Princeton, NJ 08540

Mfg Lot: 27384  
Discard After: 05/15 BW 12/1/2011 SAMPLE

**RX ONLY-KEEP OUT OF REACH OF CHILDREN**

0

BY350

Dosage: See package insert  
Store between 59-86 degrees F

**PRINCIPAL DISPLAY PANEL - 4MG**

**Tizanidine 4mg tablet(s)**

**RX only**

Packaged and Distributed By: **STAT USA** Gainesville, GA 30501

**Tizanidine**  
4mg **60 Tabs**

Generic For: **Zanaflex**

NDC 42549-597-60 Prod# 597-60 Lot# SAMPLE  
\*Each Tablet Contains: 4.576mg Tizanidine Hcl. equivalent to 4mg tizanidine base.

Mfg By: Sandoz Inc. Princeton, NJ 08540 NDC 0185-4400-51  
Mfg Lot: GKK0900 Discard After: 03/13 BW 11/29/2011 SAMPLE

**RX ONLY-KEEP OUT OF REACH OF CHILDREN**

May cause **DROWSINESS**. **ALCOHOL** may intensify this effect. Use care when operating dangerous machinery. May cause drowsiness or dizziness.

**TIZANIDINE HYDROCHLORIDE**

tizanidine hydrochloride tablet

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42549-549(NDC:0185-0034)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
TIZANIDINE HYDROCHLORIDE (UNII: B53E3NMY5C) (TIZANIDINE - UNII:6A106C00GW)	TIZANIDINE	2 mg

**Inactive Ingredients**

Ingredient Name	Strength
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	7mm
Flavor		Imprint Code	E;34
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42549-549-04	140 in 1 BOTTLE		
2	NDC:42549-549-12	120 in 1 BOTTLE		
3	NDC:42549-549-28	28 in 1 BOTTLE		
4	NDC:42549-549-30	30 in 1 BOTTLE		
5	NDC:42549-549-60	60 in 1 BOTTLE		
6	NDC:42549-549-90	90 in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076280	11/26/2002	

## TIZANIDINE HYDROCHLORIDE

tizanidine hydrochloride tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42549-597(NDC:0185-4400)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TIZANIDINE HYDROCHLORIDE (UNII: B53E3NMY5C) (TIZANIDINE - UNII:6A106C00GW)	TIZANIDINE	4 mg

### Inactive Ingredients

Ingredient Name	Strength
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	4 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	E;44
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42549-597-04	140 in 1 BOTTLE		
2	NDC:42549-597-12	120 in 1 BOTTLE		
3	NDC:42549-597-28	28 in 1 BOTTLE		
4	NDC:42549-597-30	30 in 1 BOTTLE		
5	NDC:42549-597-60	60 in 1 BOTTLE		
6	NDC:42549-597-90	90 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076280	06/27/2002	

**Labeler** - STAT Rx USA LLC (786036330)

**Registrant** - PSS World Medical Inc. (101822682)

### Establishment

Name	Address	ID/FEI	Business Operations
STAT Rx USA LLC		786036330	RELABEL, REPACK