

ANTI-DIARRHEAL- loperamide hydrochloride capsule, liquid filled
MEIJER DISTRIBUTION INC

Drug Facts

Active ingredient (in each capsule)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- in children under 12 years of age

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea.**
- **not for use in children under 12 years of age**
- adults and children 12 years and over: 2 capsules after the first loose stool; 1 capsule after each subsequent loose stool; but no more than 4 capsules in 24 hours

Other information

- store at 20°-25°C (68°-77°F). Protect from Light.
- avoid excessive heat above 40°C (104°F)
- do not use if carton or blister unit is open or torn
- see side panel for lot number and expiration date

Inactive ingredients

butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water

Questions or comments?

call toll free **1-888-235-2466** (Mon - Fri 9AM - 5PM EST)

DIST. MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

†This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Imodium[®] A-D.

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

L0000305

Rev # 11/19

Principal Display Panel

meijer

NDC 41250-210-08

Compare to Imodium[®] A-D[†]

anti-diarrheal

Loperamide HCl Soft Gelatin Capsules | 2 mg

Controls the Symptoms of Diarrhea

Suitable for adults and children 12 years and over

12 Softgels*

*soft gelatin capsules

Each softgel contains 2 mg loperamide HCl

actual size

PRINT SIDE GLUE

NO INK NO VARNISH

Drug Facts

Active ingredient (in each capsule) Loperamide HCl 2 mg..... Anti-diarrheal Purpose

Use controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings
 Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
 Heart alert: Taking more than directed can cause serious heart problems or death
 Do not use if you have bloody or black stool or in children under 12 years of age
 Ask a doctor before use if you have fever, mucus in the stool, a history of liver disease, a history of abnormal heart rhythm, or a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, dizziness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bloating, these may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.


Directions
 Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
 Adults and children 12 years and over: 2 capsules after the first loose stool, 1 capsule after each subsequent loose stool, but no more than 4 capsules in 24 hours.
 Children under 12 years of age: not for use.

Other information
 Store at 20°-25°C (68°-77°F). Protect from light.
 Avoid excessive heat above 40°C (104°F).
 Do not use if carton or blister unit is open or torn. See side panel for lot number and expiration date.

Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxy 40 hydrogenated castor oil, purified water.


Questions or comments? call toll free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

NO INK NO VARNISH



anti-diarrheal
Loperamide HCl Soft Gelatin Capsules | 2 mg


NDC 41250-210-08
Compare to
Imodium® A-D



anti-diarrheal
Loperamide HCl Soft Gelatin Capsules | 2 mg

Controls the Symptoms of Diarrhea
Suitable for adults and children 12 years and over


12 Softgels*
*soft gelatin capsules
Each softgel contains 2 mg loperamide HCl



actual size

DIST. MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com

PID 4636635



7 19283 61840 5

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT EVIDENT PACKAGE. USE ONLY IF BUSTERS & KEYS ARE USED TO OPEN. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

THIS PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED BY JOHNSON & JOHNSON, OWNER OF THE REGISTERED TRADEMARK

5



ANTI-DIARRHEAL

loperamide hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-210
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
GLYCERYL CAPRYLATE (UNII: TM2TZD4G4A)	
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)	
WATER (UNII: 059QF0K00R)	

Product Characteristics

Color	blue	Score	no score
Shape	CAPSULE	Size	10mm
Flavor		Imprint Code	LP2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-210-12	2 in 1 CARTON	02/01/2019	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41250-210-08	1 in 1 CARTON	01/13/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021855	02/01/2019	

Labeler - MEIJER DISTRIBUTION INC (006959555)**Registrant** - Bionpharma Inc. (079637826)**Establishment**

Name	Address	ID/FEI	Business Operations
Patheon Softgels Inc.		002193829	manufacture(41250-210)

Revised: 1/2020

MEIJER DISTRIBUTION INC