

COLD MEDICINE- cold medicine tablet
Allegiant Health

GreenField - Cold Medicine

Active ingredient(s)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Nasal decongestant

Use(s)

temporarily relieves

- minor aches and pains
- headaches
- nasal congestion
- sinus congestion & pressure
- runny nose
- sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 10 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- if you are allergic to acetaminophen
- more than 10 days unless directed by a doctor

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.

If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this

product

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not

sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

you have

- liver disease
- heart disease
- glaucoma
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if

- are taking the blood thinning drug warfarin
- are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- may cause
- drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- redness or swelling is present
- new symptoms occur
- you get nervous, dizzy, or sleepless
- pain or nasal congestion gets worse or lasts for more than 7 days
- fever gets worse or lasts for more than 3 days

Pregnancy/Breastfeeding

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Do not use more than directed
- **Adults and children 12 years of age and older:** take 2 tablets every 4 to 6 hours
Do not take more than 10 tablets in 24 hours
- **Children under 12 years of age:** do not use this product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage.

Other information

- store at room temperature in a dry place
- do not use if blister package is torn.

Inactive ingredients

corn starch, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid

Questions

1-888-952-0050 Monday through Friday 9 AM - 5 PM EST.

Principal Display Panel

Sinus

DAYTIME

Compare to Tylenol Sinus® Active Ingredient*

Well Fine Medus

See New Warnings Information

Sinus Cold Congestion & Pain


Well Fine Medus

Lot:
Exp:

This product is not manufactured or distributed by Novartis Consumer Health, a division of Novartis Corporation, or its subsidiaries. It is manufactured by SLD, LLC, a subsidiary of Novartis Consumer Health, located at 2000 North 1st Street, Columbus, IN 47201.

Manufactured by: SLD, LLC
2000 North 1st Street
Columbus, IN 47201

Lot: L81949 R0721
S/N: 52510 00580 4



Drug Facts
Active ingredient (in each tablet)
Chlorpheniramine maleate 2 mg
Acetaminophen 325 mg

Purposes
Relieves nasal congestion, sneezing, runny nose, and watery eyes. Relieves pain and fever.

Warnings
User warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 10 tablets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen. 3 or more alcoholic drinks every day while using this product.

Directions
Do not use more than directed. Adults and children 12 years of age and older: Take 2 tablets every 4 to 6 hours. Do not take more than 10 tablets in 24 hours. Children under 12 years of age: Do not use this product in children under 12 years of age. This will provide more than the recommended dose (overdose) and may cause liver damage.

Other information
Store at room temperature in a dry place. Do not use if blister package is torn.

Inactive ingredients
Contains: starch, croscarmellose, hydroxypropylcellulose, polyethylene glycol, polyvinylpyrrolidone, polyethylene glycol succinate, stearic acid.

Questions and Comments?
1-800-858-5000 Monday through Friday 9AM - 5PM EST.

Drug Facts (continued)
When using this product, do not use more than directed. Acetaminophen may occur, especially in children, may cause drowsiness, dizziness, and drowsiness may increase. Avoid alcoholic drinks. Be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if: redness or swelling is present, new symptoms occur, you get hives, dizziness, or drowsiness, pain or nasal congestion gets worse or lasts for more than 7 days, or your eyes, nose or ears for more than 3 days.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Think medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

LFBG7640

COLD MEDICINE

cold medicine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69168-004
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPROVIDONE (UNII: 2S7830E561)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	AZ 275
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-004-51	20 in 1 CARTON; Type 0: Not a Combination Product	11/23/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/12/2021	

