

CHILDREN PAIN RELIEF- children pain relief tablet, chewable
Allegiant Health

440 - Children Pain Relief

Active ingredient(s)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Use(s)

- temporarily relieves minor aches and pains due to:
- headache
- sore throat
- flu
- toothache
- the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if

your child has liver disease.

Ask a doctor or pharmacist before use if

your child is taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

this product does not contain directions or complete warnings for adult use

- do not give more than directed
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew before swallowing; this product will soften in mouth for ease of chewing
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (chewable tablets)**
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

**or as directed by a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions/Comments

Call 1-888-952-0050 Monday through Friday 9AM - 5PM EST

Principal Display Panel

HealthA2Z® Compare to Children's Tylenol® active ingredient* NDC 69168-440-60



Children's
Ages 2-11

PAIN RELIEF

Acetaminophen 160 mg per tablet
Pain Reliever-Fever Reducer

ASPIRIN FREE
IBUPROFEN FREE

Grape Flavor Chewables
Chew or crush tablets completely before swallowing



60 Chewable Tablets

Drug Facts

Active ingredient (in each chewable tablet) Acetaminophen 160 mg.....Pain reliever/fever reducer

Purpose Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ sore throat ■ flu ■ toothache ■ the common cold ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

Do not use if a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use if you are not sure whether a drug contains acetaminophen (prescription or over-the-counter or pharmacist). ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Drug Facts (continued on inside)

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Children's Tylenol®

Distributed by:
Alleghiant Health
Dear Park, NY 11729



LB2092
R0223



X003PNNND6D
HealthA2Z® Children's... Aspirin & Ibuprofen Free
New

Lot: _____
Exp: _____

PEEL HERE →

Drug Facts (continued)

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ find the right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ chew before swallowing; this product will soften in mouth for ease of chewing ■ repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (chewable tablets)**
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

**or as directed by a doctor

Other Information ■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING. ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive Ingredients citric acid, croscollonite, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, FD&C blue #1 aluminum lake, titanium dioxide, magnesium stearate, mannitol, polyethylene, stearic acid, sucrose

Questions or Comments?
Call 1-888-832-0350 Monday through Friday 9AM - 5PM EST

Acetaminophen 160mg

CHILDREN PAIN RELIEF

children pain relief tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69168-440
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

Inactive Ingredients

Ingredient Name	Strength

CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CROSPROVIDONE (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
DEXTRATES (UNII: G263MI44RU)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	purple	Score	2 pieces
Shape	ROUND	Size	16mm
Flavor	GRAPE	Imprint Code	44;449
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-440-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	07/18/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/18/2023	

Labeler - Allegiant Health (079501930)

Revised: 7/2023

Allegiant Health