

**GOOD NEIGHBOR PHARMACY DAY TIME- acetaminophen, dextromethorphan hbr, phenylephrine hcl solution**

**Amerisource Bergen**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Amerisource Bergen Day Time Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product**

**do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

### Other information

- **each 15 mL contains:** sodium 7 mg
- store at 20-25°C (68-77°F)

### Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

### Questions or comments?

**1-800-719-9260**

### Principal Display Panel

Compare to Vicks® DayQuil® Cold & Flu active ingredients

See New Warnings

MULTI-SYMPTOM

Non-Drowsy

Day Time

pain reliever, fever reducer, cough suppressant, nasal decongestant

Cold & Flu

Acetaminophen

Phenylephrine HCl

Dextromethorphan HBr

Aches – Fever – Cough – Nasal Congestion

Alcohol Free / Antihistamine Free

8 fl oz (237 mL)

Do not use if printed neckband is broken or missing

\*Good Neighbor Pharmacy® Day Time is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Cold & Flu.

GLUTEN FREE



Distributed By  
AmerisourceBergen  
1300 Morris Drive  
Chesterbrook, PA 19087  
Questions or Concerns?  
[www.mygnp.com](http://www.mygnp.com)



**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

: 65634 29 F3



Compare to  
Vicks® DayQuil® Cold & Flu  
active ingredients\*

NDC 24385-005-34  
See New Warnings

NO COPY AREA

NO COPY AREA

Non-Drowsy **MULTI-SYMPOM**

# Day Time

pain reliever, fever reducer,  
cough suppressant, nasal decongestant

## Cold & Flu

- Acetaminophen
- Phenylephrine HCl
- Dextromethorphan HBr

• Aches • Fever • Cough • Nasal Congestion

Alcohol Free / Antihistamine Free

8 fl oz (237 mL)

### Drug Facts

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

### Uses

- temporarily relieves common cold/flu symptoms:
- cough due to minor throat and bronchial irritation
- minor aches and pains
- sore throat
- fever
- nasal congestion
- headache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION →

### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** ■ liver disease  
■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure  
■ trouble urinating due to an enlarged prostate gland  
■ cough that occurs with too much phlegm (mucus)  
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

- Stop use and ask a doctor if** ■ new symptoms occur
- you get nervous, dizzy or sleepless
  - redness or swelling is present
  - fever gets worse or lasts more than 3 days
  - pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
  - cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### Drug Facts (continued)

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

■ take only as directed – see Overdose warning  
■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

**Other information** ■ each 15 mL contains: sodium 7 mg  
■ store at 20-25°C (68-77°F)

**Inactive ingredients** butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

**Questions or comments?** 1-800-719-9260

## GOOD NEIGHBOR PHARMACY DAY TIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:24385-005
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL

<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

### Product Characteristics

<b>Color</b>	ORANGE (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MENTHOL (with fruit)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24385-005-30	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/20/2006	
2	NDC:24385-005-38	296 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/21/2007	
3	NDC:24385-005-34	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/21/2012	
4	NDC:24385-005-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/19/2012	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/20/2006	

**Labeler** - Amerisource Bergen (007914906)