

DAYTIME SEVERE COLD AND COUGH- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride powder
Cellchem Pharmaceuticals Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

**Daytime Severe
Cold And Cough**

Drug Facts

Active Ingredients (in each packet)	Purpose
Acetaminophen 650 mg.....	Pain reliever/Fever reducer
Dextromethorphan hydrobromide 20 mg.....	Cough Suppressant
Phenylephrine Hydrochloride 10 mg.....	Nasal decongestant

Uses

- Temporarily relieves these symptoms due to cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Relieves:

- nasal congestion
- sore throat pain
- Cough
- Headache
- Bodyaches
- Fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 packets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in children under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription and non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

If pregnant or breast-feeding, ask a health care professional before use.

When using this product

- do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.

KEEP CARTON FOR REFERENCE.

DO NOT DISCARD.

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN

Directions

- do not use more than directed
- take every 4 hours, while symptoms persist.

Do not take more than 6 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years and over	one packet
children under 12 years	do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other Information

- each packet contains: potassium 10 mg, sodium 24 mg.
- phenylketonurics: contains phenylalanine 14 mg per packet.
- store at controlled room temperature 20-25°C (68 -77°F). Protect from heat and moisture.

Inactive ingredients

Acesulfame potassium, anhydrous citric acid, aspartame, FD&C Blue No.1, FD&C red No.40, maltodextrin, menthol, natural flavors, sodium citrate, starch and sugar.

Questions, Comments or Adverse Reactions?

You may report serious side effects to: +1 844-481-8884

Serious side effects can also be reported to: drugsafety@cellchempharma.ca

Distributed by:

CellChem Pharmaceuticals Inc.

Unit 1, 30 Concourse Gate

Nepean, Ontario K2E 7V7

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

SEAL AREA
INK FREE AREA
ZONE DE COLLAGE
ZONE SANS ENCRE

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

Drugs Facts

Active ingredients (in each packet)	Purpose
Acetaminophen 650 mg	Pain reliever / Fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Phenylephrine Hydrochloride 10 mg	Nasal decongestant

Uses

- Temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount. Do not take with other drugs containing acetaminophen. Do not take 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use in a child under 12 years of age if you are allergic to acetaminophen or with any other drug containing acetaminophen (prescription and non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to an enlarged prostate gland, cough that occurs with too much phlegm (mucus), cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are: taking the blood thinning drug warfarin.

When using this product: do not exceed recommended dosage.

Stop use and ask a doctor if: nervousness, dizziness, or sleeplessness occurs; fever gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; pain, cough or nasal congestion gets worse or lasts more than 7 days; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours, while symptoms persist. Do not take more than 6 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years and over	one packet
children under 12 years	do not use

- dissolve contents of one packet into 8 oz. hot water, sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg, sodium 24 mg.
- phenylephrine: contains phenylalanine 14 mg per packet.
- store at controlled room temperature 20-25°C (68-77°F). Protect from heat and moisture.

Inactive ingredients

Acetosulfame potassium, anhydrous citric acid, aspartame, FD&C Blue No.1, FD&C red No.40, maltodextrin, menthol, natural flavors, sodium citrate, starch and sugar.

NDC 73147-3110-6

Daytime Severe Cold And Cough

Acetaminophen
- Pain reliever / Fever reducer
Dextromethorphan HBr - Cough Suppressant
Phenylephrine HCl - Nasal decongestant

Relieves:
Nasal congestion
Sore throat pain
Cough
Headache
Body ache
Fever



6 Packets Berry Infused with Menthol and Green Tea flavors

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN

You may report serious side effects to:
+1 844-461-8884

Serious side effects can also be reported to:
drugsafety@celchempharma.ca

Distributed by:
CelChem
CelChem Pharmaceuticals Inc.
Unit 1, 30 Concourse Gate
Nepean, Ontario K2E 7V7

SEAL AREA
INK FREE AREA
ZONE DE COLLAGE
ZONE SANS ENCRE

LOT & EXP DD-MM-YYYY



TO OPEN: FOLD CORNER & TEAR AT SLIT OR CUT ALONG DOTTED LINE WITH SCISSORS. DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.



Daytime Severe Cold And Cough

Berry Infused with Menthol and Green Tea flavors

Active ingredients (in each packet)	Purposes
Acetaminophan 650 mg	Pain reliever / Fever reducer
Dextromethorphan hydrobromide 20 mg	cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

SEE ALLERGY AND LIVER WARNINGS ON OTHER SIDE

**THIS PACKET DOES NOT CONTAIN
FULL PRODUCT INFORMATION.
SEE CARTON FOR COMPLETE
LABELLING INFORMATION. READ
ALL WARNINGS AND DIRECTIONS
ON CARTON BEFORE USE,
DO NOT DISCARD CARTON.**

1 SINGLE DOSE

0.36 oz Powder

PC-3110-GENERIC

Distributed by:
Cell Chem
CellChem Pharmaceuticals Inc.
Unit 1, 30 Concourse Gate
Napan, Ontario K2E 7V7

Warnings
The product contains acetaminophen. Severe liver damage may occur if you take:
• more than 6 packets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

*Berry Infused with Menthol
and Green Tea flavors*

**Daytime Severe
Cold And Cough**

DAYTIME SEVERE COLD AND COUGH

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride powder

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:73147-3110

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ASPARTAME (UNII: Z0H242BBR1)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
STARCH, CORN (UNII: O8232NY3SJ)	
RAW SUGAR (UNII: 8M707QY5GH)	
MENTHOL (UNII: L7T10EIP3A)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY (Berry Infused with Menthol and Green Tea flavors)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73147-3110-6	6 in 1 BOX, UNIT-DOSE; Type 0: Not a Combination Product	07/01/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/01/2019	

Labeler - Cellchem Pharmaceuticals Inc (11518618)

Registrant - Cellchem Pharmaceuticals Inc (11518618)

Establishment

Name	Address	ID/FEI	Business Operations
Laboratoires Confab Inc		241754217	manufacture(73147-3110)

Revised: 1/2021

Cellchem Pharmaceuticals Inc