

**TRIPLE ANTIBIOTIC PLUS PAIN RELIEF- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment**  
**Taro Pharmaceuticals U.S.A., Inc.**

-----  
**Triple Antibiotic Plus Pain Relief**

<b><i>Active ingredients (each gram contains)</i></b>	<b><i>Purpose</i></b>
Bacitracin zinc 500 units	First aid antibiotic
Neomycin sulfate 3.5 mg	First aid antibiotic
Polymyxin B sulfate 10,000 units	First aid antibiotic
Pramoxine HCl 10 mg	External analgesic

**Uses**

first aid to help prevent infection and for the temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

**Warnings**

**For external use only.**

**Do not use**

- in the eyes
- over large areas of the body
- if you are allergic to any of the ingredients.

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns.

**Stop use and ask a doctor if**

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- a rash or other allergic reaction develops.

**Keep out of reach of children.** If swallowed get medical help or contact a Poison Control Center right away.

## **Directions**

Adults and children 2 years of age and older:

- clean the affected area and dry thoroughly
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage. Children under 2 years of age: ask a doctor

## **Other information**

- To open: unscrew cap, pull tab to remove foil seal
- store at 20° to 25°C (68° to 77°F)
- see carton or tube crimp for lot number and expiration date

## **Inactive ingredient**

white petrolatum

## **Questions?**

Call **1-866-923-4914**

## **PRINCIPAL DISPLAY PANEL - 28.4 g Tube Label**

**First Aid Antibiotic  
Pain Relieving Ointment**

**Maximum Strength  
Triple Antibiotic  
Ointment + Pain  
Relief**

*Bacitracin Zinc • Neomycin Sulfate  
Polymyxin B Sulfate • Pramoxine Hydrochloride*

**NET WT 1 oz (28.4 g)**

First Aid Antibiotic  
Pain Relieving Ointment

Maximum Strength

# Triple Antibiotic Ointment + Pain Relief



Bacitracin Zinc • Neomycin Sulfate  
Polymyxin B Sulfate • Pramoxine Hydrochloride

NET WT 1 oz (28.4 g)

### Active ingredients

(each gram contains)

	Purpose
Bacitracin zinc 500 units	First aid antibiotic
Neomycin sulfate 3.5 mg	First aid antibiotic
Polymyxin B sulfate 10,000 units	First aid antibiotic
Pramoxine HCl 10 mg	External analgesic

**Uses** first aid to help prevent infection and for the temporary relief of pain or discomfort in minor:  
• cuts • scrapes • burns

**Warnings** For external use only. Do not use • in the eyes • over large areas of the body • if you are allergic to any of the ingredients. **Ask a doctor before use if you have** • deep or puncture wounds • animal bites • serious burns. **Stop use and ask a doctor if** • you need to use longer than 1 week • condition persists or gets worse • symptoms persist for more

than 1 week, or clear up and occur again within a few days • a rash or other allergic reaction develops. **Keep out of reach of children.** If swallowed get medical help or contact a Poison Control Center right away.

**Directions** Adults and children 2 years of age and older: • clean the affected area and dry thoroughly • apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily • may be covered with a sterile bandage. Children under 2 years of age: ask a doctor

### Other information

• To open: unscrew cap, pull tab to remove foil seal  
• store at 20° to 25°C (68° to 77°F)  
• see carton or tube crimp for lot number and expiration date

**Inactive ingredient** white petrolatum

**Questions?** Call 1-866-923-4914

Distributed by:  
Made in Canada. LPK-0000-0 0510-0 00

## TRIPLE ANTIBIOTIC PLUS PAIN RELIEF

bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:51672-2027
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BACITRACIN ZINC</b> (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	500 [USP'U] in 1 g
<b>NEOMYCIN SULFATE</b> (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g
<b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
<b>PRAMOXINE HYDROCHLORIDE</b> (UNII: 88AYB867L5) (PRAMOXINE -	PRAMOXINE	10 mg in 1 g

UNII:068X84E056)

HYDROCHLORIDE

10 mg in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
WHITE PETROLATUM (UNII: B6E5W8RQJ4)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51672-2027-1	1 in 1 CARTON	03/31/2012	
1		14.2 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:51672-2027-2	1 in 1 CARTON	03/31/2012	
2		28.4 g in 1 TUBE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M004	03/31/2012	

**Labeler** - Taro Pharmaceuticals U.S.A., Inc. (145186370)**Establishment**

Name	Address	ID/FEI	Business Operations
Taro Pharmaceuticals Inc.		206263295	manufacture(51672-2027)

Revised: 1/2024

Taro Pharmaceuticals U.S.A., Inc.