

IBUPROFEN- ibuprofen tablet, film coated
Better Living Brands, LLC

Signature Care 44-393

Active ingredient (in each orange caplet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - toothache
 - the common cold
 - backache
 - headache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- right before or after heart surgery
- if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
 - chest pain
 - trouble breathing
 - leg swelling
 - slurred speech
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 caplet, 2 caplets may be used
 - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Signature

care®
Quality
Guaranteed

COMPARE TO

Motrin® IB Caplets
active ingredient†

NDC 21130-393-14

Ibuprofen

IBUPROFEN TABLETS USP, 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Actual Size

**500 COATED
CAPLETS**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**


†This product is not manufactured or distributed
by Kenvue Inc., distributors of Motrin® IB Caplets.
50844 ORG122139314

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0009
†1-888-723-3929

**LOVE IT
OR IT'S ON US†**

RD 23151

COMPARE TO
Motrin® IB Caplets
active ingredient†
NDC 21130-393-14




Quality
Guaranteed

Ibuprofen

IBUPROFEN TABLETS USP, 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Actual Size



**500 COATED
CAPLETS**

Drug Facts

Active ingredient
(in each orange caplet)
Ibuprofen USP,
200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose


Uses ■ temporarily relieves minor aches and pains
due to: ■ headache ■ the common cold
■ menstrual cramps ■ backache ■ toothache
■ muscular aches ■ minor pain of arthritis
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic
reaction, especially in people allergic to aspirin.
Symptoms may include:
■ facial swelling ■ blisters ■ hives ■ shock
■ skin reddening ■ asthma (wheezing) ■ rash
If an allergic reaction occurs, stop use and seek medical
help right away.
Stomach bleeding warning: This product contains an
NSAID, which may cause severe stomach bleeding. The
chance is higher if you
■ take more or for a longer time than directed
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or
nonprescription NSAIDs (aspirin, ibuprofen, naproxen,
or others) ■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ have 3 or more alcoholic drinks every day while using
this product
Heart attack and stroke warning: NSAIDs, except
aspirin, increase the risk of heart attack, heart failure,

Drug Facts (continued)
and stroke. These can be fatal. The risk is higher if you
use more than directed or for longer than directed.
Do not use ■ if you have ever had an allergic reaction
to any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as
heartburn ■ you are taking a diuretic
■ you have high blood pressure, heart disease, liver
cirrhosis, kidney disease, asthma, or had a stroke
■ you have problems or serious side effects from taking
pain relievers or fever reducers
Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because
ibuprofen may decrease this benefit of aspirin
■ taking any other drug
When using this product
■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if ■ you experience any of
the following signs of stomach bleeding: ■ feel faint
■ have bloody or black stools ■ vomit blood
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ slurred speech
■ leg swelling ■ trouble breathing
■ weakness in one part or side of body
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

Drug Facts (continued)
If pregnant or breast-feeding, ask a health
professional before use. It is especially important not
to use ibuprofen at 20 weeks or later in pregnancy
unless definitely directed to do so by a doctor because
it may cause problems in the unborn child or
complications during delivery.
Keep out of reach of children. In case of overdose,
get medical help or contact a Poison Control Center
right away.
Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 caplet
every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 caplet,
2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless
directed by a doctor
■ children under 12 years: ask a doctor
Other information
■ store between 20°-25°C (68°-77°F)
■ avoid excessive heat 40°C (104°F)
■ use by expiration date on package
Inactive ingredients carnauba wax, colloidal
silicon dioxide, corn starch, FD&C yellow #6 aluminum
lake, hypromellose, lactose anhydrous, magnesium
stearate, microcrystalline cellulose, polydextrose,
polyethylene glycol, sodium starch glycolate, stearic
acid, titanium dioxide
Questions or comments? Call 1-800-426-9391
8:30 AM-4:00 PM ET, Monday-Friday

RD 23151



50844 066122139314

LOVE IT
OR IT'S ON US†

THIS PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED
BY KENVUE INC. IT IS A DISTRIBUTOR OF MOTRIN® IB CAPLETS.
50844 066122139314

Signature Care 44-393

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-393
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	44;393
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-393-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	
2	NDC:21130-393-15	1 in 1 CARTON	03/01/1999	06/19/2023
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:21130-393-12	1 in 1 CARTON	03/01/1999	05/21/2017
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/01/1999	

Labeler - Better Living Brands, LLC (009137209)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(21130-393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(21130-393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(21130-393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(21130-393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(21130-393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(21130-393)

Revised: 1/2024

Better Living Brands, LLC