

**STOMACH RELIEF MAXIMUM STRENGTH- maximum strength pepto bismol suspension
McKesson (Sunmark)**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients

Bismuth Subsalicylate 525 mg

Purpose

upset stomach
reliever-antidiarrheal

Uses

relieves, upset stomach, heartburn
indigestion, diarrhea, nausea

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert- Contains salicylate.
Do not take if you are- taking other salicylate products,
allergic to salicylates (including aspirin)

Do not use if you have

an ulcer, a bleeding problem, bloody or black stool

Ask a doctor before use if you have

fever, mucus in stool

Ask a doctor or pharmacist before use if you are taking any drug for

anticoagulation (thinning blood)
diabetes, gout, arthritis

When using this product

a temporary, but harmless,

darkening of the stool and-or tongue may occur

Stop use and ask a doctor if

symptoms get worse
-ringing in the ears or loss of hearing occurs
-diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep this and all drugs out of the reach of children

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Directions

shake well before use
for accurate dosing, use dose cup
adults and children 12 years and over:
1 dose (2 tablespoons or 30 ml) every hour as needed

do not exceed 4 doses
(8 tablespoons or 120 ml) in 24 hours
use until diarrhea stops but not more than 2 days children under 12 years: ask a doctor
drink plenty of clear fluids to help prevent dehydration
caused by diarrhea

Other information

each tablespoon contains: sodium 6 mg
sugar free, low sodium, keep tightly closed
avoid excessive heat (over 104°F or 40°C)
protect from freezing.
Total salicylate per tablespoon, 236 mg
Visit www.peptic-drug-facts.com

Inactive ingredients

benzoic acid, flavor,
magnesium aluminum silicate, methyl cellulose, purified
water, red 22, red 28, saccharin sodium, salicylic acid,
sodium salicylate, sorbic acid

Principal Display Panel

COMPARE TO PEPTO BISMOL ACTIVE INGREDIENT

NDC 49348-923-37

STOMACH RELIEF LIQUID MAXIMUM STRENGTH
 SOOTHING RELIEF FOR UPSET STOMACH
 NAUSEA, INDIGESTION
 HEARTBURN AND DIARRHEA
 BISMUTH SUBSALICYLATE 525 mg
 PROTECTIVE COATING ACTION
 8 FL OZ (473 mL)

Drug Facts (continued)
Directions • shake well before use
 • use dosage cup provided • 1tbsp = tablespoon, mL = milliliter
 • adults and children 12 years and over:
 1 dose (2 tablespoons or 30 mL) every hour as needed
 • do not exceed 4 doses (8 tablespoons or 120 mL) in 24 hours
 • use until diarrhea stops but not more than 2 days
 • children under 12 years: ask a doctor
 • drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information
 • each 15 mL contains: sodium 15 mg
 • sugar free • low sodium • keep tightly closed
 • avoid excessive heat (over 104°F or 40°C)
 • protect from freezing • store at room temperature
 Total salicylate per tablespoon.....236 mg

Inactive ingredients benzoic acid, D&C red 22, D&C red 28, flavor, magnesium aluminum silicate, methyl cellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

Questions or comments? 1-866-534-4631

*This product is not manufactured or distributed by Procter & Gamble, the distributor of Pepto-Bismol®. TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL IS BROKEN OR MISSING.

McKesson
 Empowering Healthcare
 Another Quality Product Distributed by McKesson
 One Post Street, San Francisco, CA 94104
 Money Back Guarantee
 Please visit us at www.sunmarkbrand.com
 60-33744C

0 10939 33744 3

Drug Facts
Active ingredient (in each 15 mL, 1 tablespoon) Bismuth subsalicylate 525 mg.....upset stomach reliver/antidiarrheal
Purpose
Uses relieves • upset stomach • heartburn • indigestion • diarrhea • nausea
Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate.
Do not take if you are: • taking other salicylate products • allergic to salicylates (including aspirin)
Do not use if you have
 • an ulcer • a bleeding problem • bloody or black stool
Ask a doctor before use if you have
 • fever • mucus in stool
Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning blood) • diabetes • gout • arthritis
When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur
Stop use and ask a doctor if • symptoms get worse • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. ▶

STOMACH RELIEF MAXIMUM STRENGTH

maximum strength pepto bismol suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-923
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	

METHYLCELLULOSE (100 CPS) (UNII: 4GFU244C4J)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBIC ACID (UNII: X045WJ989B)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	MINT (mint)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-923-37	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/08/2013	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part335	11/08/2013	

Labeler - McKesson (Sunmark) (177667227)