

**25 PERSON ANSI - benzalkonium chloride, lidocaine, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, water, benzocaine, alcohol, ibuprofen, acetaminophen, aspirin, isopropyl alcohol**

**Genuine First Aid LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**25 Person ANSI**

**ACTIVE INGREDIENTS:**

Benzalkonium Chloride 0.13%

Lidocaine HCL 0.5%

Purpose: First aid antiseptic, external analgesic

Uses: First aid to help prevent infection and for the temporary relief of pain and itching associated with:

Minor Cuts

Scrapes

Burns

Warnings:

For external use only

Do not use: In eyes, in large quantities, over raw blistered areas, or on deep puncture wounds, animal bites or serious burns, for more than one week

Keep out of reach of Children.

If ingested, contact a Poison Control Center right away.

Directions: Clean affected area, Apply small amount not more than 3 times daily.

May be covered with a sterile bandage.

Other Information:

Store at room temperature

Genuine First Aid Burn Cream

Antiseptic Pain Relief With Aloe

Net Wt 0.9g (1/32 oz)

Manufactured in CHINA for

Genuine First Aid.

Reorder BC-001

# Genuine First Aid Burn Cream

**Antiseptic Pain Relief  
With Aloe**

**ACTIVE INGREDIENTS:**

**Benzalkonium**

**Chloride 0.13%**

**Lidocaine HCL 0.5%**

**Net Wt 0.9g - (1/32 oz)**



Manufactured in CHINA for  
**GENUINE FIRST AID**

**Drug Facts**

**Purpose:** First aid antiseptic,  
external analgesic

**Uses:** First aid to help prevent  
infection and for the temporary  
relief of pain and itching

associated with: • minor cuts  
• scrapes • burns

**Warnings:**

For external use only

**Do not use** • in eyes • in large  
quantities • over raw blistered  
areas, or on deep puncture  
wounds, animal bites or serious  
burns • for more than one week

**Keep out of reach of children.**

If ingested, contact a Poison  
Control Center right away.

**Directions:** • clean affected area  
• apply small amount not more  
than 3 times daily • may be  
covered with a sterile bandage

**Other information:**

Store at room temperature



600609



2015/01

Active Ingredient:

Purpose

Benzalkonium Chloride 0.40%..... First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive Ingredient: Purified water

LOT/EXP:

Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID



**DRUG FACTS - Antiseptic Towelette**

**Active Ingredient:** Benzalkonium Chloride 0.40%..First Aid Antiseptic

**Purpose:**

**Use:** For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

**Warnings:** For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If unusual redness, swelling or other symptoms occur, consult a physician immediately.

**Do not use:** In the eyes, or over large areas of the body.

**Directions:** Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

**Inactive ingredient:** Purified water.

TEAR HERE

Made in CHINA

Active Ingredient: .....Bacitracin Zinc 400 units

Neomycin Sulfate 5mg ( equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

Uses: To help prevent infection in:  
minor cuts; scrapes; burns

Warnings:

For external use only.

Do not use: in eyes; over large areas of the body;

If allergic to any of the ingredients; for more than one week unless directed by a physician.

Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg

Neomycin base)

Polymyxin B Sulfate 5000 units

Net Wt. 0.5g ; (1/64 oz)

Manufactured in CHINA for

GENUINE FIRST AID.

Reorder TAO-001

# Genuine Triple Antibiotic

**First Aid Ointment  
To Help Prevent Infection**

**Each Gram Contains:**  
**Bacitracin Zinc 400 units**  
**Neomycin Sulfate 5 mg**  
**(equivalent to 3.5 mg**  
**Neomycin base)**  
**Polymyxin B Sulfate 5000 units**  
**Net Wt 0.5g · (1/64 oz)**



Manufactured in CHINA for  
**GENUINE FIRST AID®**

## **Drug Facts**

**Uses:** To help prevent infection  
in • minor cuts • scrapes • burns

### **Warnings:**

**For external use only**

**Do not use** • in eyes • over large  
areas of the body • if allergic to  
any of the ingredients • for more  
than one week unless directed  
by a physician

**Stop use and consult a doctor**

• if the condition persists or gets  
worse • a rash or other allergic  
reaction develops

**Keep out of reach of children.**

If ingested, contact a Poison  
Control Center right away.

**Directions:** • clean affected area  
• apply small amount of this product  
(an amount equal to the surface  
area of the tip of a finger) on the  
area 1 to 3 times daily • may be  
covered with a sterile bandage

### **Other information:**

Store at room temperature

LOT

600609



2015/01

Active Ingredient:

Purified Water USP...q.s.

Inactive Ingredients:

Sodium Chloride USP 44mg

Monobasic Sodium Phosphate USP 18mg

Sodium Phosphate Dibasic USP 111mg

Edetate Disodium USP 10mg

Benzalkonium Chloride 0.5mg  
NF (as preservative)

Store in a cool place. For irrigation only.

Discard unused portion of the solution.

Not for injection.

Warning:

If you experience eye pain, changes in vision, continued redness or irritation of the eye,

or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor.

Do not use if solution changes color or becomes cloudy.

#### Directions

Remove contacts before using.

Twist top to remove.

Flush the affected area as needed. Control

Rate of flow by pressure on the bottle. Do not touch

tip of the container to any surface. Do not reuse.

If necessary continue flushing with emergency eyewash or shower.

Discard bottle after use.

#### Uses:

For flushing or irrigating the eyes to

remove loose foreign material, air pollutants,

or chlorinated water.

Code No.: GUJ/DRUG/G/1080

Batch No.:

Mfg Date:

Exp: Date:

10ml

Sterile Isotonic Buffered Genuine

Eyewash

For single use only

# 10ml

Sterile Isotonic buffered  
**GENUINE Eyewash**  
For single use only

**Directions:** Remove contact lens before using. Twist top to remove. Flush the affected area as needed. Control rate of flow by pressure on the bottle. Do not touch tip of the container to any surface. Do not reuse. If necessary, continue flushing with emergency eyewash or shower. Discard bottle after use.

**Uses:** For flushing or irrigating the eyes to remove loose foreign material, air pollutants, or chlorinated water.

**Each 10 ml contains:**

**Active Ingredient:**  
Purified Water ...q.s.

**Inactive Ingredients:**  
Sodium Chloride  
Monobasic Sodium Phosphate  
Dibasic Sodium Phosphate  
Edetate Disodium  
Benzalkonium Chloride  
NF (as preservative)

Store in a cool place. For irrigation only. Discard unused portion of the solution. Not for injection.

Mfg. Lic. No.: KTK/28/332/2002  
Batch No.:  
Mfg. date:  
Exp. date:

**Manufactured in INDIA for  
GENUINE FIRST AID**

**Warning:**  
If you experience eye pain, changes in vision, continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor. Do not use if solution changes color or becomes cloudy.

REORDER  
EYEW-10S

FOR EXTERNAL USE ONLY

**STERILE A**

Active Ingredient:

Purpose:

Benzocaine, 6% w/v..... Topical Anesthetic

SD alcohol, 60% w/v..... Antiseptic

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

Warnings: For external use only.

Avoid contact with eyes. If this happens, rinse thoroughly with water.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Flammable - keep away from fire or flame.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

Made in CHINA

LOT/EXP:

Insect Sting Relief Pad

Genuine First Aid LLC, Clearwater FL 33755

[www.GenuineFirstAid.com](http://www.GenuineFirstAid.com)

1/pouch

GENUINE FIRST AID






GENUINE FIRST AID®



ISO 9001:2010



Genuine First Aid LLC, Clearwater FL 33755  
www.GenuineFirstAid.com



REORDER ISRP-001

**DRUG FACTS - Insect Sting Relief Pad**

**Active Ingredient:** Benzocaine, 6% w/v ..... Topical Anesthetic

**Purpose:** Topical Anesthetic

SD alcohol, 60% w/v ..... Antiseptic

**Use:** For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

**Warnings:** For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Flammable - keep away from fire or flame. Avoid contact with eyes. If this happens, rinse thoroughly with water.

**Do not use:** In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

--- TEAR HERE ---

Made in CHINA

Active ingredient (in each tablet) Purpose

Ibuprofen USP (NSAID\*) 200mg ..... Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:

the common cold

headache

toothache

muscular aches

backache



minor pain of arthritis

menstrual cramps temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor before use if you are:

taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug

When using this product:

take with food or milk if stomach upset occurs

Stop use and ask a doctor If:

you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

**Adults and Children (12 years and older):** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information:** Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

Inactive ingredients: cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

## IBUPROFEN 2 Tablets

**Active ingredient (in each tablet)**      **Purpose**  
Ibuprofen USP (NSAID\*) 200mg . . . . . Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves minor aches and pains due to:  
■ the common cold ■ headache ■ toothache ■ muscular aches  
■ backache ■ minor pain of arthritis ■ menstrual cramps  
temporarily reduces fever

**Warnings Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ shock ■ facial swelling ■ asthma (wheezing) ■ rash ■ skin reddening ■ blisters ■ hives If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ taking any other drug containing an NSAID (prescription or nonprescription) ►

### Warnings (continued)

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

### When using this product

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor

**Adults and Children (12 years and older):** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information** ■ store at controlled room temperature ■ avoid excessive heat 40° C (104° F) ■ tamper evident sealed packets ■ do not use any opened or torn packets

**Inactive ingredients** cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

Active Ingredient (in each tablet)

Purpose

Acetaminophen 325 mg .....

Analgesic/antipyretic

Uses

temporary relief of minor aches and pains associated with:

common cold; headache; toothache; muscular aches; backache; arthritis; menstrual cramps; and reduction of fever

Warnings:

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if: adult takes more than 12 tablets in 24 hours, which is the maximum daily amount; child takes more than 5 doses in 24 hours; taken with other drugs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product

Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; for more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if: symptoms do not improve; pain gets worse or lasts for more than 10 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present; a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Do not exceed recommended dosage

#### Directions

Adults and Children 12 years of age or older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
--	--

Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
-------------------------------	--

Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.
----------------------------------	---

Store at 59-86 degree Farenheat (15-30 degree Celcius)  
tamper evident sealed packets; do not use any open or torn packets

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

GENUINE FIRST AID      2 Tablets

NON-ASPIRIN



GENUINE FIRST AID. 2 Tablets

# NON-ASPIRIN

**Active ingredient (in each tablet) Purpose**  
Acetaminophen 325 mg . . . . . Analgesic/antipyretic

### Uses

temporary relief of minor aches and pains associated with  
■ common cold ■ headache ■ toothache  
■ muscular aches ■ backache ■ arthritis  
■ menstrual cramps ■ and reduction of fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 12 tablets in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor

**Ask a doctor before use if the user has liver disease**

**Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin** ▶

### Warnings (continued)

**Stop use and ask a doctor if** ■ symptoms do not improve ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present ■ a rare sensitivity reaction occurs

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Do not exceed recommended dosage.**

### Directions

Adults and children 12 years of age and older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

**Other information** ■ store at 59°-86°F (15°-30°C) ■ tamper evident sealed packets ■ do not use any open or torn packets

**Inactive ingredients** corn starch, hydroxypropyl methylcellulose, polyethylene glycol, pregelatinized starch, stearic acid. May contain povidone and sodium starch glycolates.

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

Active Ingredient (in each tablet)

Purpose

Aspirin (NSAID\*) 325 mg..... Pain Reliever / fever reducer

\*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains associated with:

headache ; muscular aches ; minor arthritis pain ; backache ; common cold ; toothache ; menstrual cramps ; Temporarily reduces fever

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock, If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed

**Do not use:** if you have ever had an allergic reaction to any other pain reliever/ fever reducer; right before or after heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis

**Ask a doctor before use if:** stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

under a doctor's care for any serious condition; taking any other drug

When using this product: take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint; vomit blood; have bloody or black stools; have stomach

pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful areas; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or

contact a Poison Control Center right away.

Directions

do not use more than directed

the smallest effective dose should be used

drink a full glass of water with each dose

do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with

water every 4 hours as needed. Do not take more than 12 tablets in 24

hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); avoid

excessive heat and humidity; tamper evident sealed packets;

Do not use any opened or torn packets

Inactive Ingredients: hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

ASPIRIN 2 Tablets



# ASPIRIN

2 Tablets

**Active ingredient (in each tablet)**      **Purpose**  
Aspirin (NSAID)\* 325 mg . . . . . Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** Temporarily relieves minor aches and pains associated with  
■ headache ■ muscular aches ■ minor arthritis pain ■ backache  
■ common cold ■ toothache ■ menstrual cramps  
Temporarily reduces fever

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/ fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

## Warnings (continued)

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing  
■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear

If **pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not use more than directed  
■ the smallest effective dose should be used  
■ drink a full glass of water with each dose  
■ do not take longer than 10 days, unless directed by a doctor

**Adults and children: (12 years and older)** Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information** ■ store at 59°-86°F (15°-30°C) ■ avoid excessive heat and humidity ■ tamper evident sealed packets ■ do not use any opened or torn packets

**Inactive ingredients** hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by **GENUINE FIRST AID**  
600 Cleveland Str Suite 400, Clearwater, FL 33755

Active Ingredient: Purpose

Isopropyl Alcohol, 70% v/v..... Antiseptic

Use: For preparation of skin before injection.

Warnings: For external use only.

Flammable - keep away from fire or flame

Store at room temperature 15-30 degree Celsius (59-86 degree Fahrenheit)

Do not use: with electrocautery, in the eyes.

Stop use if irritation and redness develop. If condition persists for more than 72 hours, consult your doctor.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Wipe Injection site vigorously and discard.

Inactive Ingredient: Purified water.

LOT/EXP: Made in CHINA

20140301

Alcohol Cleansing Pad  
Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID



REORDER ACP-001

**Alcohol Cleansing Pad**  
**Toallitas Húmedas con Alcohol**



Genuine First Aid LLC, Clearwater FL 33755  
www.GenuineFirstAid.com



**GENUINE FIRST AID®**



**DRUG FACTS - Alcohol Cleansing Pad**

**Active Ingredient:** Isopropyl Alcohol, 70% v/v . . . . . **Purpose:** Antiseptic

**Use:** For preparation of the skin before injection.

**Warnings: For external use only.**

**Flammable - keep away from fire or flame.**

**Do not use:** with electrocautery, in the eyes

**Stop use** if irritation and redness develop.

If condition persists for more than 72 hours, consult your doctor.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions:** Wipe injection site vigorously and discard.

**Other information:** Store at room temperature 15°-30° C (59°-86° F)

**Inactive ingredient:** Purified water.

TEAR HERE

LOT/EXP:                      Made in CHINA  
**20140301**

Active Ingredient:

Ethyl Alcohol 62 percent

Enter section text here

Warnings:

For external use only

Flammable, keep away from fire or flame

Do not use in the eyes. In case of contact, rinse eyes thoroughly with water.

Stop use and consult a doctor if irritation and redness develop

and persist for more than 72 hours.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

Empty contents into palm. Rub hands until gel dissipates. Recommended for repeated use.

Other Information: Store at 15 to 25 degree Celsius (59 to 77 degree Fahrenheit)

Genuine Hand Sanitizer

Instant Gel

Antiseptic Handwash

with Vitamin E and Aloe

Kills 99.9 percent of germs

Without Water

Active Ingredient:

Ethyl Alcohol 62 percent

Net Wt 0.9g (1/32 oz)

Manufactured in China For

Genuine First Aid.

Reorder HS-001

# Genuine Hand Sanitizer

**Instant Gel  
Antiseptic Handwash  
with Vitamin E & Aloe**

**Kills 99.9% of Germs  
Without Water  
Active Ingredient:  
Ethyl alcohol 62%  
Net Wt 0.9g - (1/32 oz)**



Manufactured in CHINA for  
**GENUINE FIRST AID®**

## **Drug Facts**

**Uses:** For hand washing to decrease bacteria on the skin

### **Warnings:**

**For external use only**

Flammable, keep away from fire or flame

**Do not use in the eyes.** In case of contact, rinse eyes thoroughly with water

**Stop use and consult a doctor** if irritation and redness develop and persist for more than 72 hours

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions:** Empty contents into palm. Rub hands until gel dissipates. Recommended for repeated use

**Other information:** Store at 15° to 25° C (59° to 77° F)



600600



2015/01

ANSI/ISEA Z308.1-2009 TYPE III

Caution! This Kit meets ANSI/ISEA Z308.1-2009 only when required minimum fill is maintained with

first aid products marked "ANSI/ISEA Z308.1-2009."

#### CARRYING CASE

1 Easy Access Pocket System  
1 Hard Case

#### POCKET 1: SMALL CUTS AND BURNS

50 Adhesive Plastic Bandages 1"x3"  
1 Knuckle Fabric Bandage  
25 Adhesive Spot Bandages 7/8"x7/8"  
1 Fingertrip Fabric Bandage  
1 Elbow Knee Adhesive Bandage  
6 Burn Cream  
10 Antiseptic Towelettes  
6 Triple Antibiotic Ointment 0.5gr  
20 Cotton Tipped Applicators

#### POCKET 2: MEDIUM CUTS AND SCRATCHES

10 Antiseptic Towelettes  
4 Sterile Gauze Pad 3"x3"  
2 Sterile Gauze Pads 4"x4"  
2 Roller Gauze Bandage 2"x4.1yds  
1 Roller Gauze Bandage 3"x4.1yds  
1 Sterile Eye Pads  
1 Sterile Eye Wash 10ml Twist Top  
1 Butterfly Wound Closure, Medium

#### POCKET 3: SEVERE BLEEDING AND BURNS

1 First Aid Tape Roll 1/2"x5 yds.  
1 Combine Pad 5"x9"

#### POCKET 4: CPR

1 CPR Breathing Barrier

#### POCKET 5: PROTECTION

4 Medical Grade Vinyl Gloves  
1 Instant Cold Compress  
20 Alcohol Cleansing Pads  
1 Hand Sanitizer 0.9g  
1 Insect Sting Relief Pads  
1 Triangular Bandage 42"x42"x59"  
2 Ibuprofen 200mg  
2 Non Aspirin 325mg  
2 Aspirin 325mg

#### POCKET 6: INSTRUMENTS

1 Emergency First Aid Guide  
1 Plastic Tweezers  
1 Scissors

Manufactured in China for:

Genuine First Aid LLC.  
600 Cleveland Street  
Suite 400  
Clearwater FL 33755

www.GenuineFirstAid.com

GENUINE FIRST AID

Copyright c 2010 Genuine First Aid LLC. All rights reserved.

**ANSI/ISEA Z308.1-2009 TYPE III**  
Caution! This kit meets ANSI/ISEA Z308.1-2009 only when required minimum fill is maintained with first aid products marked 'ANSI/ISEA Z308.1-2009.'

**CARRYING CASE**

- 1 Easy Access Pocket System
- 1 Hard case

**POCKET 1: SMALL CUTS & BURNS**

- 50 Adhesive Plastic Bandages 1" x 3"
- 1 Knuckle Fabric Bandage
- 25 Adhesive Spot Bandages 7/8" x 7/8"
- 1 Fingertip Fabric Bandage
- 1 Elbow Knee Adhesive Bandage
- 6 Burn Cream
- 10 Antiseptic Towelettes
- 6 Triple Antibiotic Ointment 0.5g
- 20 Cotton Tipped Applicators

**POCKET 2: MEDIUM CUTS & SCRATCHES**

- 10 Antiseptic Towelettes
- 4 Sterile Gauze Pads 3" x 3"
- 2 Sterile Gauze Pads 4" x 4"
- 2 Roller Gauze Bandages 2" x 4,1yds
- 1 Roller Gauze Bandage 3" x 4,1yds
- 1 Sterile Eye Pad
- 1 Sterile Eye Wash 10ml Twist Top
- 1 Butterfly Wound Closure, Medium

**POCKET 3: SEVERE BLEEDING & BURNS**

- 1 First Aid Tape Roll 1/2" x 5yds
- 1 Combine Pad 5" x 9"

**POCKET 4: CPR**

- 1 CPR Breathing Barrier

**POCKET 5: PROTECTION**

- 4 Medical Grade Vinyl Gloves
- 1 Instant Cold Compress
- 20 Alcohol Cleansing Pads
- 1 Hand Sanitizer 0.9g
- 1 Insect Sting Relief Pad
- 1 Triangular Bandage 42" x 42" x 59"
- 2 Ibuprofen 200mg
- 2 Non Aspirin 325mg
- 2 Aspirin 325mg

**POCKET 6: INSTRUMENTS**

- 1 Emergency First Aid Guide
- 1 Plastic Tweezers
- 1 Scissors

**ESTUCHE PORTATIL**

- 1 Sistema de fundas para rapido acceso
- 1 Estuche de Plastico

**FUNDA 1: CORTADURAS Y QUEMADURAS MENORES**

- 50 Vendas Adhesivas Plasticas 1" X 3"
- 1 Vendas de Tela para Nudillos
- 25 Vendas Adhesivas Redondas 7/8" X 7/8"
- 1 Vendas de tela para yema del dedo
- 1 Vendas adhesivas para codo / rodilla
- 6 Crema para Quemaduras
- 10 Toallitas Antisepticas
- 6 Pomada de Antibiotico Triple 0.5g
- 20 Aplicadores con Puntas de Algodon

**FUNDA 2: CORTADURAS Y RASGUÑOS MEDIANOS**

- 10 Toallitas Antisepticas
- 4 Almohadilla Esteril de Gasa 3" X 3"
- 2 Almohadilla Esteril de Gasa 4" x 4"
- 2 Vendas de Gasa en Rollo 2" X 4,1 yds
- 1 Vendas de Gasa en Rollo 3" X 4,1 yds
- 1 Almohadilla Esteril para Ojos
- 1 Solucion Esteril Ocular 10ml
- 1 Vendas Adhesivas Tipo Mariposa

**FUNDA 3: SANGRADO SEVERO Y QUEMADURAS**

- 1 Cinta de Papel para Primeros Auxilios 1/2" x 5yds
- 1 Almohadilla de Uso Combinado 5" x 9"

**FUNDA 4: CPR**

- 1 Mascara CPR

**FUNDA 5: PROTECCION**


- 4 Guantes de Vinilo
- 1 Compresa de Frio Instantaneo
- 20 Toallitas Humedas con Alcohol
- 1 Desinfectante de Manos 0.9g
- 1 Toallitas para Picaduras de Insectos
- 1 Vendajes Triangulares 42" x 42" x 59"
- 2 Ibuprofen 200mg
- 2 Sin Aspirina 325mg
- 2 Aspirina 325mg

**FUNDA 6: INSTRUMENTOS**

- 1 Manual de Primeros Auxilios
- 1 Pinzas Plasticas
- 1 Tijeras

Manufactured in China for:  
Genuine First Aid LLC.  
600 Cleveland Street  
Suite 400  
Clearwater, FL 33755  
www.genuinefirstaid.com

**GENUINE FIRST AID.**



Copyright © 2010 Genuine First Aid LLC.  
All rights reserved.

## 25 PERSON ANSI

benzalkonium chloride, lidocaine, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, water, benzocaine, alcohol, ibuprofen, acetaminophen, aspirin, isopropyl alcohol kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:52124-0112
---------------------	----------------	---------------------------	----------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0112-1	1 in 1 KIT		

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	6 PACKAGE	5.4 g

<b>Part 2</b>	20 PACKAGE	16 mL
<b>Part 3</b>	6 TUBE	3 g
<b>Part 4</b>	1 BOTTLE	10 mL
<b>Part 5</b>	1 PACKAGE	0.5 mL
<b>Part 6</b>	1 PACKAGE	2
<b>Part 7</b>	1 PACKAGE	2
<b>Part 8</b>	1 PACKAGE	2
<b>Part 9</b>	20 PACKAGE	10 mL
<b>Part 10</b>	1 PACKAGE	0.9 g

## Part 1 of 10

### GENUINE FIRST AID BURN ANTISEPTIC PAIN RELIEF WITH ALOE

benzalkonium chloride, lidocaine cream

#### Product Information

**Item Code (Source)** NDC:52124-0004

**Route of Administration** TOPICAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.5 g in 100 g

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0004-1	0.9 g in 1 PACKAGE		

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part345	04/25/2010	

## Part 2 of 10

### ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

#### Product Information

<b>Item Code (Source)</b>	NDC:52124-0001
<b>Route of Administration</b>	TOPICAL

<b>Active Ingredient/Active Moiety</b>		
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.40 mL in 100 mL

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
<b>WATER</b> (UNII: 059QF0K00R)	

<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		

<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
OTC monograph final	part333	04/25/2010		

**Part 3 of 10**

**GENUINE TRIPLE ANTIBIOTIC**  
bacitracin zinc,neomycin sulfate,polymyxin b sulfat e ointment

<b>Product Information</b>	
<b>Item Code (Source)</b>	NDC:52124-0003
<b>Route of Administration</b>	TOPICAL

<b>Active Ingredient/Active Moiety</b>		
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BACITRACIN ZINC</b> (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	400 [iU] in 1 g
<b>NEOMYCIN SULFATE</b> (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	5 mg in 1 g
<b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B SULFATE	5000 [iU] in 1 g

<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>



1	NDC:52124-0003-1	0.5 g in 1 TUBE		
---	------------------	-----------------	--	--

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	04/25/2010	

### Part 4 of 10

#### STERILE ISOTONIC BUFFERED GENUINE EYEWASH

water liquid

### Product Information

Item Code (Source)	NDC:52124-0005
Route of Administration	OPHTHALMIC

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0K00R) (WATER - UNII:059QF0K00R)	WATER	98.6 mL in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, MONOBASIC (UNII: 3980JIH2SW)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0005-1	10 mL in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part349	04/25/2010	

### Part 5 of 10

#### INSECT STING RELIEF PAD

benzocaine, alcohol liquid

### Product Information

**Item Code (Source)** NDC:52124-0008

**Route of Administration** TOPICAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	6 mL in 100 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	60 mL in 100 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0008-1	0.5 mL in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part348	04/25/2010	

### Part 6 of 10

### IBUPROFEN

ibuprofen tablet

### Product Information

**Item Code (Source)** NDC:52124-0009

**Route of Administration** ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE (UNII: 3NXW29V3WO)	

LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

### Product Characteristics

Color	white (White)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;352
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0009-1	2 in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	04/25/2010	

### Part 7 of 10

#### NON-ASPIRIN

acetaminophen tablet

### Product Information

Item Code (Source)	NDC:52124-0010
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)

STEARIC ACID (UNII: 4ELV7Z65AP)

POVIDONE (UNII: FZ989GH94E)

### Product Characteristics

Color	white (WHITE)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	AZ;234
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0010-1	2 in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	04/25/2010	

## Part 8 of 10

### ASPIRIN

aspirin tablet

### Product Information

Item Code (Source)	NDC:52124-0011
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	white (White)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0011-1	2 in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	04/24/2010	

### Part 9 of 10

### ALCOHOL CLEANSING PAD

isopropyl alcohol liquid

### Product Information

<b>Item Code (Source)</b>	NDC:52124-0002
<b>Route of Administration</b>	TOPICAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	70 mL in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0002-1	0.5 mL in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	04/25/2010	

## Part 10 of 10

### GENUINE HAND SANITIZER INSTANT ANTISEPTIC HANDWASH WITH VITAMIN E AND ALOE

alcohol gel

#### Product Information

Item Code (Source) NDC:52124-0006

Route of Administration TOPICAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	62 g in 100 g

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0006-1	0.9 g in 1 PACKAGE		

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	04/25/2010	

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	04/25/2010	

**Labeler** - Genuine First Aid LLC (619609857)

#### Establishment

Name	Address	ID/FEI	Business Operations
GFA Production ( Xiamen) Co., Ltd		421256261	manufacture