

THYROSAFE- potassium iodide tablet
BTG International Inc.

Take potassium iodide (KI) only when public officials tell you. In a nuclear radiation emergency, radioactive iodine could be released into the air. KI protects only the thyroid gland from uptake of radioactive iodine. Therefore, KI should be used along with other emergency measures that will be recommended to you by public officials. If you are told to take this medicine, take it 1 time every 24 hours. Do not take it more often. More KI will not help you. Too much KI may increase the chances of side effects. Do not take this medicine if you know you are allergic to iodine (see SIDE EFFECTS below).

DESCRIPTION

Each white, round, cross-scored ThyroSafe® tablet contains 65 mg of potassium iodide.

INDICATIONS

ThyroSafe® (Potassium Iodide Tablets, USP) is a thyroid blocking medicine that is used in a nuclear radiation emergency only.

DIRECTION FOR USE

Use only as directed by public officials if a nuclear radiation emergency happens.

Dose:

Adults over 18 years	2 tablets (whole or crushed) every day (130 mg)
Children over 12 years to 18 years who weigh at least 150 pounds	2 tablets (whole or crushed) every day (130 mg)
Children over 12 years to 18 years who weigh less than 150 pounds	1 tablet (whole or crushed) or 8 teaspoons every day (65 mg)
Children over 3 years to 12 years	1 tablet (whole or crushed) or 8 teaspoons every day (65 mg)
Children over 1 month to 3 years	4 teaspoons every day (32.5 mg)
Babies at birth to 1 month	2 teaspoons every day (16.25 mg)

Tablets can be crushed and mixed in many liquids. To take the tablet in liquid solution, use dosing directions under **Making a Potassium Iodide Liquid Mixture**.

Take KI every day (every 24 hours) as directed by public officials. Do not take more than 1 dose in 24 hours. More will not help you. Too much medicine may increase the chances of side effects.

Making a Potassium Iodide Liquid Mixture:

1. Put **one** 65 mg KI tablet into a small bowl and grind it into a fine powder using the back of a metal teaspoon against the inside of the bowl. The powder should not have any large pieces.
2. Add 4 teaspoons of water to the crushed KI powder in the bowl and mix until the KI powder is dissolved in the water.
3. Take the KI water mixture solution made in step 2 and mix it with 4 teaspoons of low fat white or chocolate milk, orange juice, flat soda, raspberry syrup, or infant formula.
4. The KI liquid mixture will keep for up to 7 days in the refrigerator. It is recommended that the KI liquid mixtures be prepared weekly. Throw away unused portions.

The amount of KI (65 mg tablet) in the drink when mixed as described above is 8.125 mg per teaspoon. The number of teaspoons of the drink to give your child depends on your child's age as described in the following table:

Child's Age	Give your child this amount in teaspoons
Over 12 to 18 years old who weigh less than 150 pounds	8 teaspoons will give you a 65 mg dose
Over 3 to 12 years old	8 teaspoons will give you a 65 mg dose
Over 1 month to 3 years old	4 teaspoons will give you a 32.5 mg dose
Birth to 1 month	2 teaspoons will give you a 16.25 mg dose

Note: This is the amount to give your child for **one** single dose in **teaspoons (not** tablespoons). You should give your child **one** dose each day as recommended by the public officials.

Pregnant or breastfeeding women or babies under 1 month of age: Take as directed above and call a doctor as soon as possible. Repeat dosing should be avoided. It is recommended that thyroid function be checked in babies less than 1 month of age that take KI. Women who are pregnant or breastfeeding should also be checked by a doctor if repeat dosing is necessary. Although these precautions should be taken, the benefits of short-term use of KI to block uptake of radioactive iodine by the thyroid gland far exceed its chances of side effects.

Patients with thyroid disease: If you have both a nodular thyroid condition such as multinodular goiter with heart disease, you should not take KI. Patients with other thyroid conditions may take KI as directed above, but call a doctor if you need to take KI for more than a few days.

WARNING

People who are allergic to iodine, have dermatitis herpetiformis or hypocomplementemic vasculitis, or have nodular thyroid disease with heart disease should not take KI. Keep out of the reach of children. In case of an allergic reaction (difficulty breathing, speaking or swallowing; wheezing; shortness of breath or swelling of the mouth or throat), call 911 or get medical care right away. In case of overdose, get medical help or call a Poison Control Center right away.

HOW POTASSIUM IODIDE WORKS

Certain forms of iodine help your thyroid gland work right. Most people get the iodine they need from foods like iodized salt or fish. The thyroid can "store" or hold only a certain amount of iodine.

In a nuclear radiation emergency, radioactive iodine may be released in the air. This material may be breathed or swallowed. It may enter the thyroid gland and damage it. The damage would probably not show itself for years. Children are most likely to have thyroid damage.

If you take KI, it will block or reduce the chances that radioactive iodine will enter your thyroid gland.

WHO SHOULD NOT TAKE POTASSIUM IODIDE

People should avoid KI if they are allergic to iodine, have dermatitis herpetiformis or hypocomplementemic vasculitis, or have nodular thyroid disease with heart disease, because these conditions may increase the chances of side effects to iodine.

HOW AND WHEN TO TAKE POTASSIUM IODIDE

KI should be taken as soon as possible after public officials tell you. If you are told to repeat the dose, you should take the second dose 24 hours after the first dose. Do not take it sooner. More KI will not help you because the thyroid can "hold" only certain amounts of iodine. Taking more than 1 dose per day will increase the chances of side effects. The public officials will tell you how many days to take KI. You should take KI until the chances of major exposure to radioactive iodine by breathing or swallowing stops.

SIDE EFFECTS

Short-term use of KI at the recommended dose is safe. You should not take this drug for longer than you are told.

Possible side effects include: swelling of the salivary glands, nausea, vomiting, diarrhea, stomach ache, fever, headache, metallic taste, and allergic reactions. Allergic reactions can include

- skin rashes such as hives
- swelling of various parts of the body such as the face, lips, tongue, throat, hands or feet
- fever with joint pain
- trouble breathing, speaking or swallowing
- wheezing or shortness of breath

Get medical attention right away if you have trouble breathing, speaking or swallowing; wheezing; shortness of breath; or swelling of the mouth, tongue or throat.

Taking iodide, in rare cases, may cause overactivity of the thyroid gland, underactivity of the thyroid gland, or enlargement of the thyroid gland (goiter). Symptoms of an overactive thyroid gland may include an irregular heart beat and chest pain. Patients with thyroid disease are more likely to get these side effects. Babies under 1 month of age are more likely to get an underactive thyroid gland (hypothyroidism).

WHAT TO DO IF SIDE EFFECTS OCCUR

Stop taking KI and call a doctor if you have one or more of the following symptoms:

- swelling of the face, hands or feet

- fever and joint pain
- skin rash

Stop taking KI and get medical help right away if you have one or more of the following symptoms:

- trouble breathing, speaking or swallowing
- shortness of breath or wheezing
- swelling of the lips, tongue or throat
- irregular heart beat or chest pain

HOW SUPPLIED

ThyroSafe® (potassium iodide, USP) tablets. Packages of 10 and 20 tablets. Each white, round, cross-scored tablet contains 65 mg potassium iodide. Store at 20-25° C (68-77° F). Keep dry and foil intact.

Manufactured by Astrea Fontaine SAS,
21121 Fontaine les Dijon, France
for BTG International Inc., West Conshohocken, PA 19428 USA
1-866-849-7672.
www.thyrosafe.com.

Revised 02/2022

Drug Facts

Other information

- store at 20-25° C (68-77° F)
- protect from light
- keep dry and foil intact
- **do not throw away consumer package insert**

Inactive ingredients

lactose, magnesium stearate, microcrystalline cellulose

Questions or comments?

call toll free 1-866-849-7672

Carton Label - Principal Display Panel

NDC-50633-910-20

ThyroSafe®

Potassium Iodide Tablets, USP, 65 mg

Thyroid blocking in a radiation emergency only

20 tablets, 65 milligrams each

Carton Label - Back panel

Potassium Iodide (KI) Tablets USP, 65 mg

Drug Facts	
Active Ingredient (in each tablet)	Purpose
Potassium Iodide 65 mg	Thyroid blocking
Use helps prevent radioactive iodine from getting into the thyroid gland during a nuclear	

radiation emergency. Use along with other emergency measures recommended by public officials.

Warnings

Allergy alert: Iodine may cause an allergic reaction with 1 or more of the following symptoms:

- shortness of breath or wheezing
- swelling
- skin rash
- trouble breathing, speaking or swallowing
- fever and joint pain

Do not use if you have

- ever had an allergic reaction to iodine
- nodular thyroid disease with heart disease
- hypocomplementemic vasculitis
- dermatitis herpetiformis

Stop use and ask a doctor if you have if you have

- allergic reaction. Get medical help right away if you have trouble breathing, speaking or swallowing; shortness of breath; wheezing; swelling of the mouth, tongue or throat; or rash.
- irregular heart or chest pain. Get medical help right away.
- swelling of the hands or feet, fever, or joint pain.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- use as directed by public officials in the event of a nuclear radiation emergency.
- do not take more than 1 dose in 24 hours.
- tablets can be whole or crushed and mixed in milk, baby formula, water, orange juice, flat soda like cola, or raspberry syrup. **The liquid mixture should be given to infants, young children, and others who cannot swallow tablets; see consumer package insert on how to make a liquid mixture.**

adults over 18 years	2 tablets (whole or crushed) daily (130 mg)
children over 12 years to 18 years who weigh at least 150 pounds	2 tablets (whole or crushed) daily (130 mg)
children over 12 years to 18 years who weigh less than 150 pounds	1 tablet (whole or crushed) daily (65 mg)
children over 3 years to 12 years	1 tablet (whole or crushed) daily (65 mg)
children over 1 month to 3 years	1/2 tablet (crushed) daily (32.5 mg)
babies at birth to 1 month	16.25 mg daily as directed in consumer package insert

If pregnant, breastfeeding, have a baby up to 1 month of age, or have thyroid disease (except nodular thyroid disease with heart disease), take as directed above and contact a doctor as soon as possible

Principal Display Panel - 65 mg Carton Label

NDC-50633-910-20

ThyroSafe®

Potassium Iodide Tablets, USP, 65 mg

Thyroid blocking in a radiation emergency only

20 tablets, 65 milligrams each

BTG International Inc.

ThyroSafe[®]
Potassium Iodide Tablets,
USP, 65 mg

Potassium Iodide (KI) Tablets USP, 65 mg

Drug Facts

Active Ingredient (in each tablet)	Purpose
Potassium Iodide 65 mg	Thyroid blocking

Use helps prevent radioactive iodine from getting into the thyroid gland during a nuclear radiation emergency. Use along with other emergency measures recommended by public officials.

Warnings

Allergy alert: Iodine may cause an allergic reaction with 1 or more of the following symptoms:
 - shortness of breath or wheezing
 - swelling
 - skin rash
 - trouble breathing, speaking or swallowing
 - fever and joint pain

Do not use if you have

- ever had an allergic reaction to iodine
- nodular thyroid disease with heart disease
- hypocomplementemic vasculitis
- dermatitis herpetiformis

Stop use and ask a doctor if you have

- allergic reaction. Get medical help right away if you have trouble breathing, speaking or swallowing; shortness of breath; wheezing; swelling of the mouth, tongue or throat; or rash, irregular heart or chest pain. Get medical help right away.
- swelling of the hands or feet, fever, or joint pain

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- use as directed by public officials in the event of a nuclear radiation emergency
- do not take more than 1 dose in 24 hours
- tablets can be whole or crushed and mixed in milk, baby formula, water, orange juice, flat soda like cola, or raspberry syrup. **The liquid mixture should be given to infants, young children, and others who cannot swallow tablets; see consumer package insert on how to make a liquid mixture.**

adults over 18 years	2 tablets (whole or crushed) daily (130 mg)
children over 12 years to 18 years who weigh at least 150 pounds	2 tablets (whole or crushed) daily (130 mg)
children over 12 years to 18 years who weigh less than 150 pounds	1 tablet (whole or crushed) daily (65 mg)
children over 3 years to 12 years	1 tablet (whole or crushed) daily (65 mg)
children over 1 month to 3 years	½ tablet (crushed) daily (32.5 mg)
babies at birth to 1 month	16.25 mg daily as directed in consumer package insert

- if pregnant, breastfeeding, have a baby up to 1 month of age, or have thyroid disease (except nodular thyroid disease with heart disease), take as directed above and contact a doctor as soon as possible

1TH20 US
20221 ET

Ingredients
lactose, magnesium stearate, microcrystalline cellulose

Questions or comments?
call toll free 1-866-849-7672

Drug Facts (Continued)

Other information

- store at 20-25°C (68-77°F)
- protect from light
- keep dry and foil intact
- do not throw away consumer package insert

608152B

Thyroid blocking in a radiation emergency only
20 tablets, 65 milligrams each

ThyroSafe[®]
Potassium Iodide Tablets, USP, 65 mg

NDC-50633-910-20

ThyroSafe[®]
Potassium Iodide Tablets,
USP, 65 mg


Manufactured by Astra Fontaine SAS,
21121 Fontaine les Dijon France
for B1G International Inc.,
West Conshohocken, PA 19428 USA
www.thyrosafe.com



Batch No: Exp:



Principal Display Panel - Pack Label



Recipharm Service Packaging Artwork designer : Denis Margueritte		Client / customer Recipharm	
Désignation / Product name ALLU.THYROSAFE.65 MG.10 COMPL.200.US		Technique / technical Dimensions / cut size 110 x 43,7 - L. 200 Réf. découpe / cut ref.	
Couleurs / colors <input checked="" type="radio"/> Black <input checked="" type="radio"/> Technical color		Validation Date Name (regulatory person) Visa Pamela Ventura	
<input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/>		Code article : / Item code: 608151A Maquette n°: Draft n°: 2 Date : 24/07/20 Maq. <input checked="" type="radio"/> <input type="radio"/> Exé. <input type="radio"/>	
		Font type Helvetica Neue Font size (mini) 6,5 pt	

THYROSAFE

potassium iodide tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50633-910
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Potassium Iodide (UNII: 1C4QK22F9J) (Iodide Ion - UNII:09G416V86Q)	Potassium Iodide	65 mg

Inactive Ingredients

Ingredient Name	Strength
Lactose, Unspecified form (UNII: J2B2A4N98G)	
Microcrystalline Cellulose (UNII: OP1R32D61U)	
Magnesium Stearate (UNII: 70097M6I30)	

Product Characteristics

Color	white (white)	Score	4 pieces
Shape	ROUND (ROUND)	Size	9mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50633-910-20	160 in 1 CARTON	09/30/2002	
1		20 in 1 BOX		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076350	09/30/2002	

Labeler - BTG International Inc. (617382395)

Establishment

Name	Address	ID/FEI	Business Operations
Astrea Fontaine		260428962	MANUFACTURE(50633-910)

Revised: 3/2022

BTG International Inc.