

**NITETIME COLD AND FLU- acetaminophen, dextromethorphan hbr,  
doxylamine succinate, phenylephrine hcl solution  
Meijer Distribution Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Meijer Distribution, Inc. NiteTime Cold & Flu Drug Facts**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### **Ask doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

## Questions?

**1-800-719-9260**

### **Package/Label Principal Display Panel**

meijer®

Compare to Vicks® NyQuil® Severe active ingredients

MAXIMUM STRENGTH

nitetime cold & flu

Acetaminophen

Pain Reliever | Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Doxylamine Succinate

Antihistamine

Phenylephrine HCl

Nasal Decongestant



SEVERE

Multi-Symptom Relief

Mixed Berry Flavor

Sugar Free | Alcohol Free

12 FL OZ (355 mL)



## NITETIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-763
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:735X3R0TS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3COOX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KOOR)	

<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	RED (clear, dark)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-763-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/11/2013	
2	NDC:41250-763-50	2 in 1 PACKAGE	12/11/2013	12/11/2013
2		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/11/2013	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 8/2022

Meijer Distribution Inc