

ANTI-DIARRHEAL- loperamide hcl tablet
Kroger Company

Kroger 44-375

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

1-800-632-6900

Principal Display Panel

COMPARE TO the active ingredient of IMODIUM® A-D *See side panel

NDC 30142-775-08

OUR PHARMACIST RECOMMENDED

Kroger®

Anti-Diarrheal

Loperamide Hydrochloride

Tablets, 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea

actual
size

24 caplets

**capsule-shaped tablets

1. To open, tear at
perforations and
remove a section.

2. Use scissors to
cut through
plastic for caplet.

**TAMPER EVIDENT:
DO NOT USE IF PACKAGE
IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS
OF TAMPERING**

*Imodium® is a registered trademark of Johnson & Johnson Corporation, New Brunswick, NJ 08933. Johnson & Johnson Corporation is not affiliated with The Kroger Co. or this product.
50844 ORG061937508

DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202
QUALITY GUARANTEE
800-632-6900
www.kroger.com





Anti-Diarrheal

Loperamide Hydrochloride Tablets, 2 mg Anti-Diarrheal

Controls the Symptoms of Diarrhea



24 CAPLETS**
**capsule-shaped tablets

2. Use scissors to cut through plastic for caplet.



Registered trademark of Johnson & Johnson
Newark, NJ 08933. Johnson & Johnson
associated with The Kroger Co. or this product.
1937508



PLEASE RECYCLE

80% SIZE
12-digit UPC-A
(Not-suppressed)
For Position Only
0 41260 00995 2

DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202
QUALITY GUARANTEE
800-632-6900
www.kroger.com

**TAMPER EVIDENT:
DO NOT USE IF PACKAGE
IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS
OF TAMPERING**

B-1050-375-08
086061937508

Drug Facts (continued)

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death.
Do not use if you have bloody or black stool.

Ask a doctor before use if you have a fever, mucus in the stool, a history of liver disease, or a history of abnormal heart rhythm.

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if symptoms get worse or diarrhea lasts for more than 2 days. If you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. ■ Find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours
children 6-8 years (48-69 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number

Inactive ingredients D&C blue #1 aluminum lake, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments? 1-800-632-6900

ANTI-DIARRHEAL

loperamide hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-775
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-775-02	2 in 1 CARTON	05/06/2020	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:30142-775-08	4 in 1 CARTON	05/06/2020	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/06/2020	

Labeler - Kroger Company (006999528)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(30142-775) , pack(30142-775)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(30142-775)

Revised: 3/2024

Kroger Company