

ABIRATERONE ACETATE - abiraterone acetate tablet
Major Pharmaceuticals

BRIEF SUMMARY OF PRESCRIBING INFORMATION
These highlights do not include all of the information needed to use ABIRATERONE ACETATE TABLETS safely and effectively. See full prescribing information for ABIRATERONE ACETATE TABLETS.

ABIRATERONE ACETATE tablets, for oral use
Initial U.S. Approval: 2011

RECENT MAJOR CHANGES
Warnings and Precautions (5.1) 06/2019
Warnings and Precautions (5.4) 06/2019
Warnings and Precautions (5.5) 06/2019

INDICATIONS AND USAGE:
Abiraterone acetate tablets are a CYP17 inhibitor indicated in combination with prednisone for the treatment of patients with metastatic castration-resistant prostate cancer (CRPC).
* * * * *

DOSE AND ADMINISTRATION
Metastatic castration-resistant prostate cancer:
2. Abiraterone acetate tablets 1,000 mg orally once daily with prednisone 5 mg orally twice daily (2.1)

Patients receiving abiraterone acetate tablets should also receive a gonadotropin-releasing hormone (GnRH) analog concurrently or should have had bilateral orchiectomy. Abiraterone acetate tablets must be taken on an empty stomach with water at least one hour before or 2 hours after a meal. Do not crush or chew tablets. (2.2)
Dose Modification:
3. For patients with baseline moderate hepatic impairment (Child-Pugh Class B), reduce the abiraterone acetate starting dose to 250 mg once daily. (2.4)

4. For patients who develop hepatotoxicity during treatment, hold abiraterone acetate tablets until recovery. Re-treatment may be initiated at a reduced dose. Abiraterone acetate tablets should be discontinued if patients develop severe hepatotoxicity. (2.4)
* * * * *

CONTRAINDICATIONS
Uncoated Tablet 250 mg (3)
None (4)

WARNINGS AND PRECAUTIONS

1. Mineralocorticoid Excess: Classify moderate patients with cardiovascular disease. Control hypertension and serum hypokalemia before treatment. Monitor blood pressure, serum potassium and symptoms of fluid retention at least monthly. (5.1)
2. Adrenocortical Insufficiency: Monitor for symptoms and signs of adrenocortical insufficiency. Increased dosage of corticosteroids may be indicated before, during and after steroid withdrawal. (5.2)
3. Hepatotoxicity: Can be severe and fatal. Monitor liver function and modify, interrupt, or discontinue abiraterone acetate therapy as recommended. (5.3)
4. Increased Fractures and Mortality in Combination with radium Ra 223 dichloride: Use of abiraterone acetate plus prednisone/prednisolone in combination with radium Ra 223 dichloride is not recommended. (5.4)
5. Embryo-Fetal Toxicity: Abiraterone acetate can cause fetal harm. Advise males with female partners of reproductive potential to use effective contraception. (5.5, 6.1, 6.3)

ADVERSE REACTIONS
The most common adverse reactions (≥10%) are fatigue, arthralgia, hypertension, nausea, edema, hypokalemia, hot flash, diarrhea, vomiting, upper respiratory infection, cough, and headache. (6.1)
The most common laboratory abnormalities (≥10%) are anemia, elevated alkaline phosphatase, hypertriglyceridemia, hyperkalemia, hypokalemia, and hypocalcemia. (6.1)
To report SUSPECTED ADVERSE REACTIONS, contact AstraZeneca Corp. at 1-800-784-5375 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS
CYP3A4 Inhibitors: Avoid concomitant strong CYP3A4 inhibitors during abiraterone acetate treatment. If a strong CYP3A4 inducer must be co-administered, increase the abiraterone acetate dosing frequency. (2.5, 7.3)
CYP3A4 Substrates: Avoid co-administration of abiraterone acetate with CYP3A4 substrates that have a narrow therapeutic index. If an abiraterone treatment cannot be used, exercise caution and consider a dose reduction of the concomitant CYP3A4 substrate. (7.2)
* * * * *

Do not use abiraterone acetate tablets in patients with baseline severe hepatic impairment (Child-Pugh Class C). (6.6)
See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling. Revised: 7/20/20

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
2.1 Recommended Dose for Metastatic CRPC
2.2 Important Administration Instructions
2.3 Dose Modification Guidelines in Hepatic Impairment and Hepatotoxicity
2.4 Dose Modification Guidelines for Strong CYP3A4 Inducers
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
5.1 Hypokalemia, Fluid Retention, and Cardiovascular Adverse Reactions due to Mineralocorticoid Excess
5.2 Adrenocortical Insufficiency
5.3 Hepatotoxicity
5.4 Increased Fractures and Mortality in Combination with Radium Ra 223 Dichloride
5.5 Embryo-Fetal Toxicity
- 6 ADVERSE REACTIONS
6.1 Clinical Trial Experience
6.2 Postmarketing Experience
- 7 DRUG INTERACTIONS
7.1 Drugs that Inhibit or Induce CYP3A4 Enzymes
7.2 Effects of Abiraterone on Drug Metabolizing Enzymes
- 8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Patients with Hepatic Impairment
8.7 Patients with Renal Impairment
- 9 OVERDOSAGE
- 10 OVERDOSAGE
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility
13.2 Animal Toxicology and/or Pharmacology
- 14 CLINICAL STUDIES
- 15 HOW SUPPLIED/STORAGE AND HANDLING
- 16 PATIENT COUNSELING INFORMATION
* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE
Abiraterone acetate tablets are indicated in combination with prednisone for the treatment of patients with

1. Metastatic castration-resistant prostate cancer (CRPC).

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose for Metastatic CRPC
The recommended dose of abiraterone acetate tablets is 1,000 mg (four 250 mg tablets) orally once daily with prednisone 5 mg orally twice daily.

2.2 Important Administration Instructions
Patients receiving abiraterone acetate tablets should also receive a gonadotropin-releasing hormone (GnRH) analog concurrently or should have had bilateral orchiectomy. Abiraterone acetate tablets must be taken on an empty stomach, at least one hour before or at least two hours after a meal [see Clinical Pharmacology (7.2.3)]. The tablets should be swallowed whole with water. Do not crush or chew tablets.

2.3 Dose Modification Guidelines in Hepatic Impairment and Hepatotoxicity

Hepatic Impairment
In patients with baseline moderate hepatic impairment (Child-Pugh Class B), reduce the recommended dose of abiraterone acetate to 250 mg once daily. In patients with moderate hepatic impairment (Child-Pugh Class B), ALT, AST, and bilirubin prior to the start of treatment, every week for the first month, every two weeks for the following two months of treatment and monthly thereafter. If elevations in ALT and/or AST greater than 5X upper limit of normal (ULN) or total bilirubin greater than 3X ULN occur in patients with baseline moderate hepatic impairment, discontinue abiraterone acetate tablets and do not re-treat patients with abiraterone acetate tablets. [see Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)].

Do not use abiraterone acetate tablets in patients with baseline severe hepatic impairment (Child-Pugh Class C).

Hepatotoxicity
For patients who develop hepatotoxicity during treatment with abiraterone acetate tablets (ALT and/or AST greater than 5X ULN or total bilirubin greater than 3X ULN), interrupt treatment with abiraterone acetate tablets [see Warnings and Precautions (5.3)]. Treatment may be restarted at a reduced dose of 750 mg once daily following return of liver function tests to the patient's baseline or to AST and ALT less than or equal to 2.5X ULN and total bilirubin less than or equal to 1.5X ULN. For patients who resume treatment, monitor serum transaminases and bilirubin at a minimum of every two weeks for three months and monthly thereafter.

If hepatotoxicity recurs at the dose of 750 mg once daily, re-treatment may be restarted at a reduced dose of 500 mg once daily following return of liver function tests to the patient's baseline or to AST and ALT less than or equal to 2.5X ULN and total bilirubin less than or equal to 1.5X ULN.

If hepatotoxicity recurs at the reduced dose of 500 mg once daily, discontinue treatment with abiraterone acetate tablets. Permanently discontinue abiraterone acetate for patients who develop a concurrent elevation of ALT greater than 3 x ULN and total bilirubin greater than 2 x ULN in the absence of biliary obstruction or other causes responsible for the concurrent elevation [see Warnings and Precautions (5.3)].

2.4 Dose Modification Guidelines for Strong CYP3A4 Inducers
Avoid concomitant strong CYP3A4 inducers (e.g., phenytoin, carbamazepine, rifampin, rifabutin, rifapentine, phenobarbital) during abiraterone acetate treatment.

If a strong CYP3A4 inducer must be co-administered, increase the abiraterone acetate dosing frequency by twice a day only during the co-administration period (e.g., from 1,000 mg once daily to 1,000 mg twice a day). Reduce the dose back to the previous dose and frequency, if the concomitant strong CYP3A4 inducer is discontinued [see Drug Interactions (7.1) and Clinical Pharmacology (12.3)].

3 DOSAGE FORMS AND STRENGTHS
Abiraterone acetate 250 mg tablets, USP are white to off-white, oval, biconvex uncoated tablets. Engraved "AZ50" on one side, "APO" on the other side.

4 CONTRAINDICATIONS
None.

5 WARNINGS AND PRECAUTIONS

5.1 Hypokalemia, Fluid Retention, and Cardiovascular Adverse Reactions due to Mineralocorticoid Excess
Abiraterone acetate may cause hypertension, hypokalemia, and fluid retention as a consequence of increased mineralocorticoid levels resulting from CYP17 inhibition [see Clinical Pharmacology (12.1)]. Monitor patients for hypertension, hypokalemia, and fluid retention at least once a month.

Control hypertension and correct hypokalemia before and during treatment with abiraterone acetate. In the combined data from 4 placebo-controlled trials using prednisone 5 mg twice daily in combination with 1,000 mg abiraterone acetate daily, grades 3-4 hypokalemia were observed in 4% of patients on the abiraterone acetate arm and 2% of patients on the placebo arm. Grades 3 or 4 hypertension were observed in 2% of patients each arm and grades 3 or 4 fluid retention in 1% of patients each arm.

Caution must be exercised in patients whose underlying medical conditions might be compromised by increases in blood pressure, hypokalemia or fluid retention, such as those with heart failure, recent myocardial infarction, cardiovascular disease, or ventricular arrhythmia. In postmarketing experience, QT prolongation and Torsades de Pointes have been observed in patients who develop hypokalemia while taking abiraterone acetate.

The safety of abiraterone acetate in patients with left ventricular ejection fraction <50% or New York Heart Association (NYHA) Class III or IV heart failure (in COU-AA-301 or NYHA Class II to IV heart failure (in COU-AA-302) has not been established because these patients were excluded from these randomized clinical trials [see Clinical Studies (14)].

5.2 Adrenocortical Insufficiency
Adrenal insufficiency occurred in 0.3% of 2230 patients taking abiraterone acetate and in 0.1% of 1763 patients taking placebo in the combined data of the randomized, placebo-controlled clinical studies. Adrenocortical insufficiency was reported in patients receiving abiraterone acetate in combination with prednisone, following interruption of daily steroids and/or with concurrent infection or stress. Monitor patients for symptoms and signs of adrenocortical insufficiency, particularly if patients are withdrawn from prednisone, have prednisone dose reductions, or experience unusual stress. Symptoms

and signs of adrenal insufficiency may be masked by adverse reaction associated with mineralocorticoid excess seen in patients treated with abiraterone acetate. If clinically indicated, perform appropriate tests to confirm the diagnosis of adrenal insufficiency. Increased dosage of corticosteroids may be indicated before, during and after surgical situations. *See Warnings and Precautions (5.1).*

5.3 Hepatotoxicity

In postmarketing experience, there have been abiraterone acetate-associated severe hepatic toxicity, including fulminant hepatitis, acute liver failure and death. *See Adverse Reactions (6.2).*

In the combined data of randomized clinical trials, grade 3 to 4 ALT or AST increases (at least 5X ULN) were reported in 6% of 2230 patients who received abiraterone acetate, typically during the first 3 months after starting treatment. Patients whose baseline ALT or AST were elevated were more likely to experience liver test elevation than those beginning with normal values. Treatment discontinuation due to ALT and AST increases or abnormal hepatic function occurred in 1.1% of 2230 patients taking abiraterone acetate. In these clinical trials, no deaths clearly related to abiraterone acetate were reported due to hepatotoxicity events.

Measure serum transaminases (ALT and AST) and bilirubin levels prior to starting treatment with abiraterone acetate, every two weeks for the first three months of treatment and monthly thereafter. In patients with baseline moderate hepatic impairment receiving a reduced abiraterone acetate dose of 250 mg, measure ALT, AST, and bilirubin prior to the start of treatment, every week for the first month, every two weeks for the following two months of treatment and monthly thereafter. Promptly measure serum bilirubin, AST, and ALT if clinical symptoms or signs suggestive of hepatotoxicity develop. Elevation of AST, ALT, or bilirubin from the patient's baseline should prompt more frequent monitoring. If at any time AST or ALT rise above five times the ULN, or the bilirubin rises above three times the ULN, interrupt abiraterone acetate treatment and closely monitor liver function.

Re-treatment with abiraterone acetate at a reduced dose level may take place only after return of liver function tests to the patient's baseline or to AST and ALT 1.5 times or equal to 2.5X ULN and total bilirubin less than or equal to 1.5X ULN. *See Dosage and Administration (2.4).*

Permanently discontinue abiraterone acetate tablets for patients who develop a concurrent elevation of ALT greater than 3x ULN and total bilirubin greater than 2x ULN in the absence of biliary obstruction or other causes responsible for the concurrent elevation. *See Dosage and Administration (2.4).*

The safety of abiraterone acetate treatment of patients who develop AST or ALT greater than or equal to 20X ULN and/or bilirubin greater than or equal to 10X ULN is unknown.

5.4 Increased Fractures and Mortality in Combination with Radium Ra 223 Dichloride

Abiraterone acetate plus prednisone/prednisolone is not recommended for use in combination with radium 223 dichloride outside of clinical trials.

The clinical efficacy and safety of concurrent initiation of abiraterone acetate plus prednisone/prednisolone and radium Ra 223 dichloride was assessed in a randomized, placebo-controlled multicenter study (EKA-223 trial) in 806 patients with asymptomatic or mildly symptomatic castration-resistant prostate cancer with bone metastases. The study was unblinded early based on an Independent Data Monitoring Committee recommendation.

In the primary analysis, increased incidences of fractures (28.6% vs 11.4%) and deaths (28.2% vs 15.5%) have been observed in patients who received abiraterone acetate plus prednisone/prednisolone in combination with radium Ra 223 dichloride compared to patients who received placebo in combination with abiraterone acetate plus prednisone/prednisolone.

5.5 Embryo-Fetal Toxicity

The safety and efficacy of abiraterone acetate have not been established in females. Based on animal reproductive studies and mechanism of action, abiraterone acetate can cause fetal harm and loss of pregnancy when administered to a pregnant female. In animal reproduction studies, oral administration of abiraterone acetate to pregnant rats during organogenesis caused adverse developmental effects at maternal exposures approximately 0.03 times the human exposure (AUC) at the recommended dose. Advise males with female partners of reproductive potential to use effective contraception during treatment with abiraterone acetate and for 3 weeks after the last dose of abiraterone acetate. Use in Specific Populations (8.1, 8.3). Abiraterone acetate should not be handled by females who are or may become pregnant. *See How Supplied/Storage and Handling (16.1).*

6 ADVERSE REACTIONS

The following are discussed in more detail in other sections of the labeling:

- Hypokalemia, Fluid Retention, and Cardiovascular Adverse Reaction due to Mineralocorticoid Excess. *See Warnings and Precautions (5.1).*
- Adrenocortical Insufficiency. *See Warnings and Precautions (5.2).*
- Hepatotoxicity. *See Warnings and Precautions (5.3).*
- Increased Fractures and Mortality in Combination with Radium Ra 223 Dichloride. *See Warnings and Precautions (5.4).*

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Two randomized placebo-controlled, multicenter clinical trials (COU-AA-301 and COU-AA-302) enrolled patients who had metastatic CRPC in which abiraterone acetate was administered orally at a dose of 1200 mg daily in combination with prednisone 5 mg twice daily in the active treatment arm. Placebo plus prednisone 5 mg twice daily was given to patients on the control arm. Another randomized placebo-controlled, multicenter clinical trial enrolled patients who had another indication in which abiraterone acetate tablets was administered in combination with prednisone. Placebos were administered to patients in the control arm. Additionally, two other randomized, placebo-controlled trials were conducted in patients with metastatic CRPC. The safety data pooled from 2230 patients in the randomized controlled trials constitute the basis for the data presented in the Warnings and Precautions, Grade 1 to 4 adverse reactions, and Grade 1 to 4 laboratory abnormalities. In all trials, gynecologic releasing hormone (GnRH) analog or prior orchiectomy was required in both arms.

In the pooled data, median treatment duration was 11 months (9.1, 43) for abiraterone acetate-treated patients and 7.2 months (0.1, 43) for placebo-treated patients. The most common adverse reactions (≥10%) that occurred more commonly (>2%) in the abiraterone acetate arm were fatigue, arthralgia, hypertension, nausea, edema, hypokalemia, hot flash, diarrhea, vomiting, upper respiratory infection, cough, and headache. The most common laboratory abnormalities (>20%) that occurred more commonly (>2%) in the abiraterone acetate arm were anemia, elevated alkaline phosphatase, hyperglycemia, lymphopenia, hypochlorocholesterolemia, hypoglycemia, and hypokalemia. Grades 3 to 4 adverse events were reported for 53% of patients in the abiraterone acetate arm and 46% of patients in the placebo arm. Treatment discontinuation was reported in 14% of patients in the abiraterone acetate arm and 13% of patients in the placebo arm. The common adverse events (≥1%) resulting in discontinuation of abiraterone acetate and prednisone were hepatotoxicity and cardiac disorders.

Deaths associated with treatment-emergent adverse events were reported for 7.5% of patients in the abiraterone acetate arm and 6.6% of patients in the placebo arm. Of the patients in the abiraterone acetate arm, the most common cause of death was disease progression (3.9%). Other reported causes of death in 5 patients included pneumonia, cardio-respiratory arrest, death (no additional information), and general physical health deterioration.

COU-AA-301: Metastatic CRPC Following Chemotherapy

COU-AA-301 enrolled 1195 patients with metastatic CRPC who had received prior docetaxel chemotherapy. Patients were not eligible if AST and/or ALT >2.5X ULN in the absence of liver metastases. Patients with liver metastases were excluded if AST and/or ALT >5X ULN.

Table 1 shows adverse reactions on the abiraterone acetate arm in COU-AA-301 that occurred with a ≥2% absolute increase in frequency compared to placebo or were events of special interest. The median duration of treatment with abiraterone acetate with prednisone was 8 months.

System/Organ Class	Abiraterone Acetate with Prednisone (N=791)		Placebo with Prednisone (N=394)	
	All Grades ¹	Grade 3-4	All Grades	Grade 3-4
Adverse reaction	8	6	6	6
Musculoskeletal and connective tissue disorders				
Back pain/discomfort ²	30	23	25	14
Muscle discomfort ³	26	3.0	23	2.3
General disorders				
Edema ⁴	27	1.9	18	1.8
Vascular disorders				
Hot flash	29	2.3	17	0.3
Hypertension	15	1.3	15	0.3
Constitutional disorders				
Fatigue	18	0.6	14	0.3
Dyspepsia	11	0	13	0
Infections and infestations				
Upper respiratory tract infection	54	0	25	0
Respiratory, thoracic and mediastinal disorders				
Cough	31	0	12	0
Renal and urinary disorders				
Urinary frequency	72	0.3	51	0.3
Nocturia	62	0	41	0
Injury, poisoning and procedural complications				
Fracture ⁵	29	1.4	23	0
Cardiac disorders				
Arrhythmia ⁶	72	1.1	46	0
Acute pain chest discomfort ⁷	18	0.5	12	0
Cardiac failure ⁸	13	1.9	10	0.3

¹Adverse events graded according to CTCAE version 3.0.

²Includes terms: Arthritis, Arthralgia, Joint swelling, and Joint stiffness.

³Includes terms: Muscle spasms, Musculoskeletal pain, Myalgia, Musculoskeletal discomfort, and Musculoskeletal stiffness.

⁴Includes terms: Edema, Edema peripheral, Pitting edema, and Generalized edema.

⁵Includes all fractures with the exception of pathological fracture.

⁶Includes terms: Atrial fibrillation, Tachycardia, Atrial fibrillation, Supraventricular tachycardia, Atrial tachycardia, Ventricular tachycardia, Atrial flutter, Bradycardia, Atrioventricular block complete, Conduction disorder, and Bradycardia.

⁷Includes terms: Angina pectoris, Chest pain, and Angina unstable. Myocardial infarction or ischemia occurred more commonly in the placebo arm than in the abiraterone acetate arm (1.3% vs 1.1% respectively).

⁸Includes terms: Cardiac failure, Cardiac failure congestive, Left ventricular dysfunction, Cardiogenic shock, Cardiomegaly, Cardiomyopathy, and Ejection fraction decreased.

Table 2 shows laboratory abnormalities of interest from COU-AA-301.

Table 2: Laboratory Abnormalities of Interest in COU-AA-301

Laboratory Abnormality	Abiraterone Acetate with Prednisone (N=791)		Placebo with Prednisone (N=394)	
	All Grades	Grade 3-4 (%)	All Grades	Grade 3-4 (%)
Hyperglycemia	13	0.4	33	0
High AST	11	0.1	16	0
Hypokalemia	28	5.3	20	0
Hypophosphatemia	14	0.2	16	0
High ALT	11	0.4	10	0
High Total Bilirubin	6.6	0.1	4.6	0

COU-AA-302: Metastatic CRPC Prior to Chemotherapy

COU-AA-302 enrolled 1088 patients with metastatic CRPC who had not received prior cytotoxic chemotherapy. Patients were ineligible if AST and/or ALT >2.5X ULN and patients were excluded if they had liver metastases.

Table 3 shows adverse reactions on the abiraterone acetate arm in COU-AA-302 that occurred in ≥5% of patients with a ≥2% absolute increase in frequency compared to placebo. The median duration of treatment with abiraterone acetate with prednisone was 13.8 months.

System/Organ Class	Abiraterone Acetate with Prednisone (N=542)		Placebo with Prednisone (N=546)	
	All Grades ¹	Grade 3-4	All Grades	Grade 3-4
Adverse reaction	8	6	6	6
General disorders				
Fatigue	35	2.2	31	1.2
Edema ²	16	0.4	14	0.1
Pruritus	17	0.6	15	0.2
Musculoskeletal and connective tissue disorders				

System Organ Class	Abiraterone Acetate	Placebo	Grade 1-2	Grade 3-4
Most common adverse events	10	7.0	1.5	0.0
Headache	16	1.4	1.1	0.7
Gastrointestinal disorders				
Constipation	15	1.4	1.0	1.6
Diarrhea	15	0.0	1.0	0.0
Dyspepsia	11	0.0	1.0	0.2
Vascular disorders				
Hypotension	12	0.2	0.0	0.0
Hypertension	12	1.0	1.3	0.0
Respiratory, thoracic and mediastinal disorders				
Cough	17	0.0	1.4	0.2
Dyspnea	12	2.4	0.6	0.0
Psychiatric disorders				
Insomnia	14	1.2	1.1	0.0
Other, poisoning and procedural complications				
Cellulitis	10	0.0	1.1	0.0
Pain	10	0.0	1.3	0.0
Infections and infestations				
Upper respiratory tract infection	13	0.0	0.0	0.0
Nasopharyngitis	11	0.0	0.1	0.0
Renal and urinary disorders				
Urinary tract infection	10	1.1	1.0	1.6
Skin and subcutaneous tissue disorders				
Rash	11	0.0	1.7	0.0

¹Adverse events graded according to CTCAE version 3.0.

²Includes terms Edema peripheral, Pitting edema, and Generalized edema.

³Includes terms Arthritis, Arthralgia, Joint swelling, and Joint stiffness.

Table 4 shows laboratory abnormalities that occurred in greater than 15% of patients, and more frequently (>5%) in the abiraterone acetate arm compared to placebo in COU-AA-302.

Laboratory Abnormality	Abiraterone Acetate with Prednisone (N=540)	Placebo with Prednisone (N=540)
Hypokalemia	30	24
Hyperkalemia	37	31
High ALT	42	29
High AST	37	29
Hypertension	12	12
Hypohidrosis	17	10

¹Based on non-fasting blood draws.

Cardiovascular Adverse Reactions

In the combined data of randomized, placebo-controlled clinical studies, cardiac failure occurred more commonly in patients on the abiraterone acetate arm compared to patients on the placebo arm (2.6% versus 0.9%). Grade 1 to 4 cardiac failure occurred in 3.3% of patients taking abiraterone acetate and led to 5 treatment discontinuations and 4 deaths. Grade 1 to 4 cardiac failure occurred in 0.2% of patients taking placebo. There were no treatment discontinuations and two deaths due to cardiac failure in the placebo group.

In the same combined data, the majority of arrhythmias were grade 1 or 2. There was one death associated with arrhythmia and three patients with sudden death in the abiraterone acetate arm and five deaths in the placebo arm. There were 7 (0.2%) deaths due to cardiorespiratory events in the abiraterone acetate arm and 2 (0.1%) deaths in the placebo arm. Myocardial ischemia or myocardial infarction led to death in 3 patients in the placebo arm and 3 deaths in the abiraterone acetate arm.

6.2 Postmarketing Experience

The following additional adverse reactions have been identified during post approval use of abiraterone acetate with prednisone. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Respiratory, Thoracic and Mediastinal Disorders: non-infectious pneumonitis.

Musculoskeletal and Connective Tissue Disorders: myopathy, including rhabdomyolysis.

Hepatobiliary Disorders: fulminant hepatitis, including acute hepatic failure and death.

Cardiac Disorders: QT prolongation and Torsades de Pointes (observed in patients who developed hypokalemia or had underlying cardiovascular conditions).

7 DRUG INTERACTIONS

7.1 Drugs that Inhibit or Induce CYP3A4 Enzymes

Based on *in vitro* data, abiraterone acetate is a substrate of CYP3A4.

In a dedicated drug interaction trial, co-administration of rifampin, a strong CYP3A4 inducer, decreased exposure of abiraterone by 55%. Avoid concomitant strong CYP3A4 inducers during abiraterone acetate treatment. If a strong CYP3A4 inducer must be co-administered, increase the abiraterone acetate dosing frequency (see Dosage and Administration (2.3) and Clinical Pharmacology (12.3)).

In a dedicated drug interaction trial, co-administration of ketoconazole, a strong inhibitor of CYP3A4, had no clinically meaningful effect on the pharmacokinetics of abiraterone (see Clinical Pharmacology (12.3)).

7.2 Effects of Abiraterone on Drug Metabolizing Enzymes

Abiraterone acetate is an inhibitor of the hepatic drug-metabolizing enzymes CYP2D6 and CYP2C8. In a CYP2D6 drug-drug interaction trial, the C_{max} and AUC of dextropropriofen (CYP2D6 substrate) were increased 2.6- and 2.8-fold, respectively, when dextropropriofen was given with abiraterone acetate 1,000 mg daily and prednisone 5 mg twice daily. Avoid co-administration of abiraterone acetate with substrates of CYP2D6 with a narrow therapeutic index (e.g., antidepressants). If abiraterone treatment cannot be used, consider a dose reduction of the concomitant CYP2D6 substrate drug (see Clinical Pharmacology (12.3)).

In a CYP2C8 drug-drug interaction trial in healthy subjects, the AUC of pioglitazone (CYP2C8 substrate) was increased by 46% when pioglitazone was given together with a single dose of 1,000 mg abiraterone acetate. Therefore, patients should be monitored closely for signs of toxicity related to a CYP2C8 substrate with a narrow therapeutic index if used concomitantly with abiraterone acetate (see Clinical Pharmacology (12.3)).

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

The safety and efficacy of abiraterone acetate have not been established in females. Based on findings from animal studies and the mechanism of action, abiraterone acetate can cause fetal harm and potential loss of pregnancy.

There are no human data on the use of abiraterone acetate in pregnant women. In animal reproduction studies, oral administration of abiraterone acetate to pregnant rats during organogenesis caused adverse developmental effects at maternal exposures approximately ≥ 0.03 times the human exposure (AUC) at the recommended dose (see Data).

Data

Animal Data

In an embryo-fetal developmental toxicity study in rats, abiraterone acetate caused developmental toxicity when administered at oral doses of 10, 30 or 100 mg/kg/day throughout the period of organogenesis (gestational days 6 to 17). Findings included embryo-fetal lethality (increased post implantation loss and resorptions) and decreased number of live fetuses, fetal developmental delay (delayed effects) and urogenital effects (bilateral ureter dilation) at doses ≥ 10 mg/kg/day, decreased fetal ano-genital distance at 20 mg/kg/day, and decreased fetal body weight at 100 mg/kg/day. Doses ≥ 10 mg/kg/day caused maternal toxicity. The doses tested in rats resulted in systemic exposures (AUC) approximately 0.03, 0.1 and 0.3 times, respectively, the AUC in patients.

8.2 Lactation

Risk Summary

The safety and efficacy of abiraterone acetate have not been established in females. There is no information available on the presence of abiraterone acetate in human milk, or on the effects on the breastfed child or milk production.

8.3 Females and Males of Reproductive Potential

Contraception

Males

Based on findings in animal reproduction studies and its mechanism of action, advise males with female partners of reproductive potential to use effective contraception during treatment and for 3 weeks after the final dose of abiraterone acetate (see Use in Specific Populations (8.1)).

Infertility

Based on animal studies, abiraterone acetate may impair reproductive function and fertility in males of reproductive potential (see Nonclinical Toxicology (13.1)).

8.4 Pediatric Use

Safety and effectiveness of abiraterone acetate in pediatric patients have not been established.

8.5 Geriatric Use

Of the total number of patients receiving abiraterone acetate in randomized clinical trials, 70% of patients were 65 years and over and 27% were 75 years and over. No overall differences in safety or effectiveness were observed between these elderly patients and younger patients. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

8.6 Patients with Hepatic Impairment

The pharmacokinetics of abiraterone were examined in subjects with baseline mild (N=8) or moderate (N=8) hepatic impairment (Child-Pugh Class A and B, respectively) and in 8 healthy control subjects with normal hepatic function. The systemic exposure (AUC) of abiraterone after a single oral 1,000 mg dose of abiraterone acetate increased by approximately 1.1-fold and 3.6-fold in subjects with mild and moderate baseline hepatic impairment, respectively compared to subjects with normal hepatic function. In another trial, the pharmacokinetics of abiraterone were examined in subjects with baseline severe (N=8) hepatic impairment (Child-Pugh Class C) and in 8 healthy control subjects with normal hepatic function. The systemic exposure (AUC) of abiraterone increased by approximately 7-fold and the fraction of free drug increased 2-fold in subjects with severe baseline hepatic impairment compared to subjects with normal hepatic function.

No dosage adjustment is necessary for patients with baseline mild hepatic impairment. In patients with baseline moderate hepatic impairment (Child-Pugh Class B), reduce the recommended dose of abiraterone acetate to 250 mg once daily. Do not use abiraterone acetate in patients with baseline severe hepatic impairment (Child-Pugh Class C). If elevations in ALT, AST, SGPT, ULN or total bilirubin $>3\times$ ULN occur in patients with baseline moderate hepatic impairment, discontinue abiraterone acetate treatment (see Dosage and Administration (2.4) and Clinical Pharmacology (12.3)).

For patients who develop hepatotoxicity during treatment, interruption of treatment and dosage adjustment may be required (see Dosage and Administration (2.4), Warnings and Precautions (5.3), and Clinical Pharmacology (12.3)).

8.7 Patients with Renal Impairment

No dosage adjustment is necessary for patients with renal impairment (see Clinical Pharmacology (12.3)).


10 OVERDOSAGE

Human experience of overdose with abiraterone acetate is limited.

There is no specific antidote. In the event of an overdose, stop abiraterone acetate, undertake general supportive measures, including monitoring for arrhythmias and cardiac failure and assess liver function.

11 DESCRIPTION

Abiraterone acetate, USP is the active ingredient of abiraterone acetate tablets, USP. USP is the acetyl ester of abiraterone. Abiraterone is an inhibitor of CYP17 (17 α -hydroxylase/C17,20-lyase). Each abiraterone acetate tablet, USP contains 250 mg of abiraterone acetate, USP. Abiraterone acetate, USP is designed chemically as (3S)-17-(3-pyridinyl) androst-5,16-dien-3-yl acetate and its structure is:



Abiraterone acetate, USP is a white to off-white, non-hygroscopic, crystalline powder. Its molecular formula is $C_{26}H_{34}N_2O_2$ and it has a molecular weight of 391.55 g/mol. Abiraterone acetate, USP is a lipophilic compound with an octanol-water partition coefficient of 5.12 (Log P) and is practically insoluble in water. The pKa of the aromatic nitrogen is 5.19.

Abiraterone acetate tablets are available in 250 mg uncoated tablets with the following inactive ingredients: colloidal silicon dioxide, croscopollose, lactose monohydrate, magnesium stearate, and sodium lauryl sulfate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Abiraterone acetate is converted in vivo to abiraterone, an androgen biosynthesis inhibitor, that inhibits 17 α -hydroxylase/C17,20-lyase (CYP17). This enzyme is expressed in testicular, adrenal, and prostatic tumor tissues and is required for androgen biosynthesis. CYP17 catalyzes two sequential reactions: 1) the conversion of pregnenolone and progesterone to their 17 α -hydroxy derivatives by 17 α -hydroxylase activity and 2) the subsequent formation of dehydroepiandrosterone (DHEA) and androstenedione, respectively, by CYP17 20-lyase activity. DHEA and androstenedione are androgens and are precursors of testosterone. Inhibition of CYP17 by abiraterone can also result in increased mineralocorticoid production by the adrenals (see Warnings and Precautions (5.1)).

Androgen sensitive prostatic carcinoma responds to treatment that decreases androgen levels. Androgen deprivation therapies, such as treatment with GnRH agonists or orchiectomy, decrease androgen production in the testes but do not affect androgen production by the adrenals or in the tumor.

Abiraterone acetate decreased serum testosterone and other androgen in patients in the placebo-controlled clinical trial. It is not necessary to monitor the effect of abiraterone acetate on serum testosterone levels.

Changes in serum prostate specific antigen (PSA) levels may be observed but have not been shown to correlate with clinical benefit in individual patients.

12.2 Pharmacodynamics

Cardiac Electrophysiology

In a multi-center, open-label, single-arm trial, 33 patients with metastatic CRPC received abiraterone acetate orally at a dose of 1,000 mg once daily at least 1 hour before or 2 hours after a meal in combination with prednisone 5 mg orally twice daily. Assessments up to Cycle 2 Day 2 showed no large changes in the QTc interval (i.e., >20 ms) from baseline. However, small increases in the QTc interval (i.e., <10 ms) due to abiraterone acetate cannot be excluded due to study design limitations.

12.3 Pharmacokinetics

Following administration of abiraterone acetate, the pharmacokinetics of abiraterone and abiraterone acetate have been studied in healthy subjects and in patients with metastatic CRPC. In vivo, abiraterone acetate is converted to abiraterone. In clinical studies, abiraterone acetate plasma concentrations were below detectable levels (<0.2 ng/mL) in >99% of the analyzed samples.

Absorption

Following oral administration of abiraterone acetate to patients with metastatic CRPC, the median time to reach maximum plasma abiraterone concentrations is 2 hours. Abiraterone accumulation is observed at steady-state, with a 2-fold higher exposure (steady-state AUC) compared to a single 1,000 mg dose of abiraterone acetate.

At the dose of 1,000 mg daily in patients with metastatic CRPC, steady-state values (mean \pm SD) of C_{max} were 226 \pm 178 ng/mL and of AUC were 993 \pm 629 ng \cdot h/mL. No major deviation from dose proportionality was observed in the dose range of 250 mg to 1,000 mg. However, the exposure was not significantly increased when the dose was doubled from 1,000 to 2,000 mg (8% increase in the mean AUC).

Systemic exposure of abiraterone is increased when abiraterone acetate is administered with food. In healthy subjects abiraterone C_{max} and AUC $_{0-\infty}$ were approximately 7- and 5-fold higher, respectively, when a single dose of abiraterone acetate was administered with a low-fat meal (7% fat, 300 calories) and approximately 17- and 10-fold higher, respectively, when a single dose of abiraterone acetate was administered with a high-fat (57% fat, 825 calories) meal compared to overnight fasting. Abiraterone AUC $_{0-\infty}$ was approximately 7-fold or 1.6-fold higher, respectively, when a single dose of abiraterone acetate was administered 2 hours after or 1 hour before a medium-fat meal (25% fat, 491 calories) compared to overnight fasting.

Systemic exposures of abiraterone in patients with metastatic CRPC, after repeated dosing of abiraterone acetate were similar when abiraterone acetate was taken with low-fat meals for 7 days and increased approximately 2-fold when taken with high-fat meals for 7 days compared to when taken at least 2 hours after a meal and at least 1 hour before a meal for 7 days.

Given the normal variation in the content and composition of meals, taking abiraterone acetate with meals has the potential to result in increased and highly variable exposures. Therefore, abiraterone acetate must be taken on an empty stomach, at least one hour before or at least two hours after a meal. The tablets should be swallowed whole with water (see Dosage and Administration (2.3)).

Distribution and Protein Binding

Abiraterone is highly bound (>99%) to the human plasma protein, albumin and alpha-1 acid glycoprotein. The apparent steady-state volume of distribution (mean \pm SD) is 15,660 \pm 12,258 L. In vitro studies show that at clinically relevant concentrations, abiraterone acetate and abiraterone are not substrates of P-glycoprotein (P-gp) and that abiraterone acetate is an inhibitor of P-gp.

Metabolism

Following oral administration of ^{14}C -abiraterone acetate as capsules, abiraterone acetate is hydrolyzed to abiraterone (active metabolite). The conversion is likely through esterase activity (the esterases have not been identified) and is not CYP mediated. The two main circulating metabolites of abiraterone in human plasma are abiraterone sulphate (inactive) and N-oxide abiraterone sulphate (inactive), which account for about 47% of exposure each. CYP3A4 and SULT2A1 are the enzymes involved in the formation of N-oxide abiraterone sulphate and SULT2A1 is involved in the formation of abiraterone sulphate.

Excretion

In patients with metastatic CRPC, the mean terminal half-life of abiraterone in plasma (mean \pm SD) is 12 \pm 5 hours. Following oral administration of ^{14}C -abiraterone acetate, approximately 88% of the radioactive dose is recovered in feces and approximately 5% in urine. The major components present in feces are unchanged abiraterone acetate and abiraterone (approximately 55% and 22% of the administered dose, respectively).

Patients with Hepatic Impairment

The pharmacokinetics of abiraterone was examined in subjects with baseline mild (N=8) or moderate (N=8) hepatic impairment (Child-Pugh Class A and B, respectively) and in 8 healthy control subjects with normal hepatic function. Systemic exposure to abiraterone after a single oral 1,000 mg dose given under fasting conditions increased approximately 1.4-fold and 3.6-fold in subjects with mild and moderate baseline hepatic impairment, respectively. The mean half-life of abiraterone is prolonged to approximately 18 hours in subjects with mild hepatic impairment and to approximately 19 hours in subjects with moderate hepatic impairment.

In another trial, the pharmacokinetics of abiraterone were examined in subjects with baseline severe (N=8) hepatic impairment (Child-Pugh Class C) and in 8 healthy control subjects with normal hepatic function. The systemic exposure (AUC) of abiraterone increased by approximately 7-fold in subjects with severe baseline hepatic impairment compared to subjects with normal hepatic function. In addition, the mean protein binding was found to be lower in the severe hepatic impairment group compared to the normal hepatic function group, which resulted in a two-fold increase in the fraction of free drug in patients with severe hepatic impairment (see Dosage and Administration (2.4) and Use in Specific Populations (6.6)).

Patients with Renal Impairment

The pharmacokinetics of abiraterone were examined in patients with end-stage renal disease (ESRD) on a stable hemodialysis schedule (N=8) and in matched control subjects with normal renal function (N=8). In the ESRD cohort of the trial, a single 1,000 mg abiraterone acetate dose was given under fasting conditions 1 hour after dialysis, and samples for pharmacokinetic analysis were collected up to 96 hours post-dose. Systemic exposure to abiraterone after a single oral 1,000 mg dose did not increase in subjects with end-stage renal disease on dialysis, compared to subjects with normal renal function (see Use in Specific Populations (6.7)).

Drug Interactions

In vitro studies with human hepatic microsomes showed that abiraterone has the potential to inhibit CYP1A2, CYP2D6, CYP2C8 and to a lesser extent CYP2C3, CYP2C19 and CYP3A4/5.

In an *in vivo* drug-drug interaction trial, the C_{max} and AUC of dextromethorphan (CYP2D6 substrate) were increased 2.8- and 2.5-fold, respectively when dextromethorphan 30 mg was given with abiraterone acetate 1,000 mg daily (plus prednisone 5 mg twice daily). The AUC for dextromethorphan, the active metabolite of dextromethorphan, increased approximately 1.3-fold (see Drug Interactions (7.2)).

In a clinical study to determine the effects of abiraterone acetate 1,000 mg daily (plus prednisone 5 mg twice daily) on a single 100 mg dose of the CYP1A2 substrate theophylline, no increase in systemic exposure of theophylline was observed.

Abiraterone is a substrate of CYP3A4, *in vitro*. In a clinical pharmacokinetic interaction study of healthy subjects pretreated with a strong CYP3A4 inducer (rifampin, 600 mg daily for 6 days) followed by a single dose of abiraterone acetate 1,000 mg, the mean plasma AUC $_{0-\infty}$ of abiraterone was decreased by 55% (see Drug Interactions (7.1)).

In a separate clinical pharmacokinetic interaction study of healthy subjects, co-administration of ketoconazole, a strong inhibitor of CYP3A4, had no clinically meaningful effect on the pharmacokinetics of abiraterone (see Drug Interactions (7.1)).

In a CYP2C3 drug-drug interaction trial in healthy subjects, the AUC of pioglitazone was increased by 46% when pioglitazone was given together with a single dose of 1,000 mg abiraterone acetate (see Drug Interactions (7.2)).

In vivo, abiraterone and its major metabolites were shown to inhibit the hepatic uptake transporter OATP1B1. There are no clinical data available to confirm transporter based interaction.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility

A two-year carcinogenicity study was conducted

in rats at oral abiraterone acetate doses of 5, 15, and 50 mg/kg/day

for males and 15, 50, and 150 mg/kg/day for females.

Abiraterone acetate increased the combined incidence of interstitial cell adenomas and carcinomas in the testes at all dose levels tested. This finding is considered to be related to the pharmacological activity of abiraterone. Rats are regarded as more sensitive than humans to developing interstitial cell tumors in the testes. Abiraterone acetate was not carcinogenic in female rats at exposure levels up to 0.8 times the human clinical exposure based on AUC. Abiraterone acetate was not carcinogenic in a 6-month study in the transgenic (T_{rasH2}) mouse.

Abiraterone acetate and abiraterone was not mutagenic in an *in vitro* cytogenetic assay using primary human lymphocytes or an *in vivo* rat micronucleus assay. In repeat-dose toxicity studies in male rats (13- and 26-week) and monkeys (9 weeks), atrophy, azoemia/hypoparathyria, and hyperplasia in the reproductive system were observed at 250 mg/kg/day in rats and 250 mg/kg/day in monkeys and were consistent with the antiandrogenic pharmacological activity of abiraterone. These effects were observed in rats at systemic exposures similar to humans, and in monkeys at exposures approximately 0.6 times the AUC in humans.

In a fertility study in male rats, reduced organ weights of the reproductive system, sperm counts, sperm motility, altered sperm morphology and decreased fertility were observed in animals dosed for 4 weeks at 20 mg/kg/day orally. Mating of untreated females with males that received 20 mg/kg/day oral abiraterone acetate resulted in a reduced number of corpora lutea, implantations and live embryos and an increased incidence of pre-implantation loss. Effects on male rats were reversible after 10 weeks from the last abiraterone acetate administration.

In a fertility study in female rats, animals dosed orally for 2 weeks until day 7 of pregnancy at 20 mg/kg/day had an increased incidence of irregular or extended estrous cycles and pre-implantation loss (300 mg/kg/day). There were no differences in mating, fertility, and litter parameters in female rats that received abiraterone acetate. Effects on female rats were reversible after 4 weeks from the last abiraterone acetate administration.

The dose of 30 mg/kg/day in rats is approximately 0.3 times the recommended dose of 1,000 mg/day based on body surface area.

In 13- and 26-week studies in rats and 13- and 39-week studies in monkeys, a reduction in circulating testosterone levels occurred with abiraterone acetate at approximately one half the human clinical exposure based on AUC. As a result, decreases in organ weights and toxicities were observed in the male and female reproductive system, adrenal glands, liver, pituitary (rats only), and male mammary glands. The changes in the reproductive organs are consistent with the antiandrogenic pharmacological activity of abiraterone acetate.

13.2 Animal Toxicology and/or Pharmacology

A dose-dependent increase in carcasses was observed in rats after daily oral abiraterone acetate administration for 26 weeks starting at 150 mg/kg/day (similar to the human clinical exposure based on AUC). In a 29-week monkey study with daily oral abiraterone acetate administration, no carcasses were observed at higher doses (2 times greater than the clinical exposure based on AUC).

14 CLINICAL STUDIES

The efficacy and safety of abiraterone acetate with prednisone was established in randomized placebo-controlled international clinical studies. All patients in these studies received a GnRH analog or had prior bilateral orchiectomy. Patients with prior leuteinizing hormone treatment for prostate cancer and a history of adrenal gland or pituitary disorders were excluded from these trials. Concurrent use of spironolactone was not allowed during the study period.

COU-AA-301: Patients with metastatic CRPC who had received prior docetaxel chemotherapy
 In COU-AA-301 (NCT00638690), a total of 1195 patients were randomized 2:1 to receive either abiraterone acetate orally at a dose of 1,000 mg once daily in combination with prednisone 5 mg orally twice daily (N=797) or placebo once daily plus prednisone 5 mg orally twice daily (N=398). Patients randomized to either arm were in continue treatment until disease progression (defined as a 25% increase in PSA over the patient's baseline) together with protocol-defined radiographic progression and symptomatic or clinical progression, initiation of new treatment, unacceptable toxicity or withdrawal.

The following patient demographics and baseline disease characteristics were balanced between the treatment arms. The median age was 69 years (range 39 to 95) and the racial distribution was 53% Caucasian, 33% Black, 1.7% Asian, and 1.6% Other. Eighty-nine percent of patients enrolled had an ECOG performance status score of 0 or 1 and 45% had a Brief Pain Inventory-Short Form score of ≥ 4 (patients reported worst pain over the previous 24 hours). Ninety percent of patients had metastases in bone and 30% had visceral involvement. Seventy percent of patients had radiographic evidence of disease progression and 30% had PSA-only progression. Seventy percent of patients had previously received one cytotoxic chemotherapy regimen and 30% received two regimens.

The protocol pre-specified interim analysis was conducted after 552 deaths and showed a statistically significant improvement in overall survival (OS) in patients treated with abiraterone acetate with prednisone compared to patients in the placebo with prednisone arm (Table 7 and Figure 5). An updated survival analysis was conducted when 775 deaths (67% of the planned number of deaths for final analysis) were observed. Results from this analysis were consistent with those from the interim analysis (Table 7).


Table 7: Overall Survival of Patients Treated with Either Abiraterone Acetate or Placebo in Combination with Prednisone in COU-AA-301 (Intent-to-Treat Analysis)

	Abiraterone Acetate with Prednisone (N=797)	Placebo with Prednisone (N=398)
Primary Survival Analysis		
Deaths (%)	53 (4.7%)	119 (5.5%)
Median survival (months) (95% CI)	18.8 (14.1, 15.4)	10.9 (10.2, 12.0)
p-value ¹	0.0001	
Hazard ratio (95% CI) ²	0.646 (0.543, 0.768)	
Updated Survival Analysis		
Deaths (%)	61 (6.3%)	174 (6.9%)
Median survival (months) (95% CI)	15.8 (14.8, 17.0)	11.2 (10.4, 13.1)
Hazard ratio (95% CI) ²	0.740 (0.638, 0.859)	

¹ p-value is derived from a log-rank test stratified by ECOG performance status score (0-1 vs. 2), pain score (absent vs. present), number of prior chemotherapy regimens (1 vs. 2), and type of disease progression (PSA only vs. radiographic).

² Hazard Ratio is derived from a stratified proportional hazards model. Hazard ratio <1 favors abiraterone acetate with prednisone.

Figure 1: Kaplan Meier Overall Survival Curves in COU-AA-301 (Intent-to-Treat Analysis)



COU-AA-382: Patients with metastatic CRPC who had not received prior cytotoxic chemotherapy
 In COU-AA-382 (NCT00807388), 1088 patients were randomized 1:1 to receive either abiraterone acetate orally at a dose of 1,000 mg once daily (N=546) or placebo orally once daily (N=542). Both arms were given concomitant prednisone 5 mg twice daily. Patients continued treatment until radiographic or clinical (cytotoxic chemotherapy, radiation or surgical treatment for cancer, pain requiring chronic opioids, or ECOG performance status decline to 3 or more) disease progression, unacceptable toxicity or withdrawal. Patients with moderate or severe pain, opiate use for cancer pain, or visceral organ metastases were excluded.

Patient demographics were balanced between the treatment arms. The median age was 70 years. The racial distribution of patients treated with abiraterone acetate was 50% Caucasian, 2.8% Black, 0.7% Asian and 1.1% Other. The ECOG performance status was 0 for 70% of patients, and 1 for 24% of patients. Co-primary efficacy endpoints were overall survival and radiographic progression free survival (RPFS). Baseline pain assessment was 0-1 (asymptomatic) in 66% of patients and 2 to 3 (mildly symptomatic) in 26% of patients as defined by the Brief Pain Inventory-Short Form (worst pain over the last 24 hours).

Radiographic progression-free survival was assessed with the use of sequential imaging studies and was defined by bone scan identification of 2 or more new bone lesions with confirmation (Prostate Cancer Working Group 2 criteria) and/or modified Response Evaluation Criteria in Solid Tumors (RECIST) criteria for progression of soft tissue lesions. Analysis of RPFS utilized centrally-reviewed radiographic assessment of progression.

The planned final analysis for OS, conducted after 741 deaths (median follow up of 49 months) demonstrated a statistically significant OS improvement in patients treated with abiraterone acetate with prednisone compared to those treated with placebo with prednisone (Table 8 and Figure 2). Sixty-five percent of patients on the abiraterone acetate arm and 78% of patients on the placebo arm used subsequent therapies that may prolong OS in metastatic CRPC. Abiraterone acetate was used as a subsequent therapy in 13% of patients on the abiraterone acetate arm and 44% of patients on the placebo arm.


Table 8: Overall Survival of Patients Treated with Either Abiraterone Acetate or Placebo in Combination with Prednisone in COU-AA-382 (Intent-to-Treat Analysis)

	Abiraterone Acetate with Prednisone (N=546)	Placebo with Prednisone (N=542)
Overall Survival		
Deaths (%)	35.4 (6.2%)	37 (7.1%)
Median survival (months) (95% CI)	34.7 (32.7, 36.8)	30.3 (28.7, 33.3)
p-value ¹	0.0033	
Hazard ratio ² (95% CI)	0.81 (0.70, 0.93)	

¹ p-value is derived from a log-rank test stratified by ECOG performance status score (0 vs. 1).

² Hazard Ratio is derived from a stratified proportional hazards model. Hazard ratio <1 favors abiraterone acetate with prednisone.

Figure 2: Kaplan Meier Overall Survival Curves in COU-AA-382



At the pre-specified RPFS analysis, 150 (28%) patients treated with abiraterone acetate with prednisone and 251 (46%) patients treated with placebo with prednisone had radiographic progression. A significant difference in RPFS between treatment groups was observed (Table 9 and Figure 3).

Table 9: Radiographic Progression-free Survival of Patients Treated with Either Abiraterone Acetate or Placebo in Combination with Prednisone in COU-AA-382 (Intent-to-Treat Analysis)


	Abiraterone Acetate with Prednisone (N=546)	Placebo with Prednisone (N=542)
Radiographic Progression-free Survival		
Progression-free survival (%)	50 (28%)	51 (46%)
Median RPFS (months) (95% CI)	35 (11.66, NR)	8.26 (8.12, 8.54)
p-value ¹	0.0001	
Hazard ratio ² (95% CI) ²	0.425 (0.347, 0.522)	

NR=Not reached.

¹ p-value is derived from a log-rank test stratified by ECOG performance status score (0 vs. 1).

² Hazard Ratio is derived from a stratified proportional hazards model. Hazard ratio <1 favors abiraterone acetate with prednisone.

Figure 3: Kaplan Meier Curves of Radiographic Progression-free Survival in COU-AA-382 (Intent-to-Treat Analysis)



The primary efficacy analyses are supported by the following prospectively defined endpoints. The median time to initiation of cytotoxic chemotherapy was 25.2 months for patients in the abiraterone acetate arm and 16.8 months for patients in the placebo arm (HR=0.580; 95% CI [0.487, 0.691], p < 0.0001).

The median time to opiate use for prostate cancer pain was not reached for patients receiving abiraterone acetate and was 23.7 months for patients receiving placebo (HR=0.686; 95% CI [0.566, 0.833], p < 0.0001). The time to opiate use result was supported by a delay in patient reported pain.

progression favoring the abiraterone acetate arm.

16 HOW SUPPLIED/STORAGE AND HANDLING

Abiraterone acetate 250 mg tablets, USP are white to off-white, oval, biconvex uncoated tablets. Engraved "AZ50" on one side, "APO" on the other side. Abiraterone acetate 250 mg tablets, USP are available in high-density polyethylene bottles of 120 and 360 tablets.

Cartons of 30 tablets (10 blister packs x 3), NDC 0904-6948-04

Storage and Handling

Store at 20°C to 25°C (68°F to 77°F); excursions permitted from 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Keep out of reach of children.

Based on its mechanism of action, abiraterone acetate may harm a developing fetus. Women who are pregnant or women who may be pregnant should not handle abiraterone acetate 250 mg uncoated tablets or other abiraterone acetate tablets if broken, crushed, or damaged without protection, e.g., gloves (see Use in Specific Populations (8.1)).

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Hypokalemia, Fluid Retention, and Cardiovascular Adverse Reactions

1. Inform patients that abiraterone acetate is associated with hypokalemia, hypohalemia, and peripheral edema that may lead to QT prolongation and Torsades de Pointes in patients who develop hypokalemia while taking abiraterone acetate. Advise patients that their blood pressure, serum potassium and signs and symptoms of fluid retention will be monitored clinically at least monthly. Advise patients to adhere to corticosteroids and to report symptoms of hypokalemia, hypohalemia, or edema to their healthcare provider (see Warnings and Precautions (5.3)).

Adrenocortical Insufficiency

1. Inform patients that abiraterone acetate with prednisone is associated with adrenal insufficiency. Advise patients to report symptoms of adrenocortical insufficiency to their healthcare provider (see Warnings and Precautions (5.2)).

Hepatotoxicity

1. Inform patients that abiraterone acetate is associated with severe hepatotoxicity. Inform patients that their liver function will be monitored using blood tests. Advise patients to immediately report symptoms of hepatotoxicity to their healthcare provider (see Warnings and Precautions (5.3)).

Use in Combination with Radium Ra 223 Dichloride

Advise patients that radium Ra 223 dichloride showed an increase in mortality and an increased rate of fracture when used in combination with abiraterone acetate plus prednisone/prednisolone. Inform patients to speak with their healthcare provider about any other medications or treatment they are currently taking for prostate cancer (see Warnings and Precautions (5.4)).

Dosing and Administration

1. Inform patients that abiraterone acetate tablets are taken once daily with prednisone (twice daily according to their healthcare provider's instructions) and to not interrupt or stop either of these medications without consulting their healthcare provider.
2. Inform patients receiving GnRH therapy that they need to maintain this treatment during the course of treatment with abiraterone acetate.
3. Instruct patients to take abiraterone acetate tablets on an empty stomach, at least one hour before or at least two hours after a meal. Abiraterone acetate tablets taken with food causes increased exposure and may result in adverse reaction. Instruct patients to swallow tablets whole with water and not to crush or chew the tablets (see Dosage and Administration (2.3)).
4. Inform patients that if they miss a dose of abiraterone acetate tablets or prednisone, they should take their normal dose the following day. If more than one daily dose is skipped, inform patients to contact their healthcare provider (see Dosage and Administration (2.3)).

Embryo-Fetal Toxicity

1. Inform patients that abiraterone acetate may harm a developing fetus and can cause loss of pregnancy.
2. Advise males with female partners of reproductive potential to use effective contraception during treatment and for 3 weeks after the final dose of abiraterone acetate tablets (see Use in Specific Populations (8.1)).
3. Advise females who are pregnant or women who may be pregnant not to handle abiraterone acetate 250 mg uncoated tablets or other abiraterone tablets if broken, crushed, or damaged without protection, e.g., gloves (see Use in Specific Populations (8.1) and How Supplied/Storage and Handling (16)).

Fertility

1. Advise male patients that abiraterone acetate may impair fertility (see Use in Specific Populations (8.3)).

APOTEX INC.

Abiraterone Acetate Tablets, USP
250 mg

Manufactured by: Apotex Inc.
Apotex Inc. Apotex Corp.
Toronto, Ontario Weston, Florida
Canada M9L 1T9 USA 33326

Distributed By:

MAJOR PHARMACEUTICALS
17177 N Laurel Park Dr., Suite 233
Livonia, MI 48152
Revised: July 2019
Rev: 9

PATIENT INFORMATION

Abiraterone Acetate Tablets, USP

(a' b' r a' t e r o n e a s e' t a b l e t s)

What are Abiraterone Acetate Tablets?

Abiraterone acetate tablets are a prescription medicine that is used along with prednisone. Abiraterone acetate tablets are used to treat men with prostate cancer that has spread to other parts of the body. Abiraterone acetate tablets are not for use in women.

It is not known if abiraterone acetate tablets are safe and effective in females or children.

Before you take Abiraterone Acetate Tablets, tell your healthcare provider about all of your medical conditions, including if you:

1. have heart problems
2. have liver problems
3. have a history of adrenal problems
4. have a history of pituitary problem
5. are receiving any other treatment for prostate cancer
6. are pregnant or plan to become pregnant. Abiraterone acetate tablets can cause harm to your unborn baby and loss of pregnancy (miscarriage). Females who are or may become pregnant should not handle abiraterone acetate tablets uncoated tablets or other Abiraterone acetate tablets if broken, crushed, or damaged without protection, such as gloves.
7. have a partner who is pregnant or may become pregnant.
8. Males who have female partners who are able to become pregnant should use effective birth control (contraception) during treatment with abiraterone acetate tablets and for 3 weeks after the last dose of abiraterone acetate tablets.
9. are breastfeeding or plan to breastfeed. It is not known if abiraterone acetate passes into your breast milk.

Tell your healthcare provider about all the medicines you take or treatment you receive, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Abiraterone acetate can interact with many other medicines.

You should not start or stop any medicine before you talk with the healthcare provider that prescribed abiraterone acetate tablets.

Know the medicines you take. Keep a list of them with you to show to your healthcare provider and pharmacist when you get a new medicine.

How should I take Abiraterone Acetate Tablets?

1. Take abiraterone acetate tablets and prednisone exactly as your healthcare provider tells you.
2. Take your prescribed dose of abiraterone acetate tablets 1 time a day.
3. Your healthcare provider may change your dose if needed.
4. Do not change or stop taking your prescribed dose of abiraterone acetate tablets or prednisone without talking with your healthcare provider first.
5. Take abiraterone acetate tablets on an empty stomach, at least one hour before or at least two hours after a meal. Do not take abiraterone acetate tablets with food. Taking abiraterone acetate tablets with food may cause most of the medicine to be absorbed by the body than is needed and this may cause side effects.
6. Swallow abiraterone acetate tablets whole. Do not crush or chew tablets.
7. Take abiraterone acetate tablets with water.
8. If you miss a dose of abiraterone acetate tablets or prednisone, take your prescribed dose the following day. If you miss more than 1 dose, tell your healthcare provider right away.
9. Your healthcare provider will do blood tests to check for side effects.

What are the possible side effects of Abiraterone Acetate Tablets?

Abiraterone Acetate Tablets may cause serious side effects including:

1. **High blood pressure (hypertension), low blood potassium levels (hypokalemia), fluid retention (edema), and irregular heartbeats can happen during treatment with abiraterone acetate tablets.** This can be life threatening. To decrease the chance of this happening, you must take prednisone with abiraterone acetate tablets exactly as your healthcare provider tells you. Your healthcare provider will check your blood pressure, do blood tests to check your potassium levels, and check for any signs and symptoms of fluid retention every month during treatment with abiraterone acetate tablets. Tell your healthcare provider if you get any of the following symptoms:

• dizziness	• confusion
• fast or irregular heartbeats	• muscle weakness
• feel faint or lightheaded	• pain in your legs
• headache	• swelling in your legs or feet

1. **Adrenal problems** may happen if you stop taking prednisone, get an infection, or are under stress.
2. **Liver problems.** You may develop changes in liver function/blood test. Your healthcare provider will do blood tests to check your liver before treatment with abiraterone acetate tablets and during treatment with abiraterone acetate tablets. Liver failure may occur, which can lead to death. Tell your healthcare provider if you notice any of the following changes:
 1. yellowing of the skin or eyes
 2. darkening of the urine
 3. severe nausea or vomiting

Increased risk of bone fracture and death when abiraterone acetate tablets and prednisone or prednisolone is used in combination with a type of radiation called radium Ra 223 dichloride. Tell your healthcare provider about any other treatments you are taking for prostate cancer.

The most common side effects of abiraterone acetate include:

• feeling very tired	• vomiting
• joint pain	• infected nose, sinuses, or throat (cold)
• high blood pressure	• cough
• nausea	• headache
• •	• low red blood cells (anemia)
swelling in your legs or feet	• high blood cholesterol and levels
• low blood potassium	• triglycerides
• hot flashes	• high blood sugar levels
• diarrhea	• •
	certain other abnormal blood tests

Abiraterone acetate tablets may cause fertility problems in males, which may affect the ability to father children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of abiraterone acetate.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Abiraterone Acetate Tablets?

1. Store abiraterone acetate tablets at room temperature between 68°F to 77°F (20°C to 25°C).

Keep Abiraterone Acetate Tablets and all medicines out of the reach of children.

General information about Abiraterone Acetate Tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use abiraterone acetate tablets for a condition for which it was not prescribed. Do not give abiraterone acetate tablets to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your healthcare provider or pharmacist for information about abiraterone acetate tablets that is written for health professionals.

For more information contact Apotex Corp. at 1-800-706-5575.

What are the ingredients of Abiraterone Acetate Tablets?

Active ingredient: abiraterone acetate

Inactive ingredients: colloidal silicon dioxide, croscopollose, lactose monohydrate, magnesium stearate, and sodium lauryl sulfate.

This Patient Information has been approved by the U.S. Food and Drug Administration.

APOTEX INC.

Abiraterone Acetate Tablets, USP

250 mg

Manufactured by: Apotex Inc. Manufactured for: Apotex Corp.

Apotex Inc. Apotex Corp.
Toronto, Ontario Weston, Florida
Canada M9L 1T8 USA 33326

Distributed by:

MAJOR PHARMACEUTICALS

17177 N Laurel Park Dr., Suite 233

Livonia, MI 48152

Revised July 2019

Rev: 9

Package Label Display Panel

Abiraterone Acetate Tablets, USP

250 mg

30 Tablets



ABIRATERONE ACETATE				
abiraterone acetate tablets				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC 6904-6948-04(NDC 6905-4127)	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Abiraterone Acetate (UNII: EN6G3V7W1) (Abiraterone - UNII: 8R4652D)	Abiraterone Acetate	250 mg		
Inactive Ingredients				
	Ingredient Name	Strength		
	SILICON DIOXIDE (UNII: 81T7888M2)			
	CROSCOPOLLONE (E141MPS AT 2%) (UNII: 6840 0643M2)			
	Lactose Monohydrate (UNII: EW02720E5)			
	Magnesium Stearate (UNII: 761306C33)			
	Sodium Lauryl Sulfate (UNII: 368G6544D)			
Product Characteristics				
Color	White to off-white	Shape	Round	
Shape	Oval (Biconvex)	Size	6mm	
Flavor		Imprint Code	A25GAP0	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC 6904-6948-04	30 in 1 CARTON	10/23/08	
1		[1 in 1 BUSTER PACK; Type 0; Not a Combination Product]		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA209453	10/23/08		
Labeler - Major Pharmaceuticals (01942727)				
Registrant - Apotex Inc. (20942912)				

Revised: 7/20/20

Major Pharmaceuticals