


GLYCOPYRROLATE- glycopyrrolate tablet
Leading Pharma, LLC

GLYCOPYRROLATE TABLETS, USP
1 mg and 2 mg
Rx only

DESCRIPTION

Glycopyrrolate Tablet, USP contain the synthetic anticholinergic, glycopyrrolate. Glycopyrrolate is a quaternary ammonium compound with the following chemical name: 3-[(cyclopentylhydroxyphenylacetyl)oxy]-1, 1-dimethylpyrrolidinium bromide. Its empirical formula is $C_{19}H_{28}BrNO_3$, its molecular weight is 398.33, and its structural formula is:



Each 1 mg tablet contains: Glycopyrrolate USP, 1 mg

Each 2 mg tablet contains: Glycopyrrolate USP, 2 mg

Inactive ingredients: dibasic calcium phosphate, lactose, magnesium stearate, povidone, and sodium starch glycolate

ACTIONS

Glycopyrrolate, like other anticholinergic (antimuscarinic) agents, inhibits the action of acetylcholine on structures innervated by postganglionic cholinergic nerves and on smooth muscles that respond to acetylcholine but lack cholinergic innervation. These peripheral cholinergic receptors are present in the autonomic effector cells of smooth muscle, cardiac muscle, the sino-atrial node, the atrioventricular node, exocrine glands, and, to a limited degree, in the autonomic ganglia. Thus, it diminishes the volume and free acidity of gastric secretions and controls excessive pharyngeal, tracheal, and bronchial secretions.

Glycopyrrolate antagonizes muscarinic symptoms (e.g., bronchorrhea, bronchospasm, bradycardia, and intestinal hypermotility) induced by cholinergic drugs such as the anticholinesterases.

The highly polar quaternary ammonium group of glycopyrrolate limits its passage across lipid membranes, such as the blood-brain barrier, in contrast to atropine sulfate and scopolamine hydrobromide, which are non-polar tertiary amines which penetrate lipid barriers easily.

INDICATIONS

For use as adjunctive therapy in the treatment of peptic ulcer.

CONTRAINDICATIONS

Glaucoma; obstructive uropathy (for example, bladder neck obstruction due to prostatic hypertrophy);

obstructive disease of the gastrointestinal tract (as in achalasia, pyloroduodenal stenosis, etc.); paralytic ileus; intestinal atony of the elderly or debilitated patient; unstable cardiovascular status in acute hemorrhage; severe ulcerative colitis; toxic megacolon complicating ulcerative colitis; myasthenia gravis. Glycopyrrolate Tablet, USP are contraindicated in those patients with a *hypersensitivity to glycopyrrolate*.

WARNINGS

In the presence of a high environmental temperature, heat prostration (fever and heat stroke due to decreased sweating) can occur with use of Glycopyrrolate Tablet, USP.

Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy or colostomy. In this instance treatment with this drug would be inappropriate and possibly harmful.

Glycopyrrolate Tablet, USP may produce drowsiness or blurred vision. In this event, the patient should be warned not to engage in activities requiring mental alertness such as operating a motor vehicle or other machinery, or performing hazardous work while taking this drug.

Theoretically, with overdosage, a curare-like action may occur, i.e., neuro-muscular blockade leading to muscular weakness and possible paralysis.

Pregnancy

The safety of this drug during pregnancy has not been established. The use of any drug during pregnancy requires that the potential benefits of the drug be weighed against possible hazards to mother and child. Reproduction studies in rats revealed no teratogenic effects from glycopyrrolate; however, the potent anticholinergic action of this agent resulted in diminished rates of conception and of survival at weaning, in a dose-related manner. Other studies in dogs suggest that this may be due to diminished seminal secretion which is evident at high doses of glycopyrrolate. Information on possible adverse effects in the pregnant female is limited to uncontrolled data derived from marketing experience. Such experience has revealed no reports of teratogenic or other fetus-damaging potential. No controlled studies to establish the safety of the drug in pregnancy have been performed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. As a general rule, nursing should not be undertaken while a patient is on a drug since many drugs are excreted in human milk.

Pediatric Use

Since there is no adequate experience in pediatric patients who have received this drug, safety and efficacy in pediatric patients have not been established.

PRECAUTIONS

Use Glycopyrrolate Tablet, USP with caution in the elderly and in all patients with:

- Autonomic neuropathy.
- Hepatic or renal disease.
- Ulcerative colitis—large doses may suppress intestinal motility to the point of producing a paralytic ileus and for this reason may precipitate or aggravate “toxic megacolon,” a serious complication of the disease.
- Hyperthyroidism, coronary heart disease, congestive heart failure, cardiac tachyarrhythmias, tachycardia, hypertension and prostatic hypertrophy.

- Hiatal hernia associated with reflux esophagitis, since anticholinergic drugs may aggravate this condition.

ADVERSE REACTIONS

Anticholinergics produce certain effects, most of which are extensions of their fundamental pharmacological actions. Adverse reactions to anticholinergics in general may include xerostomia; decreased sweating; urinary hesitancy and retention; blurred vision; tachycardia; palpitations; dilatation of the pupil; cycloplegia; increased ocular tension; loss of taste; headaches; nervousness; mental confusion; drowsiness; weakness; dizziness; insomnia; nausea; vomiting; constipation; bloated feeling; impotence; suppression of lactation; severe allergic reaction or drug idiosyncrasies including anaphylaxis, urticaria and other dermal manifestations.

Glycopyrrolate is chemically a quaternary ammonium compound; hence, its passage across lipid membranes, such as the blood-brain barrier, is limited in contrast to atropine sulfate and scopolamine hydrobromide. For this reason the occurrence of CNS related side effects is lower, in comparison to their incidence following administration of anticholinergics which are chemically tertiary amines that can cross this barrier readily.

CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT SIDE EFFECTS. YOU MAY REPORT SIDE EFFECTS TO THE FDA AT 1-800-FDA-1088 OR LEADING PHARMA, LLC AT 1-844-740-7500.

OVERDOSAGE

The symptoms of overdosage of Glycopyrrolate Tablet, USP are peripheral in nature rather than central.

1. To guard against further absorption of the drug—use gastric lavage, cathartics and/or enemas.
2. To combat peripheral anticholinergic effects (residual mydriasis, dry mouth, etc.)—utilize a quaternary ammonium anticholinesterase, such as neostigmine methylsulfate.
3. To combat hypotension—use pressor amines (norepinephrine, metaraminol) i.v.; and supportive care.
4. To combat respiratory depression—administer oxygen; utilize a respiratory stimulant such as Dopram[®] i.v.; artificial respiration.

DOSAGE AND ADMINISTRATION

The dosage of Glycopyrrolate Tablet, USP should be adjusted to the needs of the individual patient to assure symptomatic control with a minimum of adverse reactions. The presently recommended maximum daily dosage of glycopyrrolate is 8 mg.

Glycopyrrolate Tablet, USP 1 mg. The recommended initial dosage of Glycopyrrolate Tablet, USP for adults is one tablet three times daily (in the morning, early afternoon, and at bedtime). Some patients may require two tablets at bedtime to assure overnight control of symptoms. For maintenance, a dosage of one tablet twice a day is frequently adequate.

Glycopyrrolate Tablet, USP 2 mg. The recommended dosage of Glycopyrrolate Tablet, USP for adults is one tablet two or three times daily at equally spaced intervals.

Glycopyrrolate Tablet, USP are not recommended for use in pediatric patients under the age of 12 years.

DRUG INTERACTIONS

There are no known drug interactions.

HOW SUPPLIED

Glycopyrrolate Tablet, USP 1 mg are bisected, compressed white, round tablets debossed “EP” above the bisect and “139” below the bisect on one side of the tablet, and plain on the other side.
Available in bottles of 100 (NDC 69315-139-01)

Glycopyrrolate Tablet, USP 2 mg are bisected, compressed white, round tablets debossed “EP” above the bisect and “140” below the bisect on one side of the tablet, and plain on the other side.
Available in bottles of 100 (NDC 69315-140-01).

Store at Controlled Room Temperature 20° - 25°C (68° - 77°F).

Dispense in a tight, light-resistant container as defined in the USP, using a child-resistant closure.

Rx only

Manufactured by:
Leading Pharma, LLC
Fairfield, NJ 07004
Rev. 03 11/17

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 69315-139-01

1 mg: Bottle of 100 Tablets

NDC 69315-139-01

**GLYCOPYRROLATE
TABLETS, USP**

1 mg

Rx only
100 Tablets

**LEADING
PHARMA**

Each Tablet Contains:
Glycopyrrolate, USP..... 1 mg

Usual Dosage: One or two tablets three times a day. See accompanying information.

Dispense in a tight, light-resistant container as defined in the USP, using a child-resistant closure.

Keep this and all Medications out of the reach of children.

Store at 20°-25°C (68°-77°F) [see USP controlled room temperature].

Manufactured by:
Leading Pharma, LLC
Fairfield, NJ 07004

Tablet Imprinted: EP 139 Rev. 01

N 3 69315-139-01 1

.6" Unvarnished Area

NDC 69315-140-01

2 mg: Bottle of 100 Tablets

NDC 69315-140-01

GLYCOPYRROLATE TABLETS, USP

2 mg

Rx only
100 Tablets

LEADING
PHARMA

Each Tablet Contains:

Glycopyrrolate, USP 2 mg

Usual Dosage: One tablet two or three times a day. See accompanying information.

Dispense in a tight, light-resistant container as defined in the USP, using a child-resistant closure.

Keep this and all Medications out of the reach of children.

Store at 20°-25°C (68°-77°F) [see USP controlled room temperature].

Manufactured by:
Leading Pharma, LLC
Fairfield, NJ 07004

Tablet Imprinted: EP 140

Rev. 01



N 3 69315-140-01 7

.6" Unvarnished Area

GLYCOPYRROLATE

glycopyrrolate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:69315-139
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GLYCOPYRROLATE (UNII: V92SO9WP2I) (GLYCOPYRROLONIUM - UNII:A14FB57V1D)	GLYCOPYRROLATE	1 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	8mm
Flavor		Imprint Code	EP;139
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69315-139-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/22/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090195	09/22/2018	

GLYCOPYRROLATE

glycopyrrolate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:69315-140
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GLYCOPYRROLATE (UNII: V92SO9WP2I) (GLYCOPYRRONIUM - UNII:A14FB57V1D)	GLYCOPYRROLATE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	EP;140
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69315-140-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/22/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090195	09/22/2018	

Labeler - Leading Pharma, LLC (079575060)

Establishment

Name	Address	ID/FEI	Business Operations
Leading Pharma, LLC		079575060	MANUFACTURE(69315-139, 69315-140)

Revised: 10/2019

Leading Pharma, LLC