

ASPIRIN- aspirin tablet, film coated
Discount Drug Mart

Drug Mart 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor

- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal display panel

DISCOUNT

drug

mart

FOOD MARKET

AN EMPLOYEE OWNERSHIP COMPANY

ASPIRIN

(NSAID)

pain reliever/fever reducer

Compare to the active ingredient of **GENUINE BAYER®** Aspirin[†]

100 TABLETS - 325 mg each

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

DISTRIBUTED BY:

DRUG MART-FOOD FAIR, 211 COMMERCE DRIVE, MEDINA, OH 44256

www.discount-drugmart.com

[†]This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin. 50844 REV0122C15712



ASPIRIN (NSAID)

pain reliever/fever reducer

Compare to the active ingredient in GENUINE BAYER® Aspirin®

100 TABLETS - 325 mg each



0 9335110160 1

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin. 50844 REV10122C15712
DISTRIBUTED BY:
DRUG MART-FOOD FAIR, 211 COMMERCE DRIVE, MEDINA, OH 44256
www.discount-drugmart.com

TAMPER EVIDENT, DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient (in each tablet)
Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

no print / no varnish area
lot no. & exp. date

Uses temporarily relieves:
 ■ minor pain of arthritis
 ■ headache ■ muscle pain ■ menstrual pain
 ■ toothache ■ pain and fever of colds

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ facial swelling
 ■ shock ■ hives ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ are age 60 or older
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ have had stomach ulcers or bleeding problems
 ■ take more or for a longer time than directed
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma
Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis
Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools
 ■ feel faint ■ have stomach pain that does not get better
 ■ ringing in the ears or a loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ dizziness or swelling is present ■ new symptoms occur
 These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

PEEL HERE FOR MORE DRUG FACTS

STOP PEELING

Questions or comments? 1-800-426-9391

Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol
Other information ■ use by expiration date on package ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Directions ■ do not take more than directed
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: do not use unless directed by a doctor

Drug Facts (continued)
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma
Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis
Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools
 ■ feel faint ■ have stomach pain that does not get better
 ■ ringing in the ears or a loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ dizziness or swelling is present ■ new symptoms occur
 These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Mart 44-157

ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-157
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-157-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2019	
2	NDC:53943-157-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2019	
3	NDC:53943-157-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/22/2019	

Labeler - Discount Drug Mart (047741335)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(53943-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(53943-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(53943-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(53943-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(53943-157)

Revised: 2/2024

Discount Drug Mart