

**MENSTRUAL COMPLETE MAXIMUM STRENGTH- acetaminophen, caffeine, pyrilamine maleate tablet, film coated**

**Meijer Distribution Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Meijer 44-390**

***Active ingredients (in each caplet)***

Acetaminophen 500 mg

Caffeine 60 mg

Pyrilamine maleate 15 mg

***Purpose***

Pain reliever

Diuretic

Antihistamine

***Uses***

for the temporary relief of these symptoms associated with menstrual periods:

- bloating
- headache
- water-weight gain
- cramps
- backache
- fatigue
- muscle aches

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- avoid alcoholic beverages
- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness
- you may get drowsy
- limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not take more than the recommended dose**
- adults and children 12 years and over:
  - take 2 caplets with water
  - repeat every 6 hours, as needed
  - do not exceed 6 caplets per day
- children under 12 years: ask a doctor

***Other information***

- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

***Inactive ingredients***

corn starch, croscarmellose sodium, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

**Questions or comments?**

**1-800-426-9391**

**Principal Display Panel**

NDC 41250-390-10

**Compare to Midol® Complete** active ingredients\*

**meijer®**

Maximum Strength

**menstrual  
complete**

**Acetaminophen I Caffeine I Pyrilamine maleate**  
Pain Reliever I Diuretic I Antihistamine

Multi-Symptom Relief of:  
Cramps, Bloating, Fatigue,  
Backache & Headache

**40 Caplets**                      actual size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING**

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Midol® Complete.

50844    REV0517B39010

**DIST. BY: MEIJER  
DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com**

**Drug Facts (continued)**

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than the recommended dose
- adults and children 12 years and over:
  - take 2 caplets with water
  - repeat every 6 hours, as needed
  - do not exceed 6 caplets per day

**Drug Facts (continued)**

children under 12 years: ask a doctor

**Other Information**

- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients**

croscarmellose sodium, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, Traceln

**Drug Facts (continued)**

If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- avoid alcoholic beverages
- excitability may occur, especially in children

**Drug Facts (continued)**

operating machinery

- alcohol, sedatives, and tranquilizers may increase drowsiness
- you may get drowsy
- limit the use of caffeine-containing medications, foods, or beverages while taking this product
- because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Midol® Complete. 50844 REV0517B39010

**OUR QUALITY GUARANTEE**

*The Meijer Family*

www.meijer.com

**DIST. BY MEIJER DISTRIBUTION, INC.**  
GRAND RAPIDS, MI 49544  
www.meijer.com

**Drug Facts (continued)**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

more than 4,000 mg of acetaminophen in 24 hours

with other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

**Warnings**

**Drug Facts (continued)**

**Active ingredients (in each caplet)**

Acetaminophen 500 mg. Pain reliever

Caffeine 60 mg. Diuretic

Pyrilamine maleate 15 mg. Antihistamine

**Uses**

for the temporary relief of these symptoms associated with menstrual periods:

- water-weight gain
- cramps
- backache
- fatigue
- bloating
- headache
- muscle aches

**Purpose**

**Drug Facts (continued)**

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

more than 4,000 mg of acetaminophen in 24 hours

with other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

**Drug Facts (continued)**

(Questions or comments? 1-800-426-9391)



no print/no varnish area  
lot no. & exp.date

B-1214-390-10-R  
REV0517B39010

NDC 41250-390-10

Compare to Midol® Complete active ingredients\*

**meijer**

**MAXIMUM STRENGTH**

**menstrual complete**

Acetaminophen | Caffeine | Pyrilamine maleate

Pain Reliever | Diuretic | Antihistamine

Multi-Symptom Relief of:  
Cramps, Bloating, Fatigue,  
Backache & Headache

**40 Caplets**

actual size

Meijer 44-390

**MENSTRUAL COMPLETE MAXIMUM STRENGTH**  
acetaminophen, caffeine, pyrilamine maleate tablet, film coated

**Product Information**

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:41250-390 |
| <b>Route of Administration</b> | ORAL           |                           |               |

**Active Ingredient/Active Moiety**

| <b>Ingredient Name</b>   | <b>Basis of Strength</b> | <b>Strength</b> |
|--|--------------------------|-----------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)   | ACETAMINOPHEN            | 500 mg          |
| CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)             | CAFFEINE                 | 60 mg           |
| PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL) | PYRILAMINE MALEATE       | 15 mg           |

**Inactive Ingredients**

| <b>Ingredient Name</b>                              | <b>Strength</b> |
|---|-----------------|
| STARCH, CORN (UNII: O8232NY3SJ)                     |                 |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48)             |                 |
| HYPROMELLOSES (UNII: 3NXW29V3WO)                    |                 |
| MAGNESIUM STEARATE (UNII: 70097M6I30)               |                 |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)      |                 |
| POLYDEXTROSE (UNII: VH2XOU12IE)                     |                 |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) |                 |
| POVIDONE (UNII: FZ989GH94E)                         |                 |
| STEARIC ACID (UNII: 4ELV7Z65AP)                     |                 |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                 |                 |
| TRIACETIN (UNII: XHX3C3X673)                        |                 |
| CROSPVIDONE (UNII: 2S7830E561)                      |                 |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                  |                 |

**Product Characteristics**

|                 |       |                     |          |
|-----------------|-------|---------------------|----------|
| <b>Color</b>    | WHITE | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL  | <b>Size</b>         | 17mm     |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | 44;390   |
| <b>Contains</b> |       |                     |          |

**Packaging**

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:41250-390-10 | 1 in 1 CARTON                                     | 04/29/2002           |                    |
| 1 |                  | 40 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |

**Marketing Information**

| <b>Marketing Category</b> | <b>Application Number or Monograph Citation</b> | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|---------------------------|---|-----------------------------|---------------------------|
| OTC MONOGRAPH NOT FINAL   | part343   | 04/29/2002                  |                           |

**Labeler** - Meijer Distribution Inc (006959555)**Establishment**

| <b>Name</b>             | <b>Address</b> | <b>ID/FEI</b> | <b>Business Operations</b> |
|-------------------------|----------------|---------------|----------------------------|
| LNK International, Inc. |                | 038154464     | PACK(41250-390)            |

**Establishment**

| <b>Name</b>             | <b>Address</b> | <b>ID/FEI</b> | <b>Business Operations</b> |
|-------------------------|----------------|---------------|----------------------------|
| LNK International, Inc. |                | 832867894     | MANUFACTURE(41250-390)     |

**Establishment**

| <b>Name</b>             | <b>Address</b> | <b>ID/FEI</b> | <b>Business Operations</b> |
|-------------------------|----------------|---------------|----------------------------|
| LNK International, Inc. |                | 832867837     | PACK(41250-390)            |

**Establishment**

| <b>Name</b>             | <b>Address</b> | <b>ID/FEI</b> | <b>Business Operations</b> |
|-------------------------|----------------|---------------|----------------------------|
| LNK International, Inc. |                | 967626305     | PACK(41250-390)            |

**Establishment**

| <b>Name</b>             | <b>Address</b> | <b>ID/FEI</b> | <b>Business Operations</b> |
|-------------------------|----------------|---------------|----------------------------|
| LNK International, Inc. |                | 868734088     | PACK(41250-390)            |

Revised: 9/2020

Meijer Distribution Inc