

MIDOL LONG LASTING RELIEF- acetaminophen tablet, extended release
Bayer HealthCare LLC.

Midol Long Lasting Relief

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

Uses

- Temporarily relieves minor aches and pains due
-
- | | | |
|------------------|-------------------------------------|---------------------------|
| • muscular aches | • backache | • minor pain of arthritis |
| • toothache | • premenstrual and menstrual cramps | |
| • headache | • the common cold | |
-
- temporarily reduces fever

Warnings

Liver warning: The product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

-
- | | | |
|------------------|------------|--------|
| • skin reddening | • blisters | • rash |
|------------------|------------|--------|
-

- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are not sure whether a drug contains acetaminophen, ask a doctor

or pharmacist.

- If you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease

As a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 8 hours with water• swallow whole; do not crush, chew, split or dissolve• do not take more than 6 caplets in 24 hours<ul style="list-style-type: none">◦ do not use for more than 10 days unless directed by a doctor
children under 12 years	<ul style="list-style-type: none">• do not use

Other information

Other information

- store at 20-25° (68-77°F). Avoid excessive heat 40°C (104°F).
- **do not use if printed foil under cap is broken or missing**

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments

Questions or comments? 1-800-331-4536

MIDOL LONG LASTING RELIEF

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	650 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CARNAUBA WAX (UNII: R12CBM0EIZ)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
MALTODEXTRIN (UNII: 7CVR7L4A2D)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
POVIDONE (UNII: FZ989GH94E)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	white	Score	no score	
Shape	OVAL	Size	18mm	
Flavor		Imprint Code	MLLR	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-8090-20	1 in 1 CARTON	03/10/2017	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA075077	03/10/2017	

Labeler - Bayer HealthCare LLC. (112117283)