


**LUXIQ- betamethasone valerate aerosol, foam**  
**Mylan Pharmaceuticals Inc.**

-----  
**DESCRIPTION**

Luxiq Foam contains betamethasone valerate, a synthetic corticosteroid, for topical dermatologic use. The corticosteroids constitute a class of primarily synthetic steroids used topically as anti-inflammatory agents.

Betamethasone valerate is 9-fluoro-11 $\beta$ ,17, 21-trihydroxy-16 $\beta$ -methylpregna-1, 4-diene-3, 20-dione 17-valerate, with the empirical formula  $C_{27}H_{37}FO_6$ , a molecular weight of 476.58. The following is the chemical structure:



**Betamethasone valerate**

Betamethasone valerate, USP is a white to practically white, odorless crystalline powder, and is practically insoluble in water, freely soluble in acetone and in chloroform, soluble in alcohol, and slightly soluble in benzene and in ether.

Luxiq<sup>®</sup> (betamethasone valerate) Foam, 0.12%, contains 1.2 mg betamethasone valerate, USP, per gram in a thermolabile hydroethanolic foam vehicle consisting of cetyl alcohol, citric acid, ethanol (60.4%), polysorbate 60, potassium citrate, propylene glycol, purified water, and stearyl alcohol pressurized with a hydrocarbon (propane/butane) propellant.

**CLINICAL PHARMACOLOGY**

Like other topical corticosteroids, betamethasone valerate foam has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A<sub>2</sub> inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A<sub>2</sub>.

## Pharmacokinetics

Topical corticosteroids can be absorbed from intact healthy skin. The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle and the integrity of the epidermal barrier. Occlusion, inflammation and/or other disease processes in the skin may also increase percutaneous absorption.

The use of pharmacodynamic endpoints for assessing the systemic exposure of topical corticosteroids is necessary due to the fact that circulating levels are well below the level of detection. Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. They are metabolized, primarily in the liver, and are then excreted by the kidneys. In addition, some corticosteroids and their metabolites are also excreted in the bile.

## CLINICAL STUDIES

The safety and efficacy of Luxiq has been demonstrated in a four-week trial. An adequate and well-controlled clinical trial was conducted in 190 patients with moderate to severe scalp psoriasis. Patients were treated twice daily for four weeks with Luxiq Foam, Placebo foam, a commercially available betamethasone valerate lotion 0.12% (formerly expressed as 0.1% betamethasone), or Placebo lotion. At four weeks of treatment, study results of 159 patients demonstrated that the efficacy of Luxiq Foam in treating scalp psoriasis is superior to that of Placebo foam, and is comparable to that of a currently marketed BMV lotion (see Table below).

Subjects with Target Lesion Parameter Clear at Endpoint	Luxiq Foam n (%)	BMV lotion n (%)	Placebo foam n (%)
Scaling	30 (47%)	22 (35%)	2 (6%)
Erythema	26 (41%)	16 (25%)	2 (6%)
Plaque Thickness	42 (66%)	25 (40%)	5 (16%)
Investigator's Global: Subjects Completely Clear or Almost Clear at Endpoint	43 (67%)	29 (46%)	6 (19%)

## INDICATIONS AND USAGE

Luxiq is a medium potency topical corticosteroid indicated for relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses of the scalp.

## CONTRAINDICATIONS

Luxiq is contraindicated in patients who are hypersensitive to betamethasone valerate, to other corticosteroids, or to any ingredient in this preparation.

## PRECAUTIONS

### General

Systemic absorption of topical corticosteroids has caused reversible hypothalamic-pituitary-adrenal (HPA) axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal of treatment. Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on treatment.

Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings.

Therefore, patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid.

Recovery of HPA axis function is generally prompt upon discontinuation of topical corticosteroids. Infrequently, signs and symptoms of glucocorticosteroid insufficiency may occur requiring supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Pediatric patients may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios. (See PRECAUTIONS: Pediatric Use.)

If irritation develops, Luxiq should be discontinued and appropriate therapy instituted. Allergic contact dermatitis with corticosteroids is usually diagnosed by observing a failure to heal rather than noting a clinical exacerbation, as with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic patch testing.

In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, use of Luxiq should be discontinued until the infection has been adequately controlled.

### **Information for Patients**

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. The treated scalp area should not be bandaged or otherwise covered or wrapped so as to be occlusive unless directed by the physician.
4. Patients should report to their physician any signs of local adverse reactions.
5. As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, contact the physician.

### **Laboratory Tests**

The following tests may be helpful in evaluating patients for HPA axis suppression:

ACTH stimulation test

A.M. plasma cortisol test

Urinary free cortisol test

### **Carcinogenesis, Mutagenesis, and Impairment of Fertility**

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of betamethasone valerate.

Betamethasone was genotoxic in the *in vitro* human peripheral blood lymphocyte chromosome aberration assay with metabolic activation and in the *in vivo* mouse bone marrow micronucleus assay.

### **Pregnancy**

Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women. Therefore, Luxiq should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

### **Nursing Mothers**

Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Because many drugs are excreted in human milk, caution should be exercised when Luxiq is administered to a nursing woman.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established. Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA axis suppression and Cushing's syndrome when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency during and/or after withdrawal of treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children.

Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome, linear growth retardation, delayed weight gain, and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include low plasma cortisol levels and an absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to children should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

### **ADVERSE REACTIONS**

The most frequent adverse event was burning/itching/stinging at the application site; the incidence and severity of this event were as follows:

Incidence and severity of burning/itching/stinging				
Product	Total incidence	Maximum severity		
		Mild	Moderate	Severe
Luxiq Foam n = 63	34 (54%)	28 (44%)	5 (8%)	1 (2%)
Betamethasone valerate lotion n = 63	33 (52%)	26 (41%)	6 (10%)	1 (2%)
Placebo Foam n = 32	24 (75%)	13 (41%)	7 (22%)	4 (12%)
Placebo Lotion n = 30	20 (67%)	12 (40%)	5 (17%)	3 (10%)

Other adverse events which were considered to be possibly, probably, or definitely related to Luxiq occurred in 1 patient each; these were paresthesia, pruritus, acne, alopecia, and conjunctivitis.

The following additional local adverse reactions have been reported with topical corticosteroids, and they may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximately decreasing order of occurrence: irritation; dryness; folliculitis; acneiform eruptions; hypopigmentation; perioral dermatitis; allergic contact dermatitis; secondary infection; skin atrophy; striae; and miliaria.

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients.

## OVERDOSAGE

Topically applied Luxiq can be absorbed in sufficient amounts to produce systemic effects. (See PRECAUTIONS.)

## DOSAGE AND ADMINISTRATION

Note: For proper dispensing of foam, can must be inverted.

For application to the scalp invert can and dispense a small amount of Luxiq onto a saucer or other cool surface. Do not dispense directly onto hands as foam will begin to melt immediately upon contact with warm skin. Pick up small amounts of foam with fingers and gently massage into affected area until foam disappears. Repeat until entire affected scalp area is treated. Apply twice daily, once in the morning and once at night.

As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, reassessment of the diagnosis may be necessary.

Luxiq should not be used with occlusive dressings unless directed by a physician.

## HOW SUPPLIED

Luxiq (betamethasone valerate) Foam, 0.12% contains 1.2 mg of betamethasone valerate, USP per gram. It is available as follows:

NDC 0378-8187-50

carton containing one 50 g aluminum can

NDC 0378-8187-01

carton containing one 100 g aluminum can

**Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]**

## WARNING

**FLAMMABLE. AVOID FIRE, FLAME OR SMOKING DURING AND IMMEDIATELY FOLLOWING APPLICATION.** Keep this and all medication out of the reach of children. Contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120°F (49°C).

## PATIENT INFORMATION

### **Luxiq<sup>®</sup>**

(betamethasone valerate) Foam, 0.12%

### **About Luxiq**

Your doctor has prescribed Luxiq (betamethasone valerate) Foam, 0.12%, for the relief of corticosteroid-responsive skin conditions of the scalp. Luxiq works because its active ingredient is betamethasone valerate, 0.12%. Betamethasone belongs to a group of medicines known as topical corticosteroids. These agents are used to reduce the inflammation, redness, swelling, itching, and tenderness associated with dermatologic conditions.

Other ingredients in Luxiq include cetyl alcohol, citric acid, ethanol, polysorbate 60, potassium citrate, propylene glycol, purified water, and stearyl alcohol. The foam is dispensed from an aluminum can that is pressurized by a hydrocarbon propellant (propane and butane).

**If you answer YES to one or more of the following questions, tell your doctor (or pharmacist) before using this medicine, so you can get advice about what to do.**

- Are you allergic to any of the ingredients contained in Luxiq?
- Are you pregnant? Planning on becoming pregnant while using Luxiq? Or are you breastfeeding?
- Do you think you have an infection on your scalp?

### **How to apply Luxiq**



*Turn the can upside down and dispense a small amount of Luxiq onto a clean saucer or other cool, clean surface. Do not dispense directly onto hands, as foam will begin to melt immediately upon contact with warm skin.*

*Pick up small amounts of foam with fingers and gently massage into affected area until foam disappears. Repeat until entire affected scalp area is treated. Apply twice daily, once in the morning and once at night. Use sparingly—only enough to cover the affected areas.*



*Gently massage the foam in until it is absorbed and allow the areas to dry naturally.*

*When applying to the scalp, move the hair away so that the foam can be applied directly to each affected area.*



*Wash your hands immediately after applying Luxiq, and discard any unused dispensed medication.*



*Do not wash or rinse the treated areas immediately after applying Luxiq.*

- Do not use this medication for any condition other than the one for which it was prescribed.
- **Luxiq is for external use only.**
- **Keep the foam away from your eyes**, as it will sting. If the foam gets into your eyes, rinse well with cold water. If the stinging continues, contact your doctor immediately.

**WHAT YOU SHOULD KNOW ABOUT LUXIQ:**

#### **What to do if you miss an application**

If you forget to apply Luxiq at the scheduled time, use it as soon as you remember, and then go back to your regular schedule. If you remember at or about the time of your next daily application, apply that

dose and continue with your normal application schedule. If you miss several doses, tell your doctor at your next appointment.

### **About side effects**

As with all medications, there may be some side effects. The most frequent side effects associated with the use of Luxiq include mild burning, stinging, or itching at the site of application. These side effects typically disappear shortly after application.

Let your doctor know if you notice any of the following:

- Any unusual effects that you do not understand.
- Affected areas that do not seem to be healing after several weeks of using the foam.

### **Important safety notes**

- The treated areas should not be bandaged or covered unless directed by your doctor.
- Keep this and all medicines out of the reach of children.
- Store the can at controlled room temperature 20° to 25°C (68° to 77°F) and protect it from direct sunlight, as this is a pressurized container.
- **Keep away from and do not spray near fire, open flame, or direct heat--this product is flammable.** Do not smoke while using or holding the can. Keep the can away from all sources of ignition. Do not pierce or burn the can, and never throw the can in a fire, even if empty.
- When you have finished your treatment, dispose of the can safely. A completely empty can is recyclable.
- Do not use the foam after the expiration date shown on the bottom of the can.
- Do not give Luxiq to anyone else. Your doctor has prescribed this medicine for your use only.

For additional information visit [www.luxiq.com](http://www.luxiq.com)

Questions: Call Mylan at 1-877-446-3679 (1-877-4-INFO-RX). Side effects should be reported to this number.

LUXIQ is a registered trademark of Stiefel Laboratories, Inc., a GSK Company, exclusively licensed to the Mylan Companies

©2018 Delcor Asset Corporation, a Mylan Company

Manufactured for:

**Mylan Pharmaceuticals Inc.**  
Morgantown, WV 26505 U.S.A.

Manufactured by:

**DPT Laboratories, Ltd.**  
San Antonio, TX 78215 U.S.A.

140989-0418

Revised: 4/2018

DPT:BETAFON:R1

**PRINCIPAL DISPLAY PANEL – 0.12%**

**NDC 0378-8187-50**

**Luxiq®**  
(betamethasone  
valerate)

Foam, 0.12%

**Rx only 50 g**

**FOR DERMATOLOGIC USE ONLY.  
NOT FOR OPHTHALMIC USE.**

Invert can and then  
press firmly to  
dispense.



**Description:** Luxiq<sup>®</sup> (betamethasone valerate) Foam, 0.12%, contains 1.2 mg betamethasone valerate, USP, per gram in a thermolabile hydroethanolic foam vehicle consisting of cetyl alcohol, citric acid, ethanol (60.4%), polysorbate 60, potassium citrate, propylene glycol, purified water, and stearyl alcohol pressurized with a hydrocarbon (propane/butane) propellant.

**Usual Dosage:** See accompanying prescribing information. For application to the scalp, invert can and dispense a small amount of Luxiq onto a saucer or other cool surface. (Do not dispense directly onto hands, as foam will begin to melt immediately upon contact with warm skin.) Pick up small amounts of foam with fingers and gently massage into affected area until foam disappears. Repeat until entire affected scalp area is treated. Apply twice daily, once in the morning and once at night.

**WARNING: FLAMMABLE. AVOID FIRE, FLAME, OR SMOKING DURING AND IMMEDIATELY FOLLOWING APPLICATION.**

Keep away from eyes or other mucous membranes. Keep this and all medication out of the reach of children.

Contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120°F (49°C).

**Store at 20° to 25°C (68° to 77°F).**



**[See USP Controlled Room  
Temperature.]**

**CFC FREE**

Manufactured for:

**Mylan Pharmaceuticals Inc.**

Morgantown, WV 26505 U.S.A.

Manufactured by:

**DPT Laboratories, Ltd.**

San Antonio, TX 78215 U.S.A.

For additional information, visit

**luxiq.com**

Questions: Call Mylan at

1-877-446-3679 (1-877-4-INFO-RX).

Side effects should be reported to  
this number.

**Mylan.com**

DPT:8187:50:1C:R1

117518-0418

©2018 Delcor Asset Corporation,  
a Mylan Company

LUXIQ is a registered trademark of Stiefel  
Laboratories, Inc., a GSK Company, exclusively  
licensed to the Mylan Companies.

(betamethasone  
valerate)  
Foam, 0.12%



NDC 0378-8187-50

NDC 0378-8187-50



(betamethasone  
valerate)  
Foam, 0.12%



(betamethasone  
valerate)  
Foam, 0.12%

FOR DERMATOLOGIC USE ONLY.  
NOT FOR OPHTHALMIC USE.

Invert can and then  
press firmly to  
dispense.



(betamethasone  
valerate)  
Foam, 0.12%



(betamethasone  
valerate)  
Foam, 0.12%

FOR DERMATOLOGIC USE ONLY.  
NOT FOR OPHTHALMIC USE.

Invert can and then  
press firmly to  
dispense.



Description: Luxiq® (betamethasone valerate) Foam, 0.12%, contains 1.2 mg betamethasone valerate, USP, per gram in a thermolabile hydroethanolic foam vehicle consisting of cetyl alcohol, citric acid, ethanol (60.4%), polysorbate 60, potassium citrate, propylene glycol, purified water, and stearyl alcohol pressurized with a hydrocarbon (propane/butane) propellant.

Usual Dosage: See accompanying prescribing information. For application to the scalp, invert can and dispense a small amount of Luxiq onto a saucer or other cool surface. (Do not dispense directly onto hands, as foam will begin to melt immediately upon contact with warm skin.) Pick up small amounts of foam with fingers and gently massage into affected area until foam disappears. Repeat until entire affected scalp area is treated. Apply twice daily, once in the morning and once at night.

**WARNING: FLAMMABLE. AVOID FIRE, FLAME, OR SMOKING DURING AND IMMEDIATELY FOLLOWING APPLICATION.**

Keep away from eyes or other mucous membranes. Keep this and all medication out of the reach of children.

Contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120°F (49°C).

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]



N 3 0378-8187-50 6

CFC FREE

Manufactured for:  
Mylan Pharmaceuticals Inc.  
Morgantown, WV 26505 U.S.A.

Manufactured by:  
DPT Laboratories, Ltd.  
San Antonio, TX 78215 U.S.A.

For additional information, visit  
luxiq.com

Questions: Call Mylan at  
1-877-446-3679 (1-877-4-INFO-RX).  
Side effects should be reported to  
this number.

Mylan | [Mylan.com](http://Mylan.com)

DPT8187-50:1C:R1  
117518-0418

©2018 Delcor Asset Corporation,  
a Mylan Company

LUXIQ is a registered trademark of Stiefel  
Laboratories, Inc., a GSK Company, exclusively  
licensed to the Mylan Companies.

Rx only 50 g

Rx only 50 g





## LUXIQ

betamethasone valerate aerosol, foam

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0378-8187
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BETAMETHASONE VALERATE (UNII: 9IFA5XM7R2) (BETAMETHASONE - UNII:9842X06Q6M)	BETAMETHASONE VALERATE	1.2 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
CETYL ALCOHOL (UNII: 936JST6JCN)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
ALCOHOL (UNII: 3K9958V90M)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
STEARYL ALCOHOL (UNII: 2KR89I4HIY)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0378-8187-50	50 g in 1 CAN; Type 0: Not a Combination Product	12/17/2018	
2	NDC:0378-8187-01	100 g in 1 CAN; Type 0: Not a Combination Product	12/17/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020934	12/17/2018	

