

## **IBUPROFEN- ibuprofen capsule, liquid filled**

Hy-Vee

-----

### **DRUG FACTS**

#### **Active ingredient (in each capsule)**

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever/fever reducer

#### **Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

#### **Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stool
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

### **Inactive ingredients**

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

### **Questions or Comments?**

Call toll free **1-800-289-8343 Monday- Friday 9AM-4PM CST**

### **Principal Display Panel**

†**COMPARE TO the Active ingredient in Advil® Liqui-Gels®**

### **Ibuprofen**

**Capsules, 200 mg**

Pain Reliever/Fever Reducer (NSAID)

Softgels\*\*

\*\*LIQUID FILLED CAPSULES

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**DISTRIBUTED BY HY-VEE, INC., WEST DES MOINES, IA 50266**

**SATISFACTION GUARANTEED, CALL 1-800-289-8343 [www.hy-vee.com](http://www.hy-vee.com)**

### **Product Label**

Exp. Date:

Rev# 03/12  
PLD-47E F0030306



DISTRIBUTED BY HY-VEE, INC., WEST DES MOINES, IA 50266  
SATISFACTION GUARANTEED. CALL 1-800-289-8343  
www.hy-vee.com

HyVee. health

NDC 42507-198-48

# Ibuprofen Capsules 200 mg

Pain Reliever/Fever Reducer (NSAID)



Actual Size

**BONUS SIZE**  
40 + 8 FREE

**48** SOFTGELS\*\*

<b>Drug Facts</b>	<b>Purpose</b>
<b>Active ingredient (in each capsule)</b> Soubulized ibuprofen equal to 200 mg ibuprofen (NSAID)*. Pain reliever/fever reducer (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug	
<b>Uses</b>	
<ul style="list-style-type: none"><li>temporarily relieves minor aches and pains due to:<ul style="list-style-type: none"><li>headache</li><li>muscular aches</li><li>minor pain of arthritis</li><li>toothache</li><li>backache</li><li>menstrual cramps</li></ul></li><li>temporarily reduces fever</li></ul>	
<b>Warnings</b>	
<b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: <ul style="list-style-type: none"><li>hives</li><li>facial swelling</li><li>skin reddening</li><li>stomach</li><li>shock</li></ul> If an allergic reaction occurs, stop use and seek medical help right away.	<ul style="list-style-type: none"><li>asthma (wheezing)</li><li>rash</li><li>blisters</li></ul>
<b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: <ul style="list-style-type: none"><li>are age 60 or older</li><li>have had stomach ulcers or bleeding problems</li><li>take a blood thinning (anticoagulant) or steroid drug</li><li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li><li>have 3 or more alcoholic drinks every day while using this product</li><li>take more or for a longer time than directed</li></ul>	
<b>Do not use</b>	
<ul style="list-style-type: none"><li>if you have ever had an allergic reaction to any other pain reliever/fever reducer</li><li>right before or after heart surgery</li></ul>	

<b>Drug Facts (continued)</b>	
<b>Ask a doctor before use if</b>	
<ul style="list-style-type: none"><li>the stomach bleeding warning applies to you</li><li>you have problems or serious side effects from taking pain relievers or fever reducers</li><li>you have a history of stomach problems, such as heartburn</li><li>you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma</li><li>you are taking a diuretic</li></ul>	
<b>Ask a doctor or pharmacist before use if you are</b>	
<ul style="list-style-type: none"><li>under a doctor's care for any serious condition</li><li>taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li><li>taking any other drug</li></ul>	
<b>When using this product</b>	
<ul style="list-style-type: none"><li>take with food or milk, if stomach upset occurs</li><li>the risk of heart attack or stroke may increase if you use more than directed or for longer than directed</li></ul>	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"><li>you experience any of the following signs of stomach bleeding:<ul style="list-style-type: none"><li>feel faint</li><li>vomit blood</li><li>have bloody or black stools</li><li>have stomach pain that does not get better</li><li>pain gets worse or lasts more than 10 days</li><li>fever gets worse or lasts more than 3 days</li><li>redness or swelling is present in the painful area</li><li>any new symptoms appear</li></ul></li></ul>	
<b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	

<b>Drug Facts (continued)</b>	
<b>Directions</b>	
<ul style="list-style-type: none"><li>do not take more than directed</li><li>the smallest effective dose should be used</li><li>adults and children 12 years and over:<ul style="list-style-type: none"><li>take 1 capsule every 4 to 6 hours while symptoms persist</li><li>if pain or fever does not respond to 1 capsule, 2 capsules may be used</li></ul></li><li>do not exceed 6 capsules in 24 hours, unless directed by a doctor</li><li>children under 12 years: ask a doctor</li></ul>	
<b>Other information</b>	
<ul style="list-style-type: none"><li>each capsule contains: potassium 20 mg</li><li>read all warnings and directions before use. Keep carbox. store at 20° to 25°C (68° to 77°F)</li><li>avoid excessive heat above 40°C (104°F). Protect from light.</li></ul>	
<b>Inactive ingredients</b>	FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol
<b>Questions or comments?</b>	
Call toll free 1-800-289-8343 Monday-Friday 9AM-4PM CST	

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

## Hy-Vee Ibuprofen Capsules 200 mg

**IBUPROFEN**

ibuprofen capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:42507-198
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

**Inactive Ingredients**

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

**Product Characteristics**

<b>Color</b>	BLUE	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	IB200
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42507-198-48	1 in 1 BOX	01/15/2013	
1		48 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:42507-198-27	80 in 1 BOTTLE; Type 0: Not a Combination Product	01/15/2013	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA078682	01/15/2013	
------	------------	------------	--

**Labeler** - Hy-Vee (006925671)

Revised: 6/2016

Hy-Vee