LIL DRUG STORE PAIN RELIEVER EXTRA STRENGTH- acetaminophen tablet Lil' Drug Store Products, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Lil' Drug Store ® Pain Reliever

Extra Strength

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- adults and children 12 years and over
 - take 2 caplets every 6 hours while symptoms last
 - do not take more than 6 caplets in 24 hours
 - do not take for more than 10 days unless directed by a doctor
- children under 12 years: do not use

Other information

store between 68-77°F (20-25°C)

Inactive ingredients

corn starch, croscarmellose sodium 1 , hypromellose 1 , lactose monohydrate 1 , magnesium stearate 1 , maltodextrin 1 , medium-chain triglycerides 1 , mineral oil 1 , polydextrose 1 , polyethylene glycol 1 , polyvinyl alcohol 1 , povidone, purified water 1 , sodium starch glycolate 1 , stearic acid 1 , talc 1 , titanium dioxide

Questions or comments?

Call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

Product distributed by:

Lil' Drug Store Products, Inc. 9300 Earhart Lane SW Cedar Rapids, IA 52404

PRINCIPAL DISPLAY PANEL - 500 mg Caplet Bottle Carton - NDC 66715-6817-4

QUALITY GUARANTEED

Compare to the Active Ingredient in Tylenol® Extra Strength**

Extra Strength

Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg

ACTUAL SIZE

24

Caplets

Lil

Drug Store ®

Extra Strength Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg

Caplets





Compare to the Active Ingredient in Tylenol® Extra Strength**

Extra Strength Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg





PLD-E108D

FC005264

Product distributed by: Lil' Drug Store Products, Inc. 9300 Earhart Lane SW Cedar Rapids, IA 52404



Lot No: Exp Date:

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSI

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg Pain reliever/ fever reducer

Purpose

Uses

- temporarily relieves minor aches and pains due to:
- headache
- the common cold
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■skin reddening ■blisters ■rash If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.


Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away

(1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- adults and children 12 years and over
- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours
- do not take for more than 10 days unless directed by a doctor
- children under 12 years: do not use







PRINCIPAL DISPLAY PANEL - 500 mg Caplet Bottle Carton - NDC 66715-6827-4

QUALITY GUARANTEED

Compare to the Active Ingredient in Tylenol® Extra Strength**

Extra Strength

Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg

ACTUAL SIZE

24

Caplets

Lil'

Drug Store ®

Extra Strength Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg

Caplets





Compare to the Active Ingredient in Tylenol® Extra Strength**

Extra Strength Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg





PLD-E108D

FC005264

Product distributed by: Lil' Drug Store Products, Inc. 9300 Earhart Lane SW Cedar Rapids, IA 52404



Lot No: Exp Date:

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSI

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg Pain reliever/ fever reducer

Purpose

Uses

- temporarily relieves minor aches and pains due to:
- headache
- the common cold
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■skin reddening ■blisters ■rash If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.


Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away

(1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- adults and children 12 years and over
- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours
- do not take for more than 10 days unless directed by a doctor
- children under 12 years: do not use







PRINCIPAL DISPLAY PANEL - 500 mg Caplet Bottle Carton - NDC 66715-6837-4

QUALITY GUARANTEED

Compare to the Active Ingredient in Tylenol® Extra Strength**

Extra Strength

Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg

ACTUAL SIZE

24

Caplets

Lil'

Drug Store ®

Extra Strength Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg

Caplets





Compare to the Active Ingredient in Tylenol® Extra Strength**

Extra Strength Pain Reliever

Acetaminophen

Pain Reliever/Fever Reducer, 500 mg





PLD-E108D

FC005264

Product distributed by: Lil' Drug Store Products, Inc. 9300 Earhart Lane SW Cedar Rapids, IA 52404



Lot No: Exp Date:

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSI

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg Pain reliever/ fever reducer

Purpose

Uses

- temporarily relieves minor aches and pains due to:
- headache
- the common cold
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■skin reddening ■blisters ■rash If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.


Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away

(1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- adults and children 12 years and over
- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours
- do not take for more than 10 days unless directed by a doctor
- children under 12 years: do not use







LIL DRUG STORE PAIN RELIEVER EXTRA STRENGTH

acetaminophen tablet

| acetaminophen tablet | | | | | |
|----------------------------|----------------|-----------------|------------|-----------|----------|
| | | | | | |
| Product Information | | | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Sour | rce) | NDC:66715 | 5-6817 |
| Route of Administration | ORAL | | | | |
| | | | | | |
| Active Ingredient/Active | Moiety | | | | |
| | <u> </u> | | | | |
| Ingre | edient Name | | Basis of S | trength | Strength |

| Inactive Ingredients | |
|--------------------------------------------------------|----------|
| Ingredient Name | Strength |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U) | |
| MINERAL OIL (UNII: T5L8T28FGP) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

| Product Characterist | roduct Characteristics | | | | |
|-----------------------------|------------------------|--------------|----------|--|--|
| Color | white | Score | no score | | |
| Shape | OVAL | Size | 18mm | | |
| Flavor | | Imprint Code | P500 | | |
| Contains | | | | | |

| Packaging | | | | | | | |
|--------------------|-----|--------------------------------------------------------|-------------------------|-----------------------|--|--|--|
| # Item C | ode | Package Description | Marketing Start Date | Marketing End Date | | | |
| NDC:6671 6817-4 | 1 i | in 1 CARTON | 04/15/2019 | 08/12/2024 | | | |
| 1 | | in 1 BOTTLE, PLASTIC; Type 0: Not a ombination Product | | | | | |

| Marketing Information | | | | |
|-------------------------|---------------------------------------------|-------------------------|-----------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC monograph not final | part343 | 01/10/2015 | 08/12/2024 | |
| | | | | |

LIL DRUG STORE PAIN RELIEVER EXTRA STRENGTH

acetaminophen tablet

| Product Information | | | | | |
|-------------------------|----------------|--------------------|----------------|--|--|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:66715-6827 | | |
| Route of Administration | ORAL | | | | |

| Active Ingredient/Active Moiety | | | | |
|--------------------------------------------------------------------|--------------------------|----------|--|--|
| Ingredient Name | Basis of Strength | Strength | | |
| ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) | ACETAMINOPHEN | 500 mg | | |

| Inactive Ingredients | |
|--------------------------------------------------------|----------|
| Ingredient Name | Strength |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U) | |
| MINERAL OIL (UNII: T5L8T28FGP) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) | |
| POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

| Product Characteristics | | | | | | |
|-------------------------|-------|--------------|----------|--|--|--|
| Color | white | Score | no score | | | |
| Shape | OVAL | Size | 18mm | | | |
| Flavor | | Imprint Code | AV;0821 | | | |
| Contains | | | | | | |

| | Packaging | | | | | | |
|---|-----------|----------------------|------------------------------------------------------------|-------------------------|-----------------------|--|--|
| | # | Item Code | Package Description | Marketing Start Date | Marketing End Date | | |
| | | NDC:66715- 6827-4 | 1 in 1 CARTON | 04/15/2019 | 08/12/2024 | | |
| | 1 | | 24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | | | |
| Н | | | | | | | |

| Marketing Information | | | | |
|-------------------------|---------------------------------------------|-------------------------|-----------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC monograph not final | part343 | 01/10/2015 | 08/12/2024 | |
| | | | | |

LIL DRUG STORE PAIN RELIEVER EXTRA STRENGTH

acetaminophen tablet

Product Type HUMAN OTC DRUG Item Code (Source) NDC:66715-6837

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 500 mg

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------------------------|----------|
| STARCH, CORN (UNII: 08232NY3SJ) | |
| CROSCARMELLOSE SODIUM (UNII: M280L1HH48) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U) | |
| MINERAL OIL (UNII: T5L8T28FGP) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B) | |

Product Characteristics

STEARIC ACID (UNII: 4ELV7Z65AP)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

TALC (UNII: 7SEV7J4R1U)

| Color | white | Score | no score |
|----------|-------|--------------|----------|
| Shape | OVAL | Size | 18mm |
| Flavor | | Imprint Code | TCL;341 |
| Contains | | | |

| P | Packaging | | | | | | |
|---|----------------------|------------------------------------------------------------|-------------------------|-----------------------|--|--|--|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date | | | |
| 1 | NDC:66715- 6837-4 | 1 in 1 CARTON | 04/15/2019 | 08/12/2024 | | | |
| 1 | | 24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | | | | |

| Marketing Information | | | | | |
|-------------------------|---------------------------------------------|-------------------------|-----------------------|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | |
| OTC monograph not final | part343 | 01/10/2015 | 08/12/2024 | | |
| | part343 | 01/10/2015 | 08/12/2024 | | |

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2021 Lil' Drug Store Products, Inc.