

**CAREONE ACETAMINOPHEN- acetaminophen tablet, film coated**  
**American Sales Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**American Sales Company Acetaminophen Drug Facts**

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 caplets every 6 hours while symptoms last</li><li>• do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, mica-based pearlescent pigment, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid

## Questions or comments?

1-800-719-9260

### Package/Label Principal Display Panel

Compare to the active ingredient in Tylenol® Extra Strength Rapid Release Gels

Extra Strength

ACETAMINOPHEN

500mg

Actual Size

Pain Reliever/Fever Reducer

Rapid Release

For Adults

Gluten Free

OUR PHARMACISTS RECOMMEND

100 CAPLETS



**DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

\*\* This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Tylenol® Extra Strength Rapid Release Gels.

DISTRIBUTED BY FOODHOLD U.S.A., LLC  
LANDOVER, MD 20785 · 1-877-846-9949  
©2017 S&S Brands, LLC

Quality guaranteed or your money back.

## Drug Facts

### Active ingredient (in each caplet)

### Purpose

Acetaminophen 500 mg.....Pain reliever/fever reducer

### Uses

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - minor pain of arthritis
  - backache
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition. ▶

## Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years


ask a doctor

### Other information

store at 20-25°C (68-77°F)

**Inactive ingredients** croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, mica-based pearlescent pigment, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid

### Questions or comments? 1-800-719-9260



## CAREONE ACETAMINOPHEN

acetaminophen tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-786
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPAS AT 5%) (UNII: 68401960MK)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

### Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	3S0
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-786-78	1 in 1 CARTON	12/21/2017	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/21/2017	

**Labeler** - American Sales Company (809183973)

Revised: 12/2019

American Sales Company