

PUBLIX STOMACH RELIEF ULTRA STRENGTH- bismuth subsalicylate suspension
PUBLIX SUPER MARKETS, INC

PUBLIX Ultra Strength Pink Bismuth Subsalicylate 525 mg Drug Facts

Active ingredient (in each 15 mL dose)

Bismuth subsalicylate 525 mg

Purposes

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- only use dose cup provided
- **adults and children 12 years and over:**
- 15 mL (1 dose) every ½ hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 15 mL (1 dose) every ½ hour as needed overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 15 mL dose cup contains:** potassium 7 mg, sodium 5 mg
- **salicylate 230 mg**
- low sodium
- sugar free

- store at room temperature
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Do not use if imprinted shrink band is missing or broken.

Principal Display Panel

NDC 41415-707-12

***Compare to the active ingredient in Pepto-Bismol® Ultra Strength**

Ultra Strength

pink bismuth

bismuth subsalicylate

Upset Stomach Reliever/Antidiarrheal

Natural Cherry flavor

Relieves:

- Nausea
- Heartburn
- Indigestion
- Upset Stomach
- Diarrhea

12 FL. OZ. (354 mL)

2X CONCENTRATED FORMULA**

Distributed by:

***This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Ultra Strength.**

7.25" Width

3.5625" Height

Drug Facts (continued)

Directions ■ shake well before use ■ only use dose cup provided
 ■ **adults and children 12 years and over** ■ 15 mL (1 dose) every ½ hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea. ■ 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 ■ do not exceed 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ **children under 12 years:** ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information
 ■ each 15 mL dose cup contains: potassium 7 mg, sodium 5 mg
 ■ salicylate 230 mg ■ low sodium ■ sugar free
 ■ store at room temperature
 ■ protect from freezing
 ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavors, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

DISTRIBUTED BY
 PUBLIX SUPER MARKETS, INC.,
 3300 PUBLIX CORPORATE PARKWAY
 LAKELAND, FL 33811
 1-888-267-3037 publix.com
 PUBLIX GUARANTEE:
 COMPLETE SATISFACTION
 OR YOUR MONEY BACK

SCAN HERE FOR MORE INFORMATION

Publix
 707CH12PUBLR

LOT: _____
 EXP: _____

0 4 14 15 54073 6



NDC 41415-707-12

STOMACH RELIEF
ultrastrength
BISMUTH SUBSALICYLATE 525 mg
 UPSET STOMACH RELIEVER/
 ANTIDIARRHEAL

Relieves

- Nausea
- Heartburn
- Indigestion
- Upset stomach
- Diarrhea

2X CONCENTRATED FORMULA**

CHERRY FLAVORED

*Compare to the active ingredient in Pepto-Bismol® Ultra Strength

12 FL OZ (354 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

Drug Facts

Active ingredient (in each 15 mL dose)	Purpose
Bismuth subsalicylate 525 mg	Upset stomach reliever and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:
 ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are
 ■ allergic to salicylates (including aspirin)
 ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
 ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. ▶

*This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Ultra Strength.

PUBLIX STOMACH RELIEF ULTRA STRENGTH

bismuth subsalicylate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41415-707
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
WATER (UNII: 059QF0KO0R)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	PINK (viscous)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41415-707-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/24/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/24/2022	

Labeler - PUBLIX SUPER MARKETS, INC (006922009)

Revised: 11/2023

PUBLIX SUPER MARKETS, INC