

**RIFABUTIN- rifabutin capsule**  
**Lupin Pharmaceuticals, Inc.**


-----  
**Rifabutin Capsules USP, 150 mg**

**Rx only**

**DESCRIPTION**

Rifabutin Capsules USP for oral administration contain 150 mg of the rifamycin antimycobacterial agent rifabutin, USP, per capsule, along with the inactive ingredients crospovidone, gelatin, iron oxide red, magnesium stearate, microcrystalline cellulose, potassium hydroxide, shellac, silicon dioxide, sodium lauryl sulphate, and titanium dioxide.

The chemical name for rifabutin is 1',4-didehydro-1-deoxy-1,4-dihydro-5'-(2-methylpropyl)-1-oxorifamycin XIV (Chemical Abstracts Service, 9th Collective Index) or (9*S*, 12*E*, 14*S*, 15*R*, 16*S*, 17*R*, 18*R*, 19*R*, 20*S*, 21*S*, 22*E*, 24*Z*)-6,16,18,20-tetrahydroxy-1'-isobutyl-14-methoxy-7,9,15,17,19,21,25-heptamethyl-spiro [9,4-(epoxypentadeca[1,11,13]trienimino)-2*H*-furo[2',3':7,8] naphth[1,2-*d*]imidazole-2,4'-piperidine]-5,10,26-(3*H*,9*H*)-trione-16-acetate. Rifabutin has a molecular formula of C<sub>46</sub>H<sub>62</sub>N<sub>4</sub>O<sub>11</sub>, a molecular weight of 847.02 and the following structure:



Rifabutin is a red-violet powder soluble in chloroform and methanol, sparingly soluble in ethanol, and very slightly soluble in water (0.19 mg/mL). Its log P value (the base 10 logarithm of the partition coefficient between n-octanol and water) is 3.2 (n-octanol/water).

**CLINICAL PHARMACOLOGY**

**Pharmacokinetics**

**Absorption**

Following a single oral dose of 300 mg to nine healthy adult volunteers, rifabutin was readily absorbed from the gastrointestinal tract with mean ( $\pm$ SD) peak plasma levels ( $C_{max}$ ) of 375 ( $\pm$ 267) ng/mL (range: 141 to 1033 ng/mL) attained in 3.3 ( $\pm$ 0.9) hours ( $T_{max}$  range: 2 to 4 hours). Absolute bioavailability assessed in five HIV-positive patients, who received both oral and intravenous doses, averaged 20%. Total recovery of radioactivity in the urine indicates that at least 53% of the orally administered rifabutin dose is absorbed from the gastrointestinal tract. The bioavailability of rifabutin from the

capsule dosage form, relative to an oral solution, was 85% in 12 healthy adult volunteers. High-fat meals slow the rate without influencing the extent of absorption from the capsule dosage form. Plasma concentrations post- $C_{max}$  declined in an apparent biphasic manner. Pharmacokinetic dose-proportionality was established over the 300 mg to 600 mg dose range in nine healthy adult volunteers (crossover design) and in 16 early symptomatic human immunodeficiency virus (HIV)-positive patients over a 300 mg to 900 mg dose range.

### **Distribution**

Due to its high lipophilicity, rifabutin demonstrates a high propensity for distribution and intracellular tissue uptake. Following intravenous dosing, estimates of apparent steady-state distribution volume ( $9.3 \pm 1.5$  L/kg) in five HIV-positive patients exceeded total body water by approximately 15-fold. Substantially higher intracellular tissue levels than those seen in plasma have been observed in both rat and man. The lung-to-plasma concentration ratio, obtained at 12 hours, was approximately 6.5 in four surgical patients who received an oral dose. Mean rifabutin steady-state trough levels ( $C_{p, min}^{SS}$ ; 24-hour post-dose) ranged from 50 to 65 ng/mL in HIV-positive patients and in healthy adult volunteers. About 85% of the drug is bound in a concentration-independent manner to plasma proteins over a concentration range of 0.05 to 1  $\mu$ g/mL. Binding does not appear to be influenced by renal or hepatic dysfunction. Rifabutin was slowly eliminated from plasma in seven healthy adult volunteers, presumably because of distribution-limited elimination, with a mean terminal half-life of 45 ( $\pm$  17) hours (range: 16 to 69 hours). Although the systemic levels of rifabutin following multiple dosing decreased by 38%, its terminal half-life remained unchanged.

### **Metabolism**

Of the five metabolites that have been identified, 25-O-desacetyl and 31-hydroxy are the most predominant, and show a plasma metabolite:parent area under the curve ratio of 0.10 and 0.07, respectively. The former has an activity equal to the parent drug and contributes up to 10% to the total antimicrobial activity.

### **Excretion**

A mass-balance study in three healthy adult volunteers with  $^{14}$ C-labeled rifabutin showed that 53% of the oral dose was excreted in the urine, primarily as metabolites. About 30% of the dose is excreted in the feces. Mean systemic clearance ( $CL_S/F$ ) in healthy adult volunteers following a single oral dose was 0.69 ( $\pm$  0.32) L/hr/kg (range: 0.46 to 1.34 L/hr/kg). Renal and biliary clearance of unchanged drug each contribute approximately 5% to  $CL_S/F$ .

## **Pharmacokinetics in Special Populations**

### **Geriatric**

Compared to healthy volunteers, steady-state pharmacokinetics of rifabutin are more variable in elderly patients (>70 years).

### **Pediatric**

The pharmacokinetics of rifabutin have not been studied in subjects under 18 years of age.

### **Renal Impairment**

The disposition of rifabutin (300 mg) was studied in 18 patients with varying degrees of renal function. Area under plasma concentration time curve (AUC) increased by about 71% in patients with severe renal impairment (creatinine clearance below 30 mL/min) compared to patients with creatinine clearance ( $Cr_{Cl}$ ) between 61 to 74 mL/min. In patients with mild to moderate renal impairment ( $Cr_{Cl}$  between 30 to 61 mL/min), the AUC increased by about 41%. In patients with severe renal impairment, carefully monitor for rifabutin associated adverse events. A reduction in the dosage of rifabutin is recommended for patients with  $Cr_{Cl} < 30$  mL/min if toxicity is suspected (see *Dosage and Administration*).

### **Hepatic Impairment**

Mild hepatic impairment does not require a dose modification. The pharmacokinetics of rifabutin in patients with moderate and severe hepatic impairment is not known.

### **Malabsorption in HIV-Infected Patients**

Alterations in gastric pH due to progressing HIV disease has been linked with malabsorption of some drugs used in HIV-positive patients (e.g., rifampin, isoniazid). Drug serum concentrations data from AIDS patients with varying disease severity (based on CD4+ counts) suggests that rifabutin absorption is not influenced by progressing HIV disease.

### **Drug-Drug Interactions** (see also *Precautions-Drug Interactions*)

Multiple dosing of rifabutin has been associated with induction of hepatic metabolic enzymes of the CYP3A subfamily. Rifabutin's predominant metabolite (25-desacetyl rifabutin: LM565), may also contribute to this effect. Metabolic induction due to rifabutin is likely to produce a decrease in plasma concentrations of concomitantly administered drugs that are primarily metabolized by the CYP3A enzymes. Similarly concomitant medications that competitively inhibit the CYP3A activity may increase plasma concentrations of rifabutin.

## **CLINICAL STUDIES**

Two randomized, double-blind clinical trials (Study 023 and Study 027) compared rifabutin (300 mg/day) to placebo in patients with CDC-defined AIDS and CD4 counts  $\leq 200$  cells/ $\mu$ L. These studies accrued patients from 2/90 through 2/92. Study 023 enrolled 590 patients, with a median CD4 cell count at study entry of 42 cells/ $\mu$ L (mean 61). Study 027 enrolled 556 patients with a median CD4 cell count at study entry of 40 cells/ $\mu$ L (mean 58).

Endpoints included the following:

- (1) MAC bacteremia, defined as at least one blood culture positive for *Mycobacterium avium* complex (MAC) bacteria.
- (2) Clinically significant disseminated MAC disease, defined as MAC bacteremia accompanied by signs or symptoms of serious MAC infection, including one or more of the following: fever, night sweats, rigors, weight loss, worsening anemia, and/or elevations in alkaline phosphatase.
- (3) Survival.

### **MAC Bacteremia**

Participants who received rifabutin were one-third to one-half as likely to develop MAC bacteremia as were participants who received placebo. These results were statistically significant (Study 023:  $p < 0.001$ ; Study 027:  $p = 0.002$ ).

In Study 023, the one-year cumulative incidence of MAC bacteremia, on an intent to treat basis, was 9% for patients randomized to rifabutin and 22% for patients randomized to placebo. In Study 027, these rates were 13% and 28% for patients receiving rifabutin and placebo, respectively.

Most cases of MAC bacteremia (approximately 90% in these studies) occurred among participants whose CD4 count at study entry was  $\leq 100$  cells/ $\mu$ L. The median and mean CD4 counts at onset of MAC bacteremia were 13 cells/ $\mu$ L and 24 cells/ $\mu$ L, respectively. These studies did not investigate the optimal time to begin MAC prophylaxis.

### **Clinically Significant Disseminated MAC Disease**

In association with the decreased incidence of bacteremia, patients on rifabutin showed reductions in the signs and symptoms of disseminated MAC disease, including fever, night sweats, weight loss, fatigue, abdominal pain, anemia, and hepatic dysfunction.

### **Survival**

The one-year survival rates in Study 023 were 77% for the group receiving rifabutin and

77% for the placebo group. In Study 027, the one-year survival rates were 77% for the group receiving rifabutin and 70% for the placebo group.

These differences were not statistically significant.

## **Microbiology**

### **Mechanism of Action**

Rifabutin inhibits DNA-dependent RNA polymerase in susceptible strains of *Escherichia coli* and *Bacillus subtilis* but not in mammalian cells. In resistant strains of *E. coli*, rifabutin, like rifampin, did not inhibit this enzyme. It is not known whether rifabutin inhibits DNA-dependent RNA polymerase in *Mycobacterium avium* or in *M. intracellulare* which comprise *M. avium* complex (MAC).

### **Susceptibility Testing**

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

## **INDICATIONS AND USAGE**

Rifabutin capsules USP are indicated for the prevention of disseminated *Mycobacterium avium* complex (MAC) disease in patients with advanced HIV infection.

## **CONTRAINDICATIONS**

Rifabutin capsules are contraindicated in patients who have had clinically significant hypersensitivity to rifabutin or to any other rifamycins.

Rifabutin Capsules are contraindicated in patients being treated with cabotegravir/rilpivirine prolonged-release injectable suspension (*see Precautions-Drug Interactions, Table 2*).

## **WARNINGS**

### **Tuberculosis**

Rifabutin capsules must not be administered for MAC prophylaxis to patients with active tuberculosis. Patients who develop complaints consistent with active tuberculosis while on prophylaxis with rifabutin should be evaluated immediately, so that those with active disease may be given an effective combination regimen of anti-tuberculosis medications. Administration of rifabutin as a single agent to patients with active tuberculosis is likely to lead to the development of tuberculosis that is resistant both to rifabutin and to rifampin.

There is no evidence that rifabutin is an effective prophylaxis against *M. tuberculosis*. Patients requiring prophylaxis against both *M. tuberculosis* and *Mycobacterium avium* complex may be given isoniazid and rifabutin concurrently.

Tuberculosis in HIV-positive patients is common and may present with atypical or extrapulmonary findings. Patients are likely to have a nonreactive purified protein derivative (PPD) despite active disease. In addition to chest X-ray and sputum culture, the following studies may be useful in the diagnosis of tuberculosis in the HIV-positive patient: blood culture, urine culture, or biopsy of a suspicious lymph node.

### **MAC Treatment with Clarithromycin**

When rifabutin is used concomitantly with clarithromycin for MAC treatment, a decreased dose of rifabutin is recommended due to the increase in plasma concentrations of rifabutin (*see Precautions-Drug Interactions, Table 2*).

### **Hypersensitivity and Related Reactions**

Hypersensitivity reactions may occur in patients receiving rifamycins. Signs and symptoms of these reactions may include hypotension, urticaria, angioedema, acute bronchospasm, conjunctivitis, thrombocytopenia, neutropenia or flu-like syndrome (weakness, fatigue, muscle pain, nausea, vomiting, headache, fever, chills, aches, rash, itching, sweats, dizziness, shortness of breath, chest pain, cough, syncope, palpitations). There have been reports of anaphylaxis with the use of rifamycins.

Monitor patients receiving rifabutin therapy for signs and/or symptoms of hypersensitivity reactions. If these symptoms occur, administer supportive measures and discontinue rifabutin.

### **Uveitis**

Due to the possible occurrence of uveitis, patients should also be carefully monitored when rifabutin is given in combination with clarithromycin (or other macrolides) and/or fluconazole and related compounds (see *Precautions-Drug Interactions, Table 2*). If uveitis is suspected, the patient should be referred to an ophthalmologist and, if considered necessary, treatment with rifabutin should be suspended (see also *Adverse Reactions*).

### ***Clostridioides difficile* Associated Diarrhea**

*Clostridioides difficile* associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including rifabutin capsules, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

### **Severe Cutaneous Adverse Reactions**

There have been reports of severe cutaneous adverse reactions (SCAR), such as Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), and acute generalized exanthematous pustulosis (AGEP) associated with rifabutin (see **ADVERSE REACTIONS**).

If patients develop a skin rash they should be monitored closely, and rifabutin discontinued if lesions progress. Specifically, for DRESS, a multi-system potential life-threatening SCAR, time to onset of the first symptoms may be prolonged. DRESS is a clinical diagnosis, and its clinical presentation remains the basis for decision making. An early withdrawal of rifabutin is essential because of the syndrome's mortality and visceral involvement (e.g., liver, bone marrow or kidney).

### **Antiretroviral and Anti-HCV Drug Interactions**

Protease inhibitors act as substrates or inhibitors of CYP3A4 mediated metabolism. Therefore, due to significant drug-drug interactions between protease inhibitors and rifabutin, their concomitant use should be based on the overall assessment of the patient and a patient-specific drug profile. The concomitant use of protease inhibitors may require at least a 50% reduction in rifabutin dose, and depending on the protease inhibitor, an adjustment of the antiretroviral drug dose. Increased monitoring for adverse events is recommended when using these drug combinations (see *Precautions-Drug Interactions*).

Rifabutin is a CYP3A inducer. Co-administration with antiretroviral drugs metabolized by CYP3A, including but not limited to products containing bicitgravir, elvitegravir, oral rilpivirine, or doravirine and anti-HCV drugs including but not limited to sofosbuvir (alone

or in combination) may decrease plasma concentrations of those drugs, which may lead to loss of virologic response and possible development of resistance. Therefore, co-administration with antiretroviral and anti-HCV drugs metabolized by CYP3A is not recommended or there may be a need to increase the dose of antiretroviral or anti-HCV drugs (see *Precautions-Drug Interactions*).

For further recommendations, please refer to the most recent prescribing information of the antiretrovirals or anti-HCV drugs or contact the specific manufacturer.

## PRECAUTIONS

### General

Because treatment with rifabutin capsules may be associated with neutropenia, and more rarely thrombocytopenia, physicians should consider obtaining hematologic studies periodically in patients receiving prophylaxis with rifabutin.

### Information for Patients

Patients should be advised of the signs and symptoms of both MAC and tuberculosis, and should be instructed to consult their physicians if they develop new complaints consistent with either of these diseases. In addition, since rifabutin may rarely be associated with myositis and uveitis, patients should be advised to notify their physicians if they develop signs or symptoms suggesting either of these disorders.

Urine, feces, saliva, sputum, perspiration, tears, and skin may be colored brown-orange with rifabutin and some of its metabolites. Soft contact lenses may be permanently stained. Patients to be treated with rifabutin should be made aware of these possibilities.

Diarrhea is a common problem caused by antibacterials which usually ends when the antibacterial is discontinued. Sometimes after starting treatment with antibacterials, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibacterial. If this occurs, patients should contact their physician as soon as possible.

### Drug Interactions

#### ***Effect of Rifabutin on the Pharmacokinetics of Other Drugs***

Rifabutin induces CYP3A enzymes and therefore may reduce the plasma concentrations of drugs metabolized by those enzymes. This effect may reduce the efficacy of standard doses of such drugs, which include itraconazole, clarithromycin, and saquinavir.

#### ***Effect of Other Drugs on Rifabutin Pharmacokinetics***

Some drugs that inhibit CYP3A may significantly increase the plasma concentration of rifabutin. Therefore, carefully monitor for rifabutin associated adverse events in those patients also receiving CYP3A inhibitors, which include fluconazole and clarithromycin. In some cases, the dosage of rifabutin may need to be reduced when it is coadministered with CYP3A inhibitors.

Table 2 summarizes the results and magnitude of the pertinent drug interactions assessed with rifabutin. The clinical relevance of these interactions and subsequent dose modifications should be judged in light of the population studied, severity of the disease, patient's drug profile, and the likely impact on the risk/benefit ratio.

**Table 2: Rifabutin Interaction Studies**

Co-administered drug	Dosing regimen of co-administered drug	Dosing regimen of rifabutin	Study population (n)	Effect on rifabutin	Effect on co-administered drug	Recommendation
<b>ANTIRETROVIRALS</b>						

Amprenavir	1200 mg twice a day for 10 days	300 mg once a day for 10 days	Healthy male subjects (6)	↑ AUC by 193%, ↑ C <sub>max</sub> by 119%	↔	Reduce rifabutin dose by at least 50%. Monitor closely for adverse reactions.
Atazanavir/ Ritonavir	300/100 mg once daily	150 mg twice weekly	Healthy adult subjects	48% ↑ in AUC, 149% ↑ C <sub>max</sub> of rifabutin. 990% ↑ in AUC, 677% ↑ C <sub>max</sub> of 25-O-desacetyl-rifabutin.	No significant change in pharmacokinetics.	A reduction in the dose of rifabutin (to 150 mg every other day or 3 times a week) is recommended. Increased monitoring for adverse reactions is warranted.
Bictegravir	75 mg once a day	300 mg once a day (fasted)	Healthy subjects	ND	↓ AUC by 38, ↓ C <sub>min</sub> by 56%, ↓ C <sub>max</sub> by 20%	Co-administration of Rifabutin with biktarvy (bictegravir/emtricitabin/ tenofovir alafenamide) is not recommended due to an expected decrease in tenofovir alafenamide in addition to the reported reduction in bictegravir. Refer to biktarvy prescribing information for additional information
Darunavir/Ritonavir	600/100 mg twice a day for 12 days	150 mg every other day for 12 days	Healthy HIV negative adults	No significant change in rifabutin pharmacokinetics.	57% ↑ in AUC, 42% ↑ C <sub>max</sub> of darunavir. 66% ↑ in AUC, 68% ↑ C <sub>max</sub> of ritonavir. 881% ↑ in AUC, 377% ↑ C <sub>max</sub> of 25-O-desacetyl-rifabutin.	A reduction in the dose of rifabutin (to 150 mg every other day or 3 times a week) is recommended. Increased monitoring for adverse reactions is warranted.
Delavirdine	400 mg three times a day	300 mg once a day	HIV-infected patients (7)	↑ AUC by 230%, ↑ C <sub>max</sub> by 128%	↓ AUC by 80%, ↓ C <sub>max</sub> by 75%, ↓ C <sub>min</sub> by 17%	<b>CONTRAINDICATED</b>
Didanosine	167 or 250 mg twice a day for 12 days	300 or 600 mg once a day for 12 days	HIV-infected patients (11)	↔	↔	
Dolutegravir	50 mg daily for 14 days	300 mg Daily for 14 days	Healthy adult subjects	ND	No significant change in dolutegravir pharmacokinetics at steady state.	
Doravirine	100 mg single dose	300 mg once a day for 16 days	Healthy subjects (12)	ND	↓ 50% in AUC, ↓ 68% in C <sub>24</sub> ↔ in C <sub>max</sub>	If concomitant use is necessary, increase the doravirine dosage as instructed in doravirine-containing product prescribing information.

Elvitegravir/ Cobicistat	150/50 mg daily	300 mg Daily Or 150 mg every other day	Healthy subjects (12)	No significant change in rifabutin pharma cokinetics. 6.3-fold ↑ in AUC, 4.8-fold ↑ C <sub>max</sub> of 25-O-desac etylrifabutin.	No change in elvitegravir except 67% ↓ C <sub>trough</sub> of Elvitegravir. No change in cobicistat exposure.	Co-administration of rifabutin with elvitegravir/ cobicistat is not recommended due to an expected decrease in elvitegravir exposure.
Etravirine	800 mg twice daily for 21 days	300 mg Daily on days 8 to 21	Healthy Volunteers (18)	No significant change in rifabutin pharmacoki netics.	37% ↓ in AUC, 37% ↓ in C <sub>max</sub> and 35% ↓ in C <sub>min</sub>	No dose adjustment of rifabutin is required when etravirine is not co-administered with protease inhibitor/ritonavir. Rifabutin should not be coadministered with etravirine and boosted PIs due to potential for decreased effectiveness of etravirine.
Fosamprenavir/ ritonavir	700 mg twice a day plus ritonavir 100 mg twice a day for 2 weeks	150 mg every other day for 2 weeks	Healthy subjects (15)	↔ AUC <sup>a</sup> ↓ C <sub>max</sub> by 15%	↑ AUC by 35% <sup>b</sup> , ↑ C <sub>max</sub> by 36%, ↑ C <sub>min</sub> by 36%	Reduce rifabutin dose by at least 75% (to a maximum 150 mg every other day or three times per week) when given with fosamprenavir/ritonavir combination.
Indinavir	800 mg three times a day for 10 days	300 mg once a day for 10 days	Healthy subjects (10)	↑ AUC by 173%, ↑ C <sub>max</sub> by 134%	↓ AUC by 34%, ↓ C <sub>max</sub> by 25%, ↓ C <sub>min</sub> by 39%	Reduce rifabutin dose by 50%, and increase indinavir dose from 800 mg to 1000 mg three times a day.
Lopinavir/ ritonavir	400/100 mg twice a day for 20 days	150 mg once a day for 10 days	Healthy subjects (14)	↑ AUC by 203% <sup>c</sup> ↓ C <sub>max</sub> by 112%	↔	Reduce rifabutin dose by at least 75% (to a maximum 150 mg every other day or three times per week) when given with lopinavir/ritonavir combination. Monitor closely for adverse reactions. Reduce rifabutin dosage further, as needed.
Saquinavir/ ritonavir	1000/100 mg twice a day for 14 or 22 days	150 mg every 3 days for 21 to 22 days	Healthy subjects	↑ AUC by 53% <sup>d</sup> ↑ C <sub>max</sub> by 88% (n=11)	↓ AUC by 13%, ↓ C <sub>max</sub> by 15%, (n=19)	Reduce rifabutin dose by at least 75% (to a maximum 150 mg every other day or three times per week) when given with saquinavir/ritonavir combination. Monitor closely for adverse



						reactions.
Rilpivirine	25 mg once a day	300 mg once a day	Healthy subjects (18)	ND	↓ AUC by 42%, ↓ C <sub>min</sub> by 48%, ↓ C <sub>max</sub> by 31%	Co-administration of rifabutin with Odefsey (rilpivirine/ tenofovir alafenamide/ emtricitabine) is not recommended, due to an expected decrease in tenofovir alafenamide in addition to the reported reduction in rilpivirine. Refer to Odefsey prescribing information for additional information. Co-administration of rifabutin with cabotegravir/rilpivirine prolonged-release injectable suspension is contraindicated.
Ritonavir	500 mg twice a day for 10 days	150 mg once a day for 16 days	Healthy subjects (5)	↑ AUC by 300%, ↑ C <sub>max</sub> by 150%	ND	Reduce rifabutin dose by at least 75% (to a maximum 150 mg every other day or three times per week) when given with lopinavir/ritonavir combination. Monitor closely for adverse reactions. Reduce rifabutin dosage further, as needed.
Tipranavir/ ritonavir	500/200 twice a day for 15 doses	150 mg single dose	Healthy subjects (20)	↑ AUC by 190%, ↑ C <sub>max</sub> by 70%	↔	Reduce rifabutin dose by at least 75% (to a maximum 150 mg every other day or three times per week) when given with tipranavir/ritonavir combination. Monitor closely for adverse reactions. Reduce rifabutin dosage further, as needed.
Nelfinavir	1250 mg twice a day for 7 to 8 days	150 mg once a day for 8 days	HIV-infected patients (11)	↑ AUC by 83%, <sup>e</sup> ↑ C <sub>max</sub> by 19%	↔	Reduce rifabutin dose by 50% (to 150 mg once a day) and increase the nelfinavir dose to 1250 mg twice a day.
Zidovudine	100 or 200 mg every four hours	300 or 450 mg once a day	HIV-infected patients (16)	↔	↓ AUC by 32%, ↓ C <sub>max</sub> by 48%	Because zidovudine levels remained within the therapeutic range during coadministration of rifabutin, dosage adjustments are not necessary.

**ANTI-HCV DRUGS**

Sofosbuvir	400 mg on day 1 and day 21	300 mg daily on day 10 to day 29	Healthy subjects (20)	ND	36% ↓ in C <sub>max</sub> and 24% ↓ AUC	Co-administration of rifabutin with sofosbuvir (alone or in combination) is not recommended.
------------	----------------------------	----------------------------------	-----------------------	----	---	--

**ANTIFUNGALS**

Fluconazole	200 mg once a day for 2 weeks	300 mg once a day for 2 weeks	HIV-infected patients (12)	↑ AUC by 82%, ↑ C <sub>max</sub> by 88%	↔	Monitor for rifabutin associated adverse events. Reduce rifabutin dose or suspend rifabutin use if toxicity is suspected.
-------------	-------------------------------	-------------------------------	----------------------------	--	---	---

Posaconazole	200 mg once a day for 10 days	300 mg once a day for 17 days	Healthy subjects (8)	↑ AUC by 72%, ↑ C <sub>max</sub> by 31%	↓ AUC by 49%, ↓ C <sub>max</sub> by 43%	If co-administration of these two drugs cannot be avoided, patients should be monitored for adverse events associated with rifabutin administration, and lack of posaconazole efficacy.
--------------	-------------------------------	-------------------------------	----------------------	--	--	---

Itraconazole	200 mg once a day	300 mg once a day	HIV-Infected patients (6)	↑ <sup>f</sup>	↓ AUC by 70%, ↓ C <sub>max</sub> by 75%	If co-administration of these two drugs cannot be avoided, patients should be monitored for adverse events associated with rifabutin administration, and lack of itraconazole efficacy. In a separate study, one case of uveitis was associated with increased serum rifabutin levels following co administration of rifabutin (300 mg once a day) with itraconazole (600 to 900 mg once a day).
--------------	-------------------	-------------------	---------------------------	----------------	--	--

Voriconazole	400 mg twice a day for 7 days (maintenance dose)	300 mg once a day for 7 days	Healthy male subjects (12)	↑ AUC by 331%, ↑ C <sub>max</sub> by 195%	↑ AUC by ~100%, ↑ C <sub>max</sub> by ~100% <sup>9</sup>	<b>CONTRAINDICATED</b>
--------------	--	------------------------------	----------------------------	--	---	------------------------

**ANTI-PCP (Pneumocystis carinii pneumonia)**

Dapsone	50 mg once a day	300 mg once a day	HIV-infected patients (16)	ND	↓ AUC by 27 to 40%	
---------	------------------	-------------------	----------------------------	----	--------------------	--

Sulfamethoxazole-Trimethoprim	800/160 mg	300 mg once a day	HIV-infected patients (12)	↔	↓ AUC by 15 to 20%	
-------------------------------	------------	-------------------	----------------------------	---	--------------------	--

**ANTI-MAC (Mycobacterium avium intracellulare complex)**

Azithromycin	500 mg once a day for 1	300 mg once a	Healthy subjects (6)	↔	↔	
--------------	-------------------------	---------------	----------------------	---	---	--

	day, then 250 mg once a day for 9 days	day				
Clarithromycin	500 mg twice a day	300 mg once a day	HIV-infected patients (12)	↑ AUC by 75%	↓ AUC by 50%	Monitor for rifabutin associated adverse events. Reduce dose or suspend use of rifabutin if toxicity is suspected. Alternative treatment for clarithromycin should be considered when treating patients receiving rifabutin
<b>ANTI-TB (Tuberculosis)</b>						
Ethambutol	1200 mg	300 mg once a day for 7 days	Healthy subjects (10)	ND	↔	
Isoniazid	300 mg	300 mg once a day for 7 days	Healthy subjects (6)	ND	↔	
Bedaquiline	400 mg daily on day 1 and day 29	300 mg daily	Healthy subjects (17)	ND	No change in bedaquiline Pharmacokinetics. 1.4-fold ↑ in M2 and approximately 3.0-fold ↑ in M3 metabolites of bedaquiline.	Avoid bedaquiline co-administration with rifabutin due to the adverse reactions associated with increased bedaquiline Metabolite concentrations.
<b>OTHER</b>						
Methadone	20 to 100 mg once a day	300 mg once a day for 13 days	HIV-infected patients (24)	ND	↔	
Ethinylestradiol (EE)/ Norethindrone (NE)	35 mg EE / 1 mg NE X 21 days	300 mg once a day for 10 days	Healthy female subjects (22)	ND	EE: ↓ AUC by 35%, ↓ C <sub>max</sub> by 20% NE: ↓ AUC by 46%	Patients should be advised to use additional or alternative methods of contraception.
Theophylline	5 mg/kg	300 mg for 14 days	Healthy subjects (11)	ND	↔	

↑ indicates increase; ↓ indicates decrease; ↔ indicates no significant change

ND -No Data

AUC -Area under the Concentration vs. Time Curve; C<sub>max</sub> -Maximum serum concentration; C<sub>min</sub> -Minimum serum concentration

<sup>a</sup> compared to rifabutin 300 mg once a day alone

<sup>b</sup> compared to historical control (fosamprenavir/ritonavir 700/100 mg twice a day)

<sup>c</sup> also taking zidovudine 500 mg once a day

<sup>d</sup> compared to rifabutin 150 mg once a day alone

<sup>e</sup> compared to rifabutin 300 mg once a day alone

<sup>f</sup> data from a case report

<sup>g</sup> compared to voriconazole 200 mg twice a day alone

### Other drugs

The structurally similar drug, rifampin, is known to reduce the plasma concentrations of a number of other drugs (see prescribing information for rifampin). Although a weaker

enzyme inducer than rifampin, rifabutin may be expected to have some effect on those drugs as well.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term carcinogenicity studies were conducted with rifabutin in mice and in rats. Rifabutin was not carcinogenic in mice at doses up to 180 mg/kg/day, or approximately 36 times the recommended human daily dose. Rifabutin was not carcinogenic in the rat at doses up to 60 mg/kg/day, about 12 times the recommended human dose.

Rifabutin was not mutagenic in the bacterial mutation assay (Ames Test) using both rifabutin-susceptible and resistant strains. Rifabutin was not mutagenic in *Schizosaccharomyces pombe* P<sub>1</sub> and was not genotoxic in V-79 Chinese hamster cells, human lymphocytes *in vitro*, or mouse bone marrow cells *in vivo*.

Fertility was impaired in male rats given 160 mg/kg (32 times the recommended human daily dose).

### **Pregnancy**

Rifabutin should be used in pregnant women only if the potential benefit justifies the potential risk to the fetus. There are no adequate and well-controlled studies in pregnant or breastfeeding women.

Reproduction studies have been carried out in rats and rabbits given rifabutin using dose levels up to 200 mg/kg (about 6 to 13 times the recommended human daily dose based on body surface area comparisons). No teratogenicity was observed in either species. In rats, given 200 mg/kg/day, (about 6 times the recommended human daily dose based on body surface area comparisons), there was a decrease in fetal viability. In rats, at 40 mg/kg/day (approximately equivalent to the recommended human daily dose based on body surface area comparisons), rifabutin caused an increase in fetal skeletal variants. In rabbits, at 80 mg/kg/day (about 5 times the recommended human daily dose based on body surface area comparisons), rifabutin caused maternotoxicity and increase in fetal skeletal anomalies. Because animal reproduction studies are not always predictive of human response, rifabutin should be used in pregnant women only if the potential benefit justifies the potential risk to the fetus.

### **Nursing Mothers**

It is not known whether rifabutin is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

### **Pediatric Use**

Safety and effectiveness of rifabutin for prophylaxis of MAC in children have not been established. Limited safety data are available from treatment use in 22 HIV-positive children with MAC who received rifabutin in combination with at least two other antimycobacterials for periods from 1 to 183 weeks. Mean doses (mg/kg) for these children were: 18.5 (range 15.0 to 25.0) for infants 1 year of age, 8.6 (range 4.4 to 18.8) for children 2 to 10 years of age, and 4.0 (range 2.8 to 5.4) for adolescents 14 to 16 years of age. There is no evidence that doses greater than 5 mg/kg daily are useful. Adverse experiences were similar to those observed in the adult population, and included leukopenia, neutropenia and rash. In addition, corneal deposits have been observed in some patients during routine ophthalmologic surveillance of HIV-positive pediatric patients receiving rifabutin as part of a multiple-drug regimen for MAC prophylaxis. These are tiny, almost transparent, asymptomatic peripheral and central corneal deposits which do not impair vision.

### **Geriatric Use**

Clinical studies of rifabutin did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the

elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (see *Clinical Pharmacology*).

## ADVERSE REACTIONS

### Adverse Reactions from Clinical Trials

Rifabutin capsules were generally well tolerated in the controlled clinical trials. Discontinuation of therapy due to an adverse event was required in 16% of patients receiving rifabutin, compared to 8% of patients receiving placebo in these trials. Primary reasons for discontinuation of rifabutin were rash (4% of treated patients), gastrointestinal intolerance (3%), and neutropenia (2%).

The following table enumerates adverse experiences that occurred at a frequency of 1% or greater, among the patients treated with rifabutin in studies 023 and 027.

**Table 3: Clinical Adverse Experiences Reported in ≥ 1% of Patients Treated With Rifabutin**

Adverse event	Rifabutin (N = 566) %	Placebo (N = 580) %
<b>Body as a whole</b>		
Abdominal pain	4	3
Asthenia	1	1
Chest pain	1	1
Fever	2	1
Headache	3	5
Pain	1	2
<b>Blood and lymphatic system</b>		
Leucopenia	10	7
Anemia	1	2
<b>Digestive system</b>		
Anorexia	2	2
Diarrhea	3	3
Dyspepsia	3	1
Eructation	3	1
Flatulence	2	1
Nausea	6	5
Nausea and vomiting	3	2
Vomiting	1	1
<b>Musculoskeletal system</b>		
Myalgia	2	1
<b>Nervous system</b>		
Insomnia	1	1
<b>Skin and appendages</b>		
Rash	11	8
<b>Special senses</b>		
Taste perversion	3	1
<b>Urogenital system</b>		
Discolored urine	30	6

### Clinical Adverse Events Reported In <1% of Patients Who Received Rifabutin

Considering data from the 023 and 027 pivotal trials, and from other clinical studies, rifabutin appears to be a likely cause of the following adverse events which occurred in less than 1% of treated patients: flu-like syndrome, hepatitis, hemolysis, arthralgia,

myositis, chest pressure or pain with dyspnea, skin discoloration, thrombocytopenia, pancytopenia and jaundice.

The following adverse events have occurred in more than one patient receiving rifabutin, but an etiologic role has not been established: seizure, paresthesia, aphasia, confusion, and non-specific T wave changes on electrocardiogram.

When rifabutin was administered at doses from 1050 mg/day to 2400 mg/day, generalized arthralgia and uveitis were reported. These adverse experiences abated when rifabutin was discontinued.

Mild to severe, reversible uveitis has been reported less frequently when rifabutin is used at 300 mg as monotherapy in MAC prophylaxis versus rifabutin in combination with clarithromycin for MAC treatment (*see also Warnings*).

Uveitis has been infrequently reported when rifabutin is used at 300 mg/day as monotherapy in MAC prophylaxis of HIV-infected persons, even with the concomitant use of fluconazole and/or macrolide antibacterials. However, if higher doses of rifabutin are administered in combination with these agents, the incidence of uveitis is higher.

Patients who developed uveitis had mild to severe symptoms that resolved after treatment with corticosteroids and/or mydriatic eye drops; in some severe cases, however, resolution of symptoms occurred after several weeks.

When uveitis occurs, temporary discontinuance of rifabutin and ophthalmologic evaluation are recommended. In most mild cases, rifabutin may be restarted; however, if signs or symptoms recur, use of rifabutin should be discontinued (Morbidity and Mortality Weekly Report, September 9, 1994).

Corneal deposits have been reported during routine ophthalmologic surveillance of some HIV-positive pediatric patients receiving rifabutin as part of a multiple drug regimen for MAC prophylaxis. The deposits are tiny, almost transparent, asymptomatic peripheral and central corneal deposits, and do not impair vision.

The following table enumerates the changes in laboratory values that were considered as laboratory abnormalities in Studies 023 and 027.

**Table 4: Percentage of Patients With Laboratory Abnormalities**

Laboratory abnormalities	Rifabutin (n = 566) %	Placebo (n = 580) %
<b>Chemistry</b>		
Increased alkaline phosphatase <sup>1</sup>	<1	3
Increased SGOT <sup>2</sup>	7	12
Increased SGPT <sup>2</sup>	9	11
<b>Hematology</b>		
Anemia <sup>3</sup>	6	7
Eosinophilia	1	1
Leukopenia <sup>4</sup>	17	16
Neutropenia <sup>5</sup>	25	20
Thrombocytopenia <sup>6</sup>	5	4

Includes grades 3 or 4 toxicities as specified:

<sup>1</sup> All values >450 U/L

<sup>2</sup> All values >150 U/L

<sup>3</sup> All hemoglobin values < 8.0 g/dL

<sup>4</sup> All WBC values < 1,500/mm

<sup>5</sup> All ANC values < 750/mm

<sup>6</sup> All platelet count values < 50,000/mm

The incidence of neutropenia in patients treated with rifabutin was significantly greater than in patients treated with placebo (p = 0.03). Although thrombocytopenia was not significantly more common among patients treated with rifabutin in these trials, rifabutin has been clearly linked to thrombocytopenia in rare cases. One patient in Study 023 developed thrombotic thrombocytopenic purpura, which was attributed to rifabutin.

## Adverse Reactions from Post-Marketing Experience

Adverse reactions identified through post-marketing surveillance by system organ class (SOC) are listed below:

### **Blood and lymphatic system disorders**

White blood cell disorders (including agranulocytosis, lymphopenia, granulocytopenia, neutropenia, white blood cell count decreased, neutrophil count decreased), platelet count decreased.

### **Immune system disorders**

Hypersensitivity, bronchospasm, rash, and eosinophilia.

### **Gastrointestinal disorders**

*Clostridioides difficile* colitis/ *Clostridioides difficile* associated diarrhea.

Pyrexia, rash and other hypersensitivity reactions such as eosinophilia and bronchospasm might occur, as has been seen with other antibacterials.

A limited occurrence of skin discoloration has been reported.

### **Severe cutaneous adverse reactions (SCARs):**

Rifabutin has been associated with the occurrence of DRESS as well as other SCARs such as SJS, TEN, and AGEP (see *WARNINGS*).

### **Rifamycin hypersensitivity reactions**

Hypersensitivity to rifamycins have been reported including flu-like symptoms, bronchospasm, hypotension, urticaria, angioedema, conjunctivitis, thrombocytopenia or neutropenia.

**To report SUSPECTED ADVERSE REACTIONS, contact Lupin Pharmaceuticals, Inc. at 1-800-399-2561 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **ANIMAL TOXICOLOGY**

Liver abnormalities (increased bilirubin and liver weight) occurred in mice, rats and monkeys at doses (respectively) 0.5, 1, and 3 times the recommended human daily dose based on body surface area comparisons. Testicular atrophy occurred in baboons at doses 2 times the recommended human dose based on body surface area comparisons, and in rats at doses 6 times the recommended human daily dose based on body surface area comparisons.

## **OVERDOSAGE**

No information is available on accidental overdosage in humans.

## **Treatment**

While there is no experience in the treatment of overdose with rifabutin capsules, clinical experience with rifamycins suggests that gastric lavage to evacuate gastric contents (within a few hours of overdose), followed by instillation of an activated charcoal slurry into the stomach, may help absorb any remaining drug from the gastrointestinal tract.

Rifabutin is 85% protein bound and distributed extensively into tissues (V<sub>ss</sub>:8 to 9 L/kg). It is not primarily excreted via the urinary route (less than 10% as unchanged drug); therefore, neither hemodialysis nor forced diuresis is expected to enhance the systemic elimination of unchanged rifabutin from the body in a patient with an overdose of rifabutin.

## **DOSAGE AND ADMINISTRATION**

It is recommended that rifabutin capsules USP be administered at a dose of 300 mg

once daily. For those patients with propensity to nausea, vomiting, or other gastrointestinal upset, administration of rifabutin at doses of 150 mg twice daily taken with food may be useful. Doses of rifabutin may be administered mixed with foods such as applesauce.

For patients with severe renal impairment (creatinine clearance less than 30 mL/min), consider reducing the dose of rifabutin by 50%, if toxicity is suspected. No dosage adjustment is required for patients with mild to moderate renal impairment. Reduction of the dose of rifabutin may also be needed for patients receiving concomitant treatment with certain other drugs (*see Precautions-Drug Interactions*).

Mild hepatic impairment does not require a dose modification. The pharmacokinetics of rifabutin in patients with moderate and severe hepatic impairment is not known.

## **HOW SUPPLIED**

Rifabutin Capsules USP are supplied as size '0' capsules having opaque red-brown cap imprinted with 'LU' in white ink and opaque red-brown body imprinted with 'R01' in white ink, containing red-violet granular powder equivalent to 150 mg of rifabutin USP.

Rifabutin Capsules USP are available as follows:

Bottles of 60 capsules - NDC 68180-285-07

Bottles of 100 capsules - NDC 68180-285-01

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Keep tightly closed and dispense in a tight container as defined in the USP. Protect from light and from excessive heat.

Manufactured for:

### **Lupin Pharmaceuticals, Inc.**

Baltimore, Maryland 21202

United States

Manufactured by:

### **Lupin Limited**

Aurangabad 431 210

INDIA

Revised: December 2024

ID#: 275817

Rifabutin Capsules USP, 150 mg

NDC 68180-285-01

A Bottle Label containing 100 capsules



**NDC 68180-285-01**

**Rifabutin Capsules USP**

**150 mg**

Each capsule contains rifabutin USP 150 mg.

**Rx only**

**LUPIN** 100 Capsules

**Usual Dosage:** See accompanying prescribing information. Dispense in tight containers (See USP).  
Keep tightly closed.


Protect from light and from excessive heat.

**Storage:** Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Manufactured for:  
**Lupin Pharmaceuticals, Inc.**  
Baltimore, Maryland 21202, United States

Manufactured by:  
**Lupin Limited**  
Aurangabad 431 210 INDIA  
Code No. MH/DRUGS/499

52 x 20 mm



**RIFABUTIN**

rifabutin capsule

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68180-285
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>RIFABUTIN</b> (UNII: 1W306TDA6S) (RIFABUTIN - UNII:1W306TDA6S)	RIFABUTIN	150 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CROSPROVIDONE (35 .MU.M)</b> (UNII: 40UAA97IT9)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE 101</b> (UNII: 7T9FYH5QMK)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>SHELLAC</b> (UNII: 46N107B710)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	BROWN (Opaque red-brown cap) , BROWN (Opaque red-brown body)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	LU;R01
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68180-285-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/26/2014	03/31/2021
2	NDC:68180-285-07	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2040	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090033	03/26/2014	

**Labeler** - Lupin Pharmaceuticals, Inc. (089153071)

**Registrant** - LUPIN LIMITED (675923163)

## Establishment

Name	Address	ID/FEI	Business Operations
LUPIN LIMITED		862272739	MANUFACTURE(68180-285)

Revised: 12/2024

Lupin Pharmaceuticals, Inc.