

AMOXICILLIN- amoxicillin tablet, film coated
AMOXICILLIN- amoxicillin powder, for suspension
AMOXICILLIN- amoxicillin tablet, chewable
AMOXICILLIN- amoxicillin capsule
Teva Pharmaceuticals USA, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use AMOXICILLIN TABLETS, AMOXICILLIN FOR ORAL SUSPENSION, AMOXICILLIN TABLETS (CHEWABLE), and AMOXICILLIN CAPSULES, safely and effectively. See full prescribing information for AMOXICILLIN TABLETS, AMOXICILLIN FOR ORAL SUSPENSION, AMOXICILLIN TABLETS (CHEWABLE), and AMOXICILLIN CAPSULES.

AMOXICILLIN tablets, for oral use
AMOXICILLIN for oral suspension,
AMOXICILLIN tablets (chewable), for oral use
AMOXICILLIN capsules, for oral use
Initial U.S. Approval: 1974

RECENT MAJOR CHANGES

Warnings and Precautions, Drug-Induced Enterocolitis Syndrome (DIES) (5.3)

5/2024

INDICATIONS AND USAGE

Amoxicillin is a penicillin-class antibacterial indicated for treatment of infections due to susceptible strains of designated microorganisms. (1)

Adults and Pediatric Patients (1)

- Upper Respiratory Tract Infections of the Ear, Nose, and Throat
- Infections of the Genitourinary Tract
- Infections of the Skin and Skin Structure
- Infections of the Lower Respiratory Tract

Adult Patients only (1)

- *Helicobacter pylori* Infection and Duodenal Ulcer Disease

Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules and other antibacterial drugs, amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria. (1)

DOSAGE AND ADMINISTRATION

- In Adults, 750 to 1750 mg/day in divided doses every 8 to 12 hours.
- In Pediatric Patients over 3 Months of Age, 20 to 45 mg/kg/day in divided doses every 8 to 12 hours. Refer to full prescribing information for specific dosing regimens. (2.2, 2.3)
- The upper dose for neonates and infants aged 3 months or younger is 30 mg/kg/day divided every 12 hours. (2.3)
- Dosing for *H. pylori* Infection (in Adults): Triple therapy: 1 gram amoxicillin, 500 mg clarithromycin, and 30 mg lansoprazole, all given twice daily (every 12 hours) for 14 days. Dual therapy: 1 gram amoxicillin and 30 mg lansoprazole, each given three times daily (every 8 hours) for 14 days. (2.4)
- Reduce the dose in patients with severe renal impairment (GFR greater than 30 mL/min). (2.5)

DOSAGE FORMS AND STRENGTHS

- Tablets: 500 mg, 875 mg (3)
- For Oral Suspension: 200 mg/5 mL, 250 mg/5 mL, 400 mg/5 mL (3)
- Tablets (Chewable): 125 mg, 250 mg (3)
- Capsules: 250 mg, 500 mg (3)

CONTRAINDICATIONS

- History of a serious hypersensitivity reaction (e.g., anaphylaxis or Stevens-Johnson syndrome) to amoxicillin or to other beta-lactams (e.g., penicillins or cephalosporins). (4)

WARNINGS AND PRECAUTIONS

- Anaphylactic reactions: Serious and occasionally fatal anaphylactic reactions have been reported in patients on penicillin therapy, including amoxicillin. Discontinue amoxicillin if a reaction occurs (5.1).
- Severe cutaneous adverse reactions (SCAR): Monitor closely. Discontinue if rash progresses. (5.2)
- Drug-induced enterocolitis syndrome (DIES) has been reported with amoxicillin use. If this occurs,

discontinue amoxicillin and institute appropriate therapy. (5.3)

- *Clostridioides difficile*-associated diarrhea (CDAD) (ranging from mild diarrhea to fatal colitis): Evaluate if diarrhea occurs. (5.4)

ADVERSE REACTIONS

The most common adverse reactions (greater than 1%) observed in clinical trials of amoxicillin capsules, amoxicillin tablets or for oral suspension were diarrhea, rash, vomiting, and nausea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Teva at 1-888-838-2872 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Coadministration with probenecid is not recommended. (7.1)
- Concomitant use of amoxicillin and oral anticoagulants may increase the prolongation of prothrombin time. (7.2)
- Coadministration with allopurinol increases the risk of rash. (7.3)
- Amoxicillin may reduce the efficacy of oral contraceptives. (7.4)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 5/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Administration Instructions
- 2.2 Dosage for Adults and Pediatric Patients Aged 3 Months (12 weeks) and Older
- 2.3 Dosage in Pediatric Patients Aged Less than 12 Weeks (3 months)
- 2.4 Dosage for *H. pylori* Infection in Adults
- 2.5 Dosage in Renal Impairment for Adults and Pediatric Patients Aged 3 Months and Older and Weight Greater than 40 kg
- 2.6 Directions for Mixing Oral Suspension

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Anaphylactic Reactions
- 5.2 Severe Cutaneous Adverse Reactions
- 5.3 Drug-Induced Enterocolitis Syndrome (DIES)
- 5.4 *Clostridioides difficile*-Associated Diarrhea (CDAD)
- 5.5 Development of Drug-Resistant Bacteria
- 5.6 Skin Rash in Patients with Mononucleosis

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Probenecid
- 7.2 Oral Anticoagulants
- 7.3 Allopurinol
- 7.4 Oral Contraceptives
- 7.5 Other Antibacterials
- 7.6 Effects on Laboratory Tests

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Dosing in Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 *H. pylori* Eradication to Reduce the Risk of Duodenal Ulcer Recurrence

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Adults and Pediatric Patients

- **Upper Respiratory Tract Infections of the Ear, Nose, and Throat:** Amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules are indicated in the treatment of infections due to susceptible (ONLY β -lactamase-negative) isolates of *Streptococcus* species. (α - and β -hemolytic isolates only), *Streptococcus pneumoniae*, *Staphylococcus* spp., or *Haemophilus influenzae*.
- **Infections of the Genitourinary Tract:** Amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules are indicated in the treatment of infections due to susceptible (ONLY β -lactamase-negative) isolates of *Escherichia coli*, *Proteus mirabilis*, or *Enterococcus faecalis*.
- **Infections of the Skin and Skin Structure:** Amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules are indicated in the treatment of infections due to susceptible (ONLY β -lactamase-negative) isolates of *Streptococcus* spp. (α - and β -hemolytic isolates only), *Staphylococcus* spp., or *E. coli*.
- **Infections of the Lower Respiratory Tract:** Amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules are indicated in the treatment of infections due to susceptible (ONLY β -lactamase-negative) isolates of *Streptococcus* spp. (α - and β -hemolytic isolates only), *S. pneumoniae*, *Staphylococcus* spp., or *H. influenzae*.

Adult Patients only

- ***Helicobacter pylori* Infection and Duodenal Ulcer Disease:**

Triple therapy for *Helicobacter pylori* (*H. pylori*) with clarithromycin and lansoprazole:
Amoxicillin, in combination with clarithromycin plus lansoprazole as triple therapy, is indicated for the treatment of patients with *H. pylori* infection and duodenal ulcer disease (active or 1-year history of a duodenal ulcer) to eradicate *H. pylori*. Eradication of *H. pylori* has been shown to reduce the risk of duodenal ulcer recurrence.

Dual therapy for *H. pylori* with lansoprazole:

Amoxicillin, in combination with lansoprazole delayed-release capsules as dual therapy, is indicated for the treatment of patients with *H. pylori* infection and duodenal ulcer disease (active or 1-year history of a duodenal ulcer) **who are either**

allergic or intolerant to clarithromycin or in whom resistance to clarithromycin is known or suspected. (See the clarithromycin package insert, MICROBIOLOGY.) Eradication of *H. pylori* has been shown to reduce the risk of duodenal ulcer recurrence.

Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of amoxicillin and other antibacterial drugs, amoxicillin should be used only to treat infections that are proven or strongly suspected to be caused by bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

To minimize the potential for gastrointestinal intolerance, amoxicillin should be taken at the start of a meal.

2.2 Dosage for Adults and Pediatric Patients Aged 3 Months (12 weeks) and Older

- Treatment should be continued for a minimum of 48 to 72 hours beyond the time that the patient becomes asymptomatic, or evidence of bacterial eradication has been obtained.
- It is recommended that there be at least 10 days' treatment for any infection caused by *Streptococcus pyogenes* to prevent the occurrence of acute rheumatic fever.
- In some infections, therapy may be required for several weeks. It may be necessary to continue clinical and/or bacteriological follow-up for several months after cessation of therapy.

Table 1. Dosage Recommendations for Adult and Pediatric Patients Aged 3 Months (12 weeks) and Older

Infection	Severity^a	Recommended Dosage for Adults and Pediatric Patients Aged 3 Months and Older and Weight Greater than 40 kg	Recommended Dosage for Pediatric Patients Aged 3 Months and Older and Weight Less than 40 kg
Ear/Nose/Throat Skin/Skin Structure Genitourinary Tract	Mild/Moderate	500 mg every 12 hours or 250 mg every 8 hours	25 mg/kg/day in divided doses every 12 hours or 20 mg/kg/day in divided doses every 8 hours
	Severe	875 mg every 12 hours or 500 mg every 8 hours	45 mg/kg/day in divided doses every 12 hours or 40 mg/kg/day in divided doses every 8 hours
Lower Respiratory Tract	Mild/Moderate or Severe	875 mg every 12 hours or 500 mg every 8 hours	45 mg/kg/day in divided doses every 12 hours or 40 mg/kg/day in divided doses every 8 hours

^a Dosage for infections caused by bacteria that are intermediate in their susceptibility to amoxicillin should follow the recommendations for severe infections.

2.3 Dosage in Pediatric Patients Aged Less than 12 Weeks (3 months)

- It is recommended that there be at least 10 days' treatment for any infection caused by *Streptococcus pyogenes* to prevent the occurrence of acute rheumatic fever.
- Due to incompletely developed renal function affecting elimination of amoxicillin in this age group, the recommended upper dose of amoxicillin is 30 mg/kg/day divided every 12 hours. There are currently no dosing recommendations for pediatric patients with impaired renal function.
- Treatment should be continued for a minimum of 48 to 72 hours beyond the time that the patient becomes asymptomatic, or evidence of bacterial eradication has been obtained.

2.4 Dosage for *H. pylori* Infection in Adults

Triple therapy: The recommended adult oral dose is 1 gram amoxicillin, 500 mg clarithromycin, and 30 mg lansoprazole, all given twice daily (every 12 hours) for 14 days.

Dual therapy: The recommended adult oral dose is 1 gram amoxicillin and 30 mg lansoprazole, each given three times daily (every 8 hours) for 14 days.

Please refer to clarithromycin and lansoprazole full prescribing information.

2.5 Dosage in Renal Impairment for Adults and Pediatric Patients Aged 3 Months and Older and Weight Greater than 40 kg

- Patients with impaired renal function do not generally require a reduction in dose unless the impairment is severe. Renal impairment patients with a glomerular filtration rate of less than 30 mL/min should *NOT* receive the 875 mg dose. See dosage regimens in patients with severe renal impairment provided in Table 2.

Table 2. Dosing in Patients with Severe Renal Impairment

Patients with Renal Impairment	Dosage Regimen
GFR 10 to 30 mL/min	500 mg or 250 mg every 12 hours, depending on the severity of the infection
GFR less than 10 mL/min	500 mg or 250 mg every 24 hours, depending on severity of the infection
Hemodialysis	500 mg or 250 mg every 24 hours, depending on severity of the infection Administer an additional dose both during and at the end of dialysis

2.6 Directions for Mixing Oral Suspension

Prepare a suspension at time of dispensing as follows: Tap bottle until all powder flows freely. Measure the total amount of water (see Table 3). Add approximately 1/3 of the water to powder. Replace cap and *shake vigorously* to wet powder. Add remaining water. Replace cap and *shake vigorously*.

Table 3. Amount of Water for Mixing For Oral Suspension

Strength	Bottle Size	Total Amount of Water
		Required for Reconstitution
For Oral Suspension 200 mg/5 mL	50 mL	39 mL
	75 mL	57 mL
	100 mL	75 mL
For Oral Suspension 250 mg/5 mL	80 mL	47 mL
	100 mL	60 mL
	150 mL	90 mL
For Oral Suspension 400 mg/5 mL	50 mL	35 mL
	75 mL	51 mL
	100 mL	67 mL

After reconstitution, the required amount of suspension should be placed directly on the child's tongue for swallowing. Alternate means of administration are to add the required amount of suspension to formula, milk, fruit juice, water, ginger ale, or cold drinks. These preparations should then be taken immediately.

SHAKE ORAL SUSPENSION WELL BEFORE USING. Keep bottle tightly closed. Any unused portion of the reconstituted suspension must be discarded after 14 days. Refrigeration is preferable, but not required.

3 DOSAGE FORMS AND STRENGTHS

Amoxicillin:

- **Tablets:**

500 mg: Each tablet contains 500 mg of amoxicillin, USP as the trihydrate. Each film-coated, capsule-shaped, off-white, tablet is debossed with "93" on one side and "2263" on the other side.

875 mg: Each tablet contains 875 mg of amoxicillin, USP as the trihydrate. Each film-coated, capsule-shaped, off-white tablet is scored on one side, debossed with "93" on one side of the score and "2264" on the other side of the score.

- **For Oral Suspension:**

200 mg/5 mL: Each 5 mL of reconstituted pink, fruit gum flavored suspension contains 200 mg amoxicillin, USP as the trihydrate.

250 mg/5 mL: Each 5 mL of reconstituted mixed berry flavored suspension contains 250 mg amoxicillin, USP as the trihydrate.

400 mg/5 mL: Each 5 mL of reconstituted pink, fruit gum flavored suspension contains 400 mg amoxicillin, USP as the trihydrate.

- **Tablets (Chewable):**

125 mg: Each white to off-white, capsule-shaped tablet, debossed 93 on one side and 2267 on the other side and contains 125 mg of amoxicillin, USP as the trihydrate.

250 mg: Each white to off-white, capsule-shaped tablet, debossed 93 (partial bisect between 9 and 3) on one side and 2268 on the other side and contains 250 mg of amoxicillin, USP as the trihydrate.

- **Capsules:**

250 mg: Opaque caramel cap and opaque buff body, hard gelatin capsule. Printed black "TEVA" on cap and "3107" on body portions of the capsule and contain 250 mg amoxicillin, USP as the trihydrate.

500 mg: Opaque buff cap and opaque buff body, hard gelatin capsules. Printed black "TEVA" on cap and "3109" on body portions of the capsules and contain 500 mg amoxicillin, USP as the trihydrate.

4 CONTRAINDICATIONS

Amoxicillin is contraindicated in patients who have experienced a serious hypersensitivity reaction (e.g., anaphylaxis or Stevens-Johnson syndrome) to amoxicillin or to other β -lactam antibacterial drugs (e.g., penicillins and cephalosporins).

5 WARNINGS AND PRECAUTIONS

5.1 Anaphylactic Reactions

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients on penicillin therapy including amoxicillin. Although anaphylaxis is more frequent following parenteral therapy, it has occurred in patients on oral penicillins. These reactions are more likely to occur in individuals with a history of penicillin hypersensitivity and/or a history of sensitivity to multiple allergens. There have been reports of individuals with a history of penicillin hypersensitivity who have experienced severe reactions when treated with cephalosporins. Before initiating therapy with amoxicillin, careful inquiry should be made regarding previous hypersensitivity reactions to penicillins, cephalosporins, or other allergens. If an allergic reaction occurs, amoxicillin should be discontinued, and appropriate therapy instituted.

5.2 Severe Cutaneous Adverse Reactions

Amoxicillin may cause severe cutaneous adverse reactions (SCAR), such as Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), and acute generalized exanthematous pustulosis (AGEP). If patients develop skin rash they should be monitored closely, and amoxicillin discontinued if lesions progress.

5.3 Drug-Induced Enterocolitis Syndrome (DIES)

Drug-induced enterocolitis syndrome (DIES) has been reported with amoxicillin use [see *Adverse Reactions (6.2)*], with most cases occurring in pediatric patients ≤ 18 years of age. DIES is a non-IgE mediated hypersensitivity reaction characterized by protracted vomiting occurring 1 to 4 hours after drug ingestion in the absence of skin or respiratory symptoms. DIES may be associated with pallor, lethargy, hypotension, shock, diarrhea within 24 hours after ingesting amoxicillin, and leukocytosis with neutrophilia. If DIES occurs, discontinue amoxicillin and institute appropriate therapy.

5.4 *Clostridioides difficile*-Associated Diarrhea (CDAD)

Clostridioides difficile-associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including amoxicillin, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin-producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial use. Careful medical history is necessary since CDAD has been reported to occur over 2 months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

5.5 Development of Drug-Resistant Bacteria

Prescribing amoxicillin in the absence of a proven or strongly suspected bacterial infection or prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

5.6 Skin Rash in Patients with Mononucleosis

A high percentage of patients with mononucleosis who receive amoxicillin develop an erythematous skin rash. Thus, amoxicillin should not be administered to patients with mononucleosis.

6 ADVERSE REACTIONS

The following are discussed in more detail in other sections of the labeling:

- Anaphylactic reactions [*see Warnings and Precautions (5.1)*]
- Severe Cutaneous Adverse Reactions [*see Warnings and Precautions (5.2)*]
- Drug-Induced Enterocolitis Syndrome (DIES) [*see Warnings and Precautions (5.3)*]
- *Clostridioides difficile*-Associated Diarrhea (CDAD) [*see Warnings and Precautions (5.4)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The most common adverse reactions (greater than 1%) observed in clinical trials of amoxicillin capsules, amoxicillin tablets or for oral suspension were diarrhea, rash, vomiting, and nausea.

Triple therapy: The most frequently reported adverse events for patients who received triple therapy (amoxicillin/clarithromycin/ lansoprazole) were diarrhea (7%), headache (6%), and taste perversion (5%).

Dual therapy: The most frequently reported adverse events for patients who received double therapy amoxicillin/lansoprazole were diarrhea (8%) and headache (7%). For more information on adverse reactions with clarithromycin or lansoprazole, refer to the Adverse Reactions section of their package inserts.

6.2 Postmarketing Experience

In addition to adverse events reported from clinical trials, the following events have been identified during postmarketing use of penicillins. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to amoxicillin.

- **Infections and Infestations:** Mucocutaneous candidiasis.
- **Gastrointestinal:** Drug-induced enterocolitis syndrome (DIES), black hairy tongue, and hemorrhagic/pseudomembranous colitis. Onset of pseudomembranous colitis symptoms may occur during or after antibacterial treatment [*see Warnings and Precautions (5.4)*].
- **Immune:** Hypersensitivity reactions, anaphylactic/anaphylactoid reactions (including shock), angioedema, serum sickness-like reactions (urticaria or skin rash accompanied by arthritis, arthralgia, myalgia, and frequently fever), hypersensitivity vasculitis [*see Warnings and Precautions (5.1)*].
- **Skin and Appendages:** Rashes, pruritus, urticaria, erythema multiforme, SJS, TEN, DRESS, AGEP, exfoliative dermatitis, and linear IgA bullous dermatosis.

- **Liver:** A moderate rise in AST and/or ALT has been noted, but the significance of this finding is unknown. Hepatic dysfunction including cholestatic jaundice, hepatic cholestasis and acute cytolytic hepatitis have been reported.
- **Renal:** Crystalluria has been reported [see *Overdosage (10)*].
- **Hemic and Lymphatic Systems:** Anemia, including hemolytic anemia, thrombocytopenia, thrombocytopenic purpura, eosinophilia, leukopenia, and agranulocytosis have been reported. These reactions are usually reversible on discontinuation of therapy and are believed to be hypersensitivity phenomena.
- **Central Nervous System:** Reversible hyperactivity, agitation, anxiety, insomnia, confusion, convulsions, behavioral changes, aseptic meningitis, and/or dizziness have been reported.
- **Miscellaneous:** Tooth discoloration (brown, yellow, or gray staining) has been reported. Most reports occurred in pediatric patients. Discoloration was reduced or eliminated with brushing or dental cleaning in most cases.

7 DRUG INTERACTIONS

7.1 Probenecid

Probenecid decreases the renal tubular secretion of amoxicillin. Concurrent use of amoxicillin and probenecid may result in increased and prolonged blood levels of amoxicillin.

7.2 Oral Anticoagulants

Abnormal prolongation of prothrombin time (increased international normalized ratio [INR]) has been reported in patients receiving amoxicillin and oral anticoagulants. Appropriate monitoring should be undertaken when anticoagulants are prescribed concurrently. Adjustments in the dose of oral anticoagulants may be necessary to maintain the desired level of anticoagulation.

7.3 Allopurinol

The concurrent administration of allopurinol and amoxicillin increases the incidence of rashes in patients receiving both drugs as compared to patients receiving amoxicillin alone. It is not known whether this potentiation of rashes is due to allopurinol or the hyperuricemia present in these patients.

7.4 Oral Contraceptives

Amoxicillin may affect the intestinal flora, leading to lower estrogen reabsorption and reduced efficacy of combined oral estrogen/progesterone contraceptives.

7.5 Other Antibacterials

Chloramphenicol, macrolides, sulfonamides, and tetracyclines may interfere with the bactericidal effects of penicillin. This has been demonstrated *in vitro*; however, the clinical significance of this interaction is not well documented.

7.6 Effects on Laboratory Tests

High urine concentrations of ampicillin may result in false-positive reactions when testing for the presence of glucose in urine using CLINITEST[®], Benedict's Solution, or Fehling's Solution. Since this effect may also occur with amoxicillin, it is recommended that glucose tests based on enzymatic glucose oxidase reactions (such as CLINISTIX[®]) be used.

Following administration of ampicillin or amoxicillin to pregnant women, a transient decrease in plasma concentration of total conjugated estriol, estriol-glucuronide,

conjugated estrone, and estradiol has been noted.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Teratogenic Effects: Pregnancy Category B.

Reproduction studies have been performed in mice and rats at doses up to 2000 mg/kg (3 and 6 times the 3 g human dose, based on body surface area). There was no evidence of harm to the fetus due to amoxicillin. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, amoxicillin should be used during pregnancy only if clearly needed.

8.2 Labor and Delivery

Oral ampicillin is poorly absorbed during labor. It is not known whether use of amoxicillin in humans during labor or delivery has immediate or delayed adverse effects on the fetus, prolongs the duration of labor, or increases the likelihood of the necessity for an obstetrical intervention.

8.3 Nursing Mothers

Penicillins have been shown to be excreted in human milk. Amoxicillin use by nursing mothers may lead to sensitization of infants. Caution should be exercised when amoxicillin is administered to a nursing woman.

8.4 Pediatric Use

The safety and effectiveness of amoxicillin for the treatment of upper respiratory tract infections, and infections of the genitourinary tract, skin and skin structure and lower respiratory tract have been established in pediatric patients.

The safety and effectiveness of amoxicillin for the treatment of *H. Pylori* infection have not been established in pediatric patients.

Because of incompletely developed renal function in neonates and young infants, the elimination of amoxicillin may be delayed. Dosing of amoxicillin should be modified in pediatric patients 12 weeks or younger (3 months or younger) [see *Dosage and Administration* (2.3)].

8.5 Geriatric Use

An analysis of clinical studies of amoxicillin was conducted to determine whether subjects aged 65 and over respond differently from younger subjects. These analyses have not identified differences in responses between the elderly and younger patients, but a greater sensitivity of some older individuals cannot be ruled out.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

8.6 Dosing in Renal Impairment

Amoxicillin is primarily eliminated by the kidney and dosage adjustment is usually required in patients with severe renal impairment (GFR less than 30 mL/min). See *Dosing in Renal Impairment* (2.5) for specific recommendations in patients with renal impairment.

10 OVERDOSAGE

In case of overdosage, discontinue amoxicillin, treat symptomatically, and institute supportive measures as required. A prospective study of 51 pediatric patients at a poison-control center suggested that overdosages of less than 250 mg/kg of amoxicillin are not associated with significant clinical symptoms.


Interstitial nephritis resulting in oliguric renal failure has been reported in a small number of patients after overdosage with amoxicillin¹.

Crystalluria, in some cases leading to renal failure, has also been reported after amoxicillin overdosage in adult and pediatric patients. In case of overdosage, adequate fluid intake and diuresis should be maintained to reduce the risk of amoxicillin crystalluria.

Renal impairment appears to be reversible with cessation of drug administration. High blood levels may occur more readily in patients with impaired renal function because of decreased renal clearance of amoxicillin. Amoxicillin may be removed from circulation by hemodialysis.

11 DESCRIPTION

Amoxicillin, USP is a semisynthetic antibacterial (amoxicillin), an analog of ampicillin, with a broad spectrum of bactericidal activity against many Gram-positive and Gram-negative microorganisms. Chemically, it is (2*S*,5*R*,6*R*)-6-[(*R*)-(-)-2-amino-2-(*p*-hydroxyphenyl)acetamido]-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylic acid trihydrate. It may be represented structurally as:



The amoxicillin, USP molecular formula is $C_{16}H_{19}N_3O_5S \cdot 3H_2O$, and the molecular weight is 419.45.

Tablets: Each tablet contains 500 mg or 875 mg of amoxicillin, USP as the trihydrate. Each film-coated tablet contains: colloidal silicon dioxide, crospovidone, D&C Yellow No. 10 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, lactose monohydrate, sodium starch glycolate, titanium dioxide and triacetin.

For Oral Suspension: Each 5 mL of reconstituted suspension contains 200 mg, 250 mg or 400 mg of amoxicillin, USP as the trihydrate. Each 5 mL of the 200 mg reconstituted suspension contains 0.09 mEq (2.11 mg) of sodium. Each 5 mL of the 250 mg reconstituted suspension contains 0.14 mEq (3.24 mg) of sodium. Each 5 mL of the 400 mg reconstituted suspension contains 0.12 mEq (2.69 mg) of sodium.

Inactive ingredients for 200 mg and 400 mg: Colloidal silicon dioxide, FD&C Red No. 40, natural and artificial fruit gum flavor, sodium benzoate, sodium citrate, spray dried mask flavor, sucrose, and xanthan gum.

Inactive ingredients for 250 mg: FD&C Red No. 40, mixed berry flavoring, silicon dioxide,

sodium benzoate, sodium citrate, sucrose, and xanthan gum.

Tablets (Chewable): Each tablet contains 125 mg or 250 mg of amoxicillin, USP as the trihydrate. Inactive ingredients in each 125 mg or 250 mg chewable tablet: cherry flavor, lactose anhydrous, magnesium stearate, mannitol, microcrystalline cellulose, sodium citrate, and sucrose.

Capsules: Each capsule, for oral administration, contains 250 mg or 500 mg amoxicillin, USP as the trihydrate. The capsules contain the following inactive ingredients: magnesium stearate and talc. The capsule shell contains D&C Yellow No. 10, FD&C Red No. 40, gelatin, and titanium dioxide. In addition, the 250 mg contains D&C Red No. 28 and FD&C Blue No. 1. The 500 mg may also contain methylparaben, propylparaben, and sodium lauryl sulfate. The printing ink contains black iron oxide, propylene glycol, shellac, and strong ammonia solution. In addition, the 250 mg may contain D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, and FD&C Red No. 40 Aluminum Lake. The 500 mg may also contain potassium hydroxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Amoxicillin is an antibacterial drug [see *Microbiology (12.4)*].

12.3 Pharmacokinetics

Absorption: Amoxicillin is stable in the presence of gastric acid and is rapidly absorbed after oral administration. The effect of food on the absorption of amoxicillin from the tablets and suspension of amoxicillin has been partially investigated; 400 mg and 875 mg formulations have been studied only when administered at the start of a light meal.

Orally administered doses of 250 mg and 500 mg amoxicillin capsules result in average peak blood levels 1 to 2 hours after administration in the range of 3.5 mcg/mL to 5.0 mcg/mL and 5.5 mcg/mL to 7.5 mcg/mL, respectively.

Mean amoxicillin pharmacokinetic parameters from an open, two-part, single-dose crossover bioequivalence study in 27 adults comparing 875 mg of amoxicillin with 875 mg of AUGMENTIN® showed that the 875 mg tablet of amoxicillin produces an $AUC_{0-\infty}$ of 35.4 ± 8.1 mcg•hr/mL and a C_{max} of 13.8 ± 4.1 mcg/mL. Dosing was at the start of a light meal following an overnight fast.

Orally administered doses of amoxicillin suspension, 125 mg/5 mL and 250 mg/5 mL, result in average peak blood levels 1 to 2 hours after administration in the range of 1.5 mcg/mL to 3.0 mcg/mL and 3.5 mcg/mL to 5.0 mcg/mL, respectively.

Oral administration of single doses of 400 mg chewable tablets and 400 mg/5 mL suspension of amoxicillin to 24 adult volunteers yielded comparable pharmacokinetic data:

Table 4: Mean Pharmacokinetic Parameters of Amoxicillin (400 mg chewable tablets and 400 mg/5 mL suspension) in Healthy Adults

Dose*	$AUC_{0-\infty}$ (mcg•hr/mL) Amoxicillin (\pm S.D.)	C_{max} (mcg/mL) [†] Amoxicillin (\pm S.D.)
400 mg (5 mL of suspension)	17.1 (3.1)	5.92 (1.62)
400 mg (1 chewable tablet)	17.9 (2.4)	5.18 (1.64)

* Administered at the start of a light meal.

[†] Mean values of 24 normal volunteers. Peak concentrations occurred approximately 1 hour after the

dose.

Distribution: Amoxicillin diffuses readily into most body tissues and fluids, with the exception of brain and spinal fluid, except when meninges are inflamed. In blood serum, amoxicillin is approximately 20% protein-bound. Following a 1-gram dose, and utilizing a special skin window technique to determine levels of the antibiotic, it was noted that therapeutic levels were found in the interstitial fluid.

Metabolism and Excretion: The half-life of amoxicillin is 61.3 minutes. Approximately 60% of an orally administered dose of amoxicillin is excreted in the urine within 6 to 8 hours. Detectable serum levels are observed up to 8 hours after an orally administered dose of amoxicillin. Since most of the amoxicillin is excreted unchanged in the urine, its excretion can be delayed by concurrent administration of probenecid [see *Drug Interactions (7.1)*].

12.4 Microbiology

Mechanism of Action

Amoxicillin is similar to penicillin in its bactericidal action against susceptible bacteria during the stage of active multiplication. It acts through the inhibition of cell wall biosynthesis that leads to the death of the bacteria.

Resistance

Resistance to Amoxicillin is mediated primarily through enzymes called beta-lactamases that cleave the beta-lactam ring of amoxicillin, rendering it inactive.

Antimicrobial Activity

Amoxicillin has been shown to be active against most isolates of the following microorganisms, both *in vitro* and in clinical infections [see *Indications and Usage (1)*].

Gram-Positive Bacteria

Enterococcus faecalis

Staphylococcus spp.

Streptococcus pneumoniae

Streptococcus spp. (alpha and beta-hemolytic)

Gram-Negative Bacteria

Escherichia coli

Haemophilus influenzae

Helicobacter pylori

Proteus mirabilis

Susceptibility Testing:

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed to evaluate carcinogenic

potential. Studies to detect mutagenic potential of amoxicillin alone have not been conducted; however, the following information is available from tests on a 4:1 mixture of AUGMENTIN®. AUGMENTIN® was non-mutagenic in the Ames bacterial mutation assay, and the yeast gene conversion assay. AUGMENTIN® was weakly positive in the mouse lymphoma assay, but the trend toward increased mutation frequencies in this assay occurred at doses that were also associated with decreased cell survival. AUGMENTIN® was negative in the mouse micronucleus test and in the dominant lethal assay in mice. Potassium clavulanate alone was tested in the Ames bacterial mutation assay and in the mouse micronucleus test and was negative in each of these assays. In a multi-generation reproduction study in rats, no impairment of fertility or other adverse reproductive effects were seen at doses up to 500 mg/kg (approximately 2 times the 3 g human dose based on body surface area).

14 CLINICAL STUDIES

14.1 *H. pylori* Eradication to Reduce the Risk of Duodenal Ulcer Recurrence

Randomized, double-blind clinical studies performed in the United States in patients with *H. pylori* and duodenal ulcer disease (defined as an active ulcer or history of an ulcer within 1 year) evaluated the efficacy of lansoprazole in combination with amoxicillin capsules and clarithromycin tablets as triple 14-day therapy, or in combination with amoxicillin capsules as dual 14-day therapy, for the eradication of *H. pylori*. Based on the results of these studies, the safety and efficacy of 2 different eradication regimens were established: **Triple therapy:** Amoxicillin 1 gram twice daily/clarithromycin 500 mg twice daily/lansoprazole 30 mg twice daily (see Table 5). **Dual therapy:** Amoxicillin 1 gram three times daily/lansoprazole 30 mg three times daily (see Table 6). All treatments were for 14 days. *H. pylori* eradication was defined as 2 negative tests (culture and histology) at 4 to 6 weeks following the end of treatment. Triple therapy was shown to be more effective than all possible dual therapy combinations. Dual therapy was shown to be more effective than both monotherapies. Eradication of *H. pylori* has been shown to reduce the risk of duodenal ulcer recurrence.

Table 5. *H. pylori* Eradication Rates When Amoxicillin is Administered as Part of a Triple Therapy Regimen

Study	Triple Therapy	Triple Therapy
	Evaluable Analysis ^a [95% Confidence Interval] (number of patients)	Intent-to-Treat Analysis ^b [95% Confidence Interval] (number of patients)
Study 1	92 [80.0 to 97.7] (n equals 48)	86 [73.3 to 93.5] (n equals 55)
Study 2	86 [75.7 to 93.6] (n equals 66)	83 [72.0 to 90.8] (n equals 70)

^a This analysis was based on evaluable patients with confirmed duodenal ulcer (active or within 1 year) and *H. pylori* infection at baseline defined as at least 2 of 3 positive endoscopic tests from CLOtest®, histology, and/or culture. Patients were included in the analysis if they completed the study. Additionally, if patients dropped out of the study due to an adverse event related to the study drug, they were included in the analysis as failures of therapy.

^b Patients were included in the analysis if they had documented *H. pylori* infection at baseline as defined above and had a confirmed duodenal ulcer (active or within 1 year). All dropouts were included as failures of therapy.

Table 6. H. pylori Eradication Rates When Amoxicillin is Administered as Part of a Dual Therapy Regimen

Study	Dual Therapy	Dual Therapy
	Evaluable Analysis ^a [95% Confidence Interval] (number of patients)	Intent-to-Treat Analysis ^b [95% Confidence Interval] (number of patients)
Study 1	77 [62.5 to 87.2] (n equals 51)	70 [56.8 to 81.2] (n equals 60)
Study 2	66 [51.9 to 77.5] (n equals 58)	61 [48.5 to 72.9] (n equals 67)

^a This analysis was based on evaluable patients with confirmed duodenal ulcer (active or within 1 year) and *H. pylori* infection at baseline defined as at least 2 of 3 positive endoscopic tests from CLOtest[®], histology, and/or culture. Patients were included in the analysis if they completed the study. Additionally, if patients dropped out of the study due to an adverse event related to the study drug, they were included in the analysis as failures of therapy.

^b Patients were included in the analysis if they had documented *H. pylori* infection at baseline as defined above and had a confirmed duodenal ulcer (active or within 1 year). All dropouts were included as failures of therapy.

15 REFERENCES

1. Swanson-Biearman B, Dean BS, Lopez G, Krenzelok EP. The effects of penicillin and cephalosporin ingestions in children less than six years of age. *Vet Hum Toxicol.* 1988; 30: 66-67.

16 HOW SUPPLIED/STORAGE AND HANDLING

Amoxicillin tablets, USP:

500 mg: Each tablet contains 500 mg of amoxicillin, USP as the trihydrate. Each film-coated, capsule-shaped, off-white, tablet is debossed with “93” on one side and “2263” on the other side. They are available in bottles of 100 (NDC 0093-2263-01) tablets.

875 mg: Each tablet contains 875 mg of amoxicillin, USP as the trihydrate. Each film-coated, capsule-shaped, off-white tablet is scored on one side, debossed with “93” on one side of the score and “2264” on the other side of the score. They are available in bottles of 100 (NDC 0093-2264-01) tablets.

Amoxicillin for oral suspension, USP:

200 mg/5 mL: Each 5 mL of reconstituted pink, fruit gum flavored suspension contains 200 mg amoxicillin, USP as the trihydrate. It is available in bottles of 50 mL (NDC 0093-4160-76), 75 mL (NDC 0093-4160-78), and 100 mL (NDC 0093-4160-73).

250 mg/5 mL: Each 5 mL of reconstituted mixed berry flavored suspension contains 250 mg amoxicillin, USP as the trihydrate. It is available in bottles of 80 mL (NDC 0093-4155-79), 100 mL (NDC 0093-4155-73), and 150 mL (NDC 0093-4155-80).

400 mg/5 mL: Each 5 mL of reconstituted pink, fruit gum flavored suspension contains 400 mg amoxicillin, USP as the trihydrate. It is available in bottles of 50 mL

(NDC 0093-4161-76), 75 mL (NDC 0093-4161-78), and 100 mL (NDC 0093-4161-73).

Amoxicillin Tablets (Chewable), USP:

125 mg: Each white to off-white, capsule-shaped tablet, debossed 93 on one side and 2267 on the other side and contain 125 mg amoxicillin, USP as the trihydrate. They are available in bottles of 100 tablets (NDC 0093-2267-01).

250 mg: Each white to off-white, capsule-shaped tablet, debossed 93 (partial bisect between 9 and 3) on one side and 2268 on the other side and contain 250 mg amoxicillin, USP as the trihydrate. They are available in bottles of 100 tablets (NDC 0093-2268-01).

Amoxicillin Capsules, USP

250 mg: Opaque caramel cap and opaque buff body, hard gelatin capsule. Printed black "TEVA" on cap and "3107" on body portions of the capsule and contain 250 mg amoxicillin, USP as the trihydrate. They are available in bottles of 100 (NDC 0093-3107-01) and 500 (NDC 0093-3107-05) capsules.

500 mg: Opaque buff cap and opaque buff body, hard gelatin capsules. Printed black "TEVA" on cap and "3109" on body portions of the capsules and contain 500 mg amoxicillin, USP as the trihydrate. They are available in bottles of 50 (NDC 0093-3109-53), 60 (NDC 0093-3109-06), 100 (NDC 0093-3109-01) and 500 (NDC 0093-3109-05) capsules.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Keep this and all medications out of the reach of children.

17 PATIENT COUNSELING INFORMATION

Administration Instructions

Advise patients that amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules may be taken every 8 hours or every 12 hours, depending on the dose prescribed.

Allergic Reactions

Counsel patients that amoxicillin contains a penicillin class drug product that can cause allergic reactions in some individuals.

Severe Cutaneous Adverse Reactions (SCAR)

Advise patients about the signs and symptoms of serious skin manifestations. Instruct patients to stop taking amoxicillin immediately and promptly report the first signs or symptoms of skin rash, mucosal lesions, or any other sign of hypersensitivity [see *Warnings and Precautions (5.2)*].

Diarrhea

Counsel patients that diarrhea is a common problem caused by antibacterial drugs which usually ends when the antibacterial drug is discontinued. Sometimes after starting treatment with antibacterial drugs, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as 2 or more months after having taken their last dose of the antibacterial drug. If this occurs, patients should contact their physician as soon as possible.

Antibacterial Resistance

Patients should be counseled that antibacterial drugs, including amoxicillin, should only

be used to treat or prevent bacterial infections. Antibacterial drugs do not treat viral infections (e.g., the common cold). When amoxicillin tablets, amoxicillin for oral suspension, amoxicillin tablets (chewable), and amoxicillin capsules are prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may: (1) decrease the effectiveness of the immediate treatment, and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by amoxicillin or other antibacterial drugs in the future.

Storage and Special Handling Instructions

It is preferable to refrigerate amoxicillin suspensions, but not required. Shake oral suspensions well before each use. Keep bottle tightly closed. When dosing a child with the suspension (liquid), use a calibrated oral syringe. Be sure to rinse the calibrated oral syringe after each use. Bottles of suspension of amoxicillin may contain more liquid than required. Follow your doctor's instructions about the amount to use and the days of treatment your child requires. Discard any unused portion of the suspension after 14 days.

Brands listed are the trademarks of their respective owners.

Manufactured In Canada By:

Teva Canada Limited

Toronto, Canada M1B 2K9

Manufactured For:

Teva Pharmaceuticals

Parsippany, NJ 07054

Rev. 5/2024

Package Label.Principal Display Panel

NDC 0093-2263-01

Amoxicillin Tablets, USP

500 mg

AMOXICILLIN USP, 500 mg as the trihydrate

Rx only

100 Tablets

NDC 0093-2263-01

Amoxicillin Tablets, USP

500 mg

AMOXICILLIN USP, 500 mg as the trihydrate

Rx only
100 Tablets

teva

GTIN 00300932263017

Each tablet contains 500 mg amoxicillin, USP as the trihydrate.

Dosage: See accompanying prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Use only if inner seal is intact.

Keep this and all medications out of the reach of children.

Manufactured in Canada By:
Teva Canada Limited
Toronto, Canada M1B 2K9

Manufactured For:
Teva Pharmaceuticals
 Parsippany, NJ 07054

361-32- 669786240A Rev 04 Rev. E 12/2022

3 0 0 9 3 - 2 2 6 3 - 0 1 7

Serialization Coding Area

Package Label.Principal Display Panel

NDC 0093-2264-01
Amoxicillin Tablets, USP
875 mg
AMOXICILLIN USP, 875 mg as the trihydrate
Rx only
100 Tablets

NDC 0093-2264-01

Amoxicillin Tablets, USP

875 mg

AMOXICILLIN USP, 875 mg as the trihydrate

Rx only
100 Tablets

teva

GTIN 00300932264014

Each tablet contains 875 mg amoxicillin, USP as the trihydrate.

Dosage: See accompanying prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Use only if inner seal is intact.

Keep this and all medications out of the reach of children.

Manufactured in Canada By:
Teva Canada Limited
Toronto, Canada M1B 2K9

Manufactured For:
Teva Pharmaceuticals
 Parsippany, NJ 07054

361-32- 669796240A Rev 04 Rev. E 12/2022

3 0 0 9 3 - 2 2 6 4 - 0 1 4

Serialization Coding Area

Package Label.Principal Display Panel

NDC 0093-4160-73
Amoxicillin for Oral Suspension, USP
200 mg per 5 mL
When reconstituted, each 5 mL contains: AMOXICILLIN USP, 200 mg as the trihydrate
Rx only
100 mL (when reconstituted)

NDC 0093-4160-73

Amoxicillin for Oral Suspension, USP

200 mg per 5 mL

When reconstituted, each 5 mL contains: AMOXICILLIN USP, 200 mg as the trihydrate

Rx only

100 mL
(when reconstituted)

CTIN 00300934160734

Net contents: Equivalent to 4 grams amoxicillin, USP

Directions for mixing:

1. Tap bottle until all powder flows freely.
2. Measure 75 mL of water (total).
3. Add approximately 1/3 of the water to powder. Replace cap; shake vigorously.
4. Add rest of water. Replace cap; shake vigorously.

Each 5 mL (1 teaspoonful) will contain amoxicillin trihydrate equivalent to 200 mg amoxicillin, USP

Dosage: See accompanying prescribing information.

Use only if inner seal is intact.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Keep this and all medications out of the reach of children.

Keep tightly closed. Shake well before each use. Refrigeration preferable but not required.

Date reconstituted: / /

Discard after 14 days.


Manufactured in Canada By: Teva Canada Limited, Toronto, Canada M1B 2K9

Manufactured For: Teva Pharmaceuticals Parsippany, NJ 07054

0 0 9 3 - 4 1 6 0 - 7 3

361-32- 669816240A Rev 06 Rev. F 6/2023

Serialization Coding Area



Package Label.Principal Display Panel

NDC 0093-4155-73
 Amoxicillin for Oral Suspension, USP
 250 mg per 5 mL
 When reconstituted, each 5 mL contains: AMOXICILLIN USP, 250 mg as the trihydrate
 Rx only
 100 mL (when reconstituted)

NDC 0093-4155-73

Amoxicillin for Oral Suspension, USP

250 mg per 5 mL

When reconstituted, each 5 mL contains: AMOXICILLIN USP, 250 mg as the trihydrate

Rx only

100 mL
(when reconstituted)

CTIN 00300934155730

Net contents: Equivalent to 5 grams amoxicillin, USP

Directions for mixing:

1. Tap bottle until all powder flows freely.
2. Measure 50 mL of water (total).
3. Add approximately 1/3 of the water to powder. Replace cap; shake vigorously.
4. Add rest of water. Replace cap; shake vigorously.

Each 5 mL (1 teaspoonful) will contain amoxicillin trihydrate equivalent to 250 mg amoxicillin, USP

Dosage: See accompanying prescribing information.

Use only if inner seal is intact.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Keep this and all medications out of the reach of children.

Keep tightly closed. Shake well before each use. Refrigeration preferable but not required.

Date reconstituted: / /

Discard after 14 days.


Manufactured in Canada By: Teva Canada Limited, Toronto, Canada M1B 2K9

Manufactured For: Teva Pharmaceuticals Parsippany, NJ 07054

0 0 9 3 - 4 1 5 5 - 7 3 0

361-32- 669656240A Rev 05 Rev. H 12/2022

Serialization Coding Area



Package Label.Principal Display Panel

NDC 0093-4161-73
 Amoxicillin for Oral Suspension, USP
 400 mg per 5 mL
 When reconstituted, each 5 mL contains: AMOXICILLIN USP, 400 mg as the trihydrate
 Rx only

100 mL (when reconstituted)

361-32- 669826240A Rev 06 Rev. F 6/2023

Manufactured in Canada By: Teva Canada Limited Toronto, Canada M1B 2K9
Manufactured For: Teva Pharmaceuticals Parsippany, NJ 07054

0 0 9 3 - 4 1 6 1 - 7 3

3 1

Amoxicillin for Oral Suspension, USP
400 mg per 5 mL

NDC 0093-4161-73
CTIN 00300934161731

Net contents: Equivalent to 8 grams amoxicillin, USP

Directions for mixing:
1. Tap bottle until all powder flows freely.
2. Measure 67 mL of water (total).
3. Add approximately 1/3 of the water to powder. Replace cap; shake vigorously.
4. Add rest of water. Replace cap; shake vigorously.

Each 5 mL (1 teaspoonful) will contain amoxicillin trihydrate equivalent to 400 mg amoxicillin, USP.

Dosage: See accompanying prescribing information.


Use only if inner seal is intact.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
Keep this and all medications out of the reach of children.
Keep tightly closed. Shake well before each use. Refrigeration preferable but not required.

Date reconstituted: _____
Discard after 14 days.

When reconstituted, each 5 mL contains: **AMOXICILLIN USP, 400 mg** as the trihydrate

Rx only
100 mL
(when reconstituted)

Serialization Coding Area



Package Label.Principal Display Panel

NDC 0093-2267-01
Amoxicillin Tablets, USP
(Chewable)
125 mg
AMOXICILLIN USP, 125 mg as the trihydrate
Rx only
100 Tablets

361-32- 669676240A Rev 04 Rev. D 12/2022

Manufactured in Canada By: Teva Canada Limited Toronto, Canada M1B 2K9
Manufactured For: Teva Pharmaceuticals Parsippany, NJ 07054

0 0 9 3 - 2 2 6 7 - 0 1 5

3 1

Amoxicillin Tablets, USP (Chewable)
125 mg

NDC 0093-2267-01
CTIN 00300932267015

Each chewable tablet contains 125 mg amoxicillin, USP as the trihydrate.

Dosage: See accompanying prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
Dispense in a tight light-resistant container as defined in the USP with a child-resistant closure (as required).


Use only if inner seal is intact.
Keep this and all medications out of the reach of children.

Manufactured in Canada By: Teva Canada Limited Toronto, Canada M1B 2K9
Manufactured For: Teva Pharmaceuticals Parsippany, NJ 07054

AMOXICILLIN USP, 125 mg as the trihydrate

Rx only
100 Tablets

Serialization Coding Area



Package Label.Principal Display Panel

NDC 0093-2268-01
Amoxicillin Tablets, USP
(Chewable)

250 mg
AMOXICILLIN USP, 250 mg as the trihydrate
Rx only
100 Tablets

3 0093-2268-01 2

361-32- 669686240A Rev 04 Rev. D 12/2022

Serialization Coding Area

Each chewable tablet contains 250 mg amoxicillin, USP as the trihydrate.

Dosage: See accompanying prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Use only if inner seal is intact. Keep this and all medications out of the reach of children.

Manufactured In Canada By: Teva Canada Limited Toronto, Canada M1B 2K9

Manufactured For: Teva Pharmaceuticals Parsippany, NJ 07054

NDC 0093-2268-01

Amoxicillin Tablets, USP (Chewable)

250 mg

AMOXICILLIN USP, 250 mg as the trihydrate

Rx only
100 Tablets

teva

Package Label.Principal Display Panel

NDC 0093-3107-05
Amoxicillin Capsules, USP
250 mg
Rx only
500 Capsules

3 0093-3107-05 1

361-32- 669666270 Rev 01 Rev. K 12/2022

Serialization Coding Area

Each capsule contains amoxicillin trihydrate equivalent to 250 mg amoxicillin, USP.

Usual Adult Dosage: 1 or 2 capsules every 8 hours.

See accompanying literature.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Keep this and all medications out of the reach of children.

Manufactured In Canada By: Teva Canada Limited Toronto, Canada M1B 2K9

Manufactured For: Teva Pharmaceuticals Parsippany, NJ 07054

NDC 0093-3107-05

Amoxicillin Capsules, USP

250 mg

Rx only
500 Capsules

teva

Package Label.Principal Display Panel

NDC 0093-3109-05
Amoxicillin Capsules, USP

500 mg
 Rx only
 500 Capsules

NDC 0093-3109-05

Amoxicillin Capsules, USP
500 mg

Rx only
 500 Capsules

CTIN 00300933109055

Each capsule contains amoxicillin trihydrate equivalent to 500 mg amoxicillin, USP.

Usual Adult Dosage: 1 capsule every 8 hours.

See accompanying literature.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required). Keep this and all medications out of the reach of children.

Manufactured in Canada By:
 Teva Canada Limited
 Toronto, Canada M1B 2K9

Manufactured For:
 Teva Pharmaceuticals
 Parsippany, NJ 07054

361-32- 669076270 Rev 04 Rev. K 12/2022

Serialization Coding Area

AMOXICILLIN

amoxicillin tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-2263
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII: 9EM05410Q9)	AMOXICILLIN ANHYDROUS	500 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 36SFW2JZ0W)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white (off-white)	Score	no score
Shape	OVAL (capsule-shaped)	Size	17mm
Flavor		Imprint Code	93;2263
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-2263-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2000	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065056	11/21/2000	

AMOXICILLIN

amoxicillin tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-2264
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	875 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 365FW2JZ0W)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white (off-white)	Score	2 pieces
Shape	OVAL (capsule-shaped)	Size	21mm
Flavor		Imprint Code	93;2264
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-2264-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/26/2000	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065056	09/26/2000	

AMOXICILLIN				
amoxicillin powder, for suspension				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-4160	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)		AMOXICILLIN ANHYDROUS	200 mg in 5 mL	
Inactive Ingredients				
Ingredient Name		Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SODIUM CITRATE (UNII: 1Q73Q2JULR)				
SUCROSE (UNII: C151H8M554)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	pink	Score		
Shape		Size		
Flavor	FRUIT (fruit gum)	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-4160-76	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2003	
2	NDC:0093-4160-78	75 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2003	
3	NDC:0093-4160-73	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/02/2003	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065119	04/02/2003	

AMOXICILLIN

amoxicillin powder, for suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-4155
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	250 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-4155-79	80 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/30/1990	
2	NDC:0093-4155-73	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/30/1990	
3	NDC:0093-4155-80	150 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/30/1990	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA061931	09/30/1990	

AMOXICILLIN

amoxicillin powder, for suspension

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-4161
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	400 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	pink	Score	
Shape		Size	
Flavor	FRUIT (fruit gum)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-4161-76	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2003	
2	NDC:0093-4161-78	75 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2003	
3	NDC:0093-4161-73	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/26/2003	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065119	02/26/2003	

AMOXICILLIN

amoxicillin tablet, chewable

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-2267

Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	125 mg	
Inactive Ingredients				
	Ingredient Name	Strength		
	ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
	MAGNESIUM STEARATE (UNII: 70097M6I30)			
	MANNITOL (UNII: 3OWL53L36A)			
	CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
	SODIUM CITRATE (UNII: 1Q73Q2JULR)			
	SUCROSE (UNII: C151H8M554)			
Product Characteristics				
Color	white (white to off-white)	Score	no score	
Shape	OVAL (capsule-shaped)	Size	15mm	
Flavor	CHERRY	Imprint Code	93;2267	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-2267-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/1995	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA064013	10/01/1995		

AMOXICILLIN			
amoxicillin tablet, chewable			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-2268
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	250 mg
Inactive Ingredients			

Ingredient Name		Strength		
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MANNITOL (UNII: 3OWL53L36A)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
SODIUM CITRATE (UNII: 1Q73Q2JULR)				
SUCROSE (UNII: C151H8M554)				
Product Characteristics				
Color	white (white to off-white)	Score	2 pieces	
Shape	OVAL (capsule-shaped)	Size	19mm	
Flavor	CHERRY	Imprint Code	9;3;2268	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-2268-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/1993	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA064013	01/01/1993		

AMOXICILLIN

amoxicillin capsule

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-3107
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	250 mg	
Inactive Ingredients			
Ingredient Name	Strength		
MAGNESIUM STEARATE (UNII: 70097M6I30)			
TALC (UNII: 7SEV7J4R1U)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GELATIN (UNII: 2G86QN327L)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
D&C RED NO. 28 (UNII: 767IP0Y5NH)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FERROSFERRIC OXIDE (UNII: XM0M87F357)			

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SHELLAC (UNII: 46N107B71O)				
AMMONIA (UNII: 5138Q19F1X)				
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)				
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)				
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)				
Product Characteristics				
Color	brown (caramel) , brown (buff)	Score no score		
Shape	CAPSULE	Size 18mm		
Flavor		Imprint Code TEVA;3107		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-3107-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/1990	
2	NDC:0093-3107-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/1990	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA061926	09/30/1990		

AMOXICILLIN			
amoxicillin capsule			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0093-3109
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)		AMOXICILLIN ANHYDROUS	500 mg
Inactive Ingredients			
Ingredient Name			Strength
MAGNESIUM STEARATE (UNII: 70097M6I3O)			
TALC (UNII: 7SEV7J4R1U)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C RED NO. 40 (UNII: WZ B9127XOA)			
GELATIN (UNII: 2G86QN327L)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
METHYLPARABEN (UNII: A2I8C7HI9T)			
PROPYLPARABEN (UNII: Z8IX2SC1OH)			

SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
AMMONIA (UNII: 5138Q19F1X)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	

Product Characteristics

Color	brown (buff)	Score	no score
Shape	CAPSULE	Size	22mm
Flavor		Imprint Code	TEVA;3109
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0093-3109-53	50 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/1990	
2	NDC:0093-3109-06	60 in 1 BOTTLE; Type 0: Not a Combination Product	12/15/2023	
3	NDC:0093-3109-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/15/2023	
4	NDC:0093-3109-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/1990	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA061926	09/30/1990	

Labeler - Teva Pharmaceuticals USA, Inc. (001627975)

Revised: 5/2024

Teva Pharmaceuticals USA, Inc.