

PRED MILD - prednisolone acetate suspension/ drops
Physicians Total Care, Inc.


PRED MILD®
(prednisolone acetate ophthalmic suspension, USP) 0.12%

sterile

DESCRIPTION

PRED MILD® (prednisolone acetate ophthalmic suspension, USP) 0.12% is a topical anti-inflammatory agent for ophthalmic use.

Structural Formula:



prednisolone acetate

Chemical Name:

11β, 17, 21-Trihydroxypregna-1, 4-diene-3, 20-dione 21-acetate

Contains:

Active: prednisolone acetate (microfine suspension) 0.12%. **Preservative:** benzalkonium chloride.
Inactives: boric acid; edetate disodium; hypromellose; polysorbate 80; purified water; sodium bisulfite; sodium chloride; and sodium citrate. The pH during its shelf life ranges from 5.0 - 6.0.

CLINICAL PHARMACOLOGY

Prednisolone acetate is a glucocorticoid that, on the basis of weight, has 3 to 5 times the anti-inflammatory potency of hydrocortisone. Glucocorticoids inhibit the edema, fibrin deposition, capillary dilation, and phagocytic migration of the acute inflammatory response, as well as capillary proliferation, deposition of collagen, and scar formation.

INDICATIONS AND USAGE

PRED MILD® is indicated for the treatment of mild to moderate noninfectious allergic and inflammatory disorders of the lid, conjunctiva, cornea, and sclera (including chemical and thermal burns).

CONTRAINDICATIONS

PRED MILD® suspension is contraindicated in acute untreated purulent ocular infections, in most viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella, and also in mycobacterial infection of the eye and fungal diseases of ocular structures. PRED MILD® suspension is also contraindicated in individuals with known or suspected

hypersensitivity to any of the ingredients of this preparation and to other corticosteroids.

WARNINGS

Prolonged use of corticosteroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and in posterior subcapsular cataract formation. Prolonged use may also suppress the host immune response and thus increase the hazard of secondary ocular infections.

Various ocular diseases and long-term use of topical corticosteroids have been known to cause corneal and scleral thinning. Use of topical corticosteroids in the presence of thin corneal or scleral tissue may lead to perforation.

Acute purulent infections of the eye may be masked or activity enhanced by the presence of corticosteroid medication.

If this product is used for 10 days or longer, intraocular pressure should be routinely monitored even though it may be difficult in children and uncooperative patients. Steroids should be used with caution in the presence of glaucoma. Intraocular pressure should be checked frequently.

The use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation.

Use of ocular steroids may prolong the course and may exacerbate the severity of many viral infections of the eye (including herpes simplex). Employment of a corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution; frequent slit lamp microscopy is recommended.

Corticosteroids are not effective in mustard gas keratitis and Sjögren's keratoconjunctivitis.

Contains sodium bisulfite, a sulfite that may cause allergic-type reactions, including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people.

PRECAUTIONS

General:

The initial prescription and renewal of the medication order beyond 20 milliliters of PRED MILD[®] should be made by a physician only after examination of the patient with the aid of magnification, such as slit lamp biomicroscopy, and, where appropriate, fluorescein staining. If signs and symptoms fail to improve after 2 days, the patient should be re-evaluated.

As fungal infections of the cornea are particularly prone to develop coincidentally with long-term local corticosteroid applications, fungal invasion should be suspected in any persistent corneal ulceration where a corticosteroid has been used or is in use. Fungal cultures should be taken when appropriate.

If this product is used for 10 days or longer, intraocular pressure should be monitored (see **WARNINGS**).

Information for patients :

If inflammation or pain persists longer than 48 hours or becomes aggravated, the patient should be advised to discontinue use of the medication and consult a physician.

This product is sterile when packaged. To prevent contamination, care should be taken to avoid touching the bottle tip to eyelids or to any other surface. The use of this bottle by more than one person may spread infection. Keep bottle tightly closed when not in use. Keep out of the reach of children.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

No studies have been conducted in animals or in humans to evaluate the potential of these effects.

Pregnancy Category C:

Prednisolone has been shown to be teratogenic in mice when given in doses 1-10 times the human dose. There are no adequate well-controlled studies in pregnant women. Prednisolone should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Dexamethasone, hydrocortisone, and prednisolone were ocularly applied to both eyes of pregnant mice five times per day on days 10 through 13 of gestation. A significant increase in the incidence of cleft palate was observed in the fetuses of the treated mice.

Nursing Mothers:

It is not known whether topical ophthalmic administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. Because of the potential for serious adverse reactions in nursing infants from prednisolone, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use:

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use:

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS

Adverse reactions include, in decreasing order of frequency, elevation of intraocular pressure (IOP) with possible development of glaucoma and infrequent optic nerve damage, posterior subcapsular cataract formation, and delayed wound healing.

Although systemic effects are extremely uncommon, there have been rare occurrences of systemic hypercorticism after use of topical steroids.

Corticosteroid-containing preparations have also been reported to cause acute anterior uveitis and perforation of the globe. Keratitis, conjunctivitis, corneal ulcers, mydriasis, conjunctival hyperemia, loss of accommodation and ptosis have occasionally been reported following local use of corticosteroids.

The development of secondary ocular infection (bacterial, fungal, and viral) has occurred. Fungal and viral infections of the cornea are particularly prone to develop coincidentally with long-term applications of steroids. The possibility of fungal invasion should be considered in any persistent corneal ulceration where steroid treatment has been used (see **WARNINGS**).

Transient burning and stinging upon instillation and other minor symptoms of ocular irritation have been reported with the use of PRED MILD[®] suspension.

OVERDOSAGE

Overdosage will not ordinarily cause acute problems. If accidentally ingested, drink fluids to dilute.

DOSAGE AND ADMINISTRATION

Shake well before using. Instill one to two drops into the conjunctival sac two to four times daily. During the initial 24 to 48 hours, the dosing frequency may be increased if necessary. Care should be taken not to discontinue therapy prematurely.

If signs and symptoms fail to improve after 2 days, the patient should be re-evaluated (see **PRECAUTIONS**).

HOW SUPPLIED

PRED MILD® (prednisolone acetate ophthalmic suspension, USP) 0.12% is supplied sterile in opaque white LDPE plastic bottles with droppers with white high impact polystyrene (HIPS) caps as follows:

5 mL in 10 mL bottle - NDC 54868-6355-0

Note: Store at 15° - 30°C (59° - 86°F) in an upright position. Protect from freezing.

Rx Only

Revised June 2004

© 2004 Allergan, Inc.

Irvine, CA 92612, U.S.A.

® Marks owned by Allergan, Inc.

6295X

71739US10P

Distributed by:

Physicians Total Care, Inc.

Tulsa, Oklahoma 74146



NDC 54868-6355-0

Rx Only

PRED MILD®

(prednisolone

acetate ophthalmic

suspension, USP)

0.12%

sterile

5 mL

PRED MILD

prednisolone acetate suspension/ drops

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:54868-6355(NDC:11980-174)
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
prednisolone acetate (UNII: 8B2807733D) (prednisolone - UNII:9PHQ9Y1OLM)	prednisolone acetate	1.2 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
benzalkonium chloride (UNII: F5UM2KM3W7)	
boric acid (UNII: R57ZHV85D4)	
edetate disodium (UNII: 7FLD91C86K)	
hypromelloses (UNII: 3NXW29V3WO)	
polysorbate 80 (UNII: 6OZP39ZG8H)	
water (UNII: 059QF0KO0R)	
sodium bisulfite (UNII: TZX5469Z6I)	
sodium chloride (UNII: 451W47IQ8X)	
sodium citrate (UNII: 1Q73Q2JULR)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54868-6355-0	1 in 1 CARTON		
1		5 mL in 1 BOTTLE, DROPPER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017100	11/01/2012	

Labeler - Physicians Total Care, Inc. (194123980)

Establishment

Name	Address	ID/FEI	Business Operations
Physicians Total Care, Inc.		194123980	relabel(54868-6355)

