

LEADER COUGH DM- dextromethorphan polistirex suspension
Cardinal Health 110, LLC. dba Leader

Cardinal Health Cough DM Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **shake bottle well before use**
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

- **each 5 mL contains: sodium 5 mg**
- store at 20° to 25° C (68° to 77° F)
- dosing cup provided

Inactive ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Day or Night

Cough DM

Dextromethorphan Polistirex

Extended-Release Oral Suspension

Cough Suppressant

Orange Flavored Liquid

COMPARE TO DELSYM® active ingredient

12 Hour Cough Relief

Alcohol-Free

Dosing Cup Included

100% Money Back Guarantee

Contains Sodium Metabisulfite, a Sulfite That May Cause Allergic-Type Reactions

3 FL OZ (89 mL)

DO NOT USE IF CARTON IS OPENED OR PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Day or Night Cough DM
Dextromethorphan Polistirex
Extended-Release Oral Suspension
Cough Suppressant

LEADER

Day or Night Cough DM
Dextromethorphan Polistirex
Extended-Release Oral Suspension
Cough Suppressant

Orange Flavored Liquid
12 Hour Cough Relief
Alcohol-Free
Dosing Cup Included

COMPARE TO DELSYM® active ingredient*

100% Money Back Guarantee

3 FL OZ (89 mL)

Orange Flavored Liquid
12 Hour Cough Relief
Alcohol-Free
Dosing Cup Included
Contains No Fever Reducer or Pain Reliever

100% Money Back Guarantee

3 FL OZ (89 mL)

DO NOT USE IF CARTON IS OPENED OR PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient (in each 5 mL) Purpose
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant

Uses temporarily relieves
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have
■ chronic cough that lasts as occurs with smoking, asthma or emphysema
■ cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if
■ side effects occur: You may report side effects to FDA at 1-800-FDA-1088.
■ cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ shake bottle well before use
■ measure only with dosing cup provided
■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information
■ each 5 mL contains: sodium 5 mg
■ store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

Drug Facts (continued)
Inactive ingredients D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?
1-800-719-9260

DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided. Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

CIN 5297841 REV. 6/20

0 96295 13035 5

Dist. by CVL, Dublin, OH 43007 1-800-204-6335
100% Money Back Guarantee
Return to place of purchase if not satisfied.
©2020 Cardinal Health, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

38421 E9 C3

LEADER COUGH DM

dextromethorphan polistirex suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0187
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
POLISTIREX (UNII: 5H9W9GTW27)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)	
SUCROSE (UNII: C151H8M554)	
TARTARIC ACID (UNII: W4888I119H)	
TRAGACANTH (UNII: 2944357O2O)	
TRIACETIN (UNII: XHX3C3X673)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0187-1	1 in 1 CARTON	12/05/2016	

1	89 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	12/05/2016	

Labeler - Cardinal Health 110, LLC. dba Leader (063997360)

Revised: 6/2021

Cardinal Health 110, LLC. dba Leader