

AFTERPILL - levonorgestrel tablet

Syzygy Healthcare Solutions, LLC PO Box 588, Westport, CT 06880

AfterPill®

(levonorgestrel) Tablet, 1.5 mg

Drug Facts

Active Ingredient

Levonorgestrel USP 1.5 mg

Purpose

Emergency contraceptive

Indications

Use for women to reduce chance of pregnancy after unprotected sex (if a contraceptive failed or if you did not use birth control).

Warnings

Allergy alert: Do not use if you have ever had an allergic reaction to levonorgestrel.

Sexually transmitted diseases (STDs) alert: This product does **not** protect against HIV/AIDS or other STDs

Do not use

- if you are already pregnant (because it will not work)
- for regular birth control

Ask a doctor or pharmacist before use if you are taking efavirenz (HIV medication) or rifampin (tuberculosis treatment) or medication for seizures (epilepsy). These medications may reduce the effectiveness of levonorgestrel.

When using this product you may have

- menstrual changes
- nausea
- lower stomach (abdominal) pain
- tiredness
- headache
- dizziness
- breast pain
- vomiting

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-

1222) right away.

Directions

- **take as soon as possible within 72 hours (3 days) after unprotected sex. The sooner you take it the better it will work.**
- **if you vomit within 2 hours after taking the medication, call a healthcare professional to find out if you should repeat the dose.**

Other Information

- **read the instructions, warnings and enclosed product leaflet before use .**
- this product works mainly by preventing ovulation (egg release). It may also prevent fertilization of a released egg (joining of sperm and egg) or attachment of a fertilized egg to the uterus (implantation).
- **tablet is enclosed in wallet seal. Do not use if the wallet seal is broken .**
- store at 25°C (77°C); excursions permitted to 15 to 30°C (59 to 86°C) [see USP Controlled Room temperature].

Inactive Ingredients

colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, and povidone.

Questions or comments?

For more information, ask your healthcare professional or call 1-855-889-3012

M-F 8:00 am -4:00 pm (CST)

Visit our website at www.afterpill.com

AfterPill®

(levonorgestrel) Tablet, 1.5 mg

Emergency Contraceptive

One Tablet. One Dose.

What You Need to Know

What is AfterPill®?

AfterPill® is emergency contraception that helps prevent pregnancy after birth control failure or unprotected sex. It is a **backup** method of preventing pregnancy and should not be used as regular birth control.

What AfterPill Solo is not.

AfterPill will not work if you are already pregnant and will not affect an existing pregnancy. AfterPill will not protect you from HIV infection (the virus that causes AIDS) and other sexually transmitted diseases (STDs).

When should I use AfterPill?

The sooner you take emergency contraception, the better it works. You should use AfterPill within 72 hours (3 days) **after you have had unprotected sex.**

AfterPill is a backup or emergency method of birth control you can use when:

- your regular birth control was used incorrectly or failed
- you did not use any birth control method

When not to use AfterPill?

AfterPill should not be used:

- as a regular birth control method, because it's not as effective as regular birth control.
- if you are already pregnant, because it will not work.
- if you are allergic to levonorgestrel or any other ingredients in AfterPill.

When should I talk to a doctor or pharmacist?

Ask a doctor or pharmacist before use if you are taking efavirenz (HIV medication) or rifampin (tuberculosis treatment) or medication for seizures (epilepsy). These medications may reduce the effectiveness of AfterPill and increase your chance of becoming pregnant. Your doctor may prescribe another form of emergency contraception that may not be affected by these medications.

How does AfterPill work?

AfterPill works before release of an egg from the ovary. As a result, AfterPill usually stops or delays the release of an egg from the ovary. AfterPill is one tablet that contains a higher dose of levonorgestrel than birth control pills and works in a similar way to prevent pregnancy.

How can I get the best results from AfterPill?

You have 72 hours (3 days) to try to prevent pregnancy after birth control failure or unprotected sex. **The sooner you take AfterPill, the better it works.**

How effective is AfterPill?

If AfterPill is taken as directed, it can significantly decrease the chance that you will get pregnant. About 7 out of every 8 women who would have gotten pregnant will not become pregnant.

How will I know AfterPill worked?

You will know AfterPill has been effective when you get your next period, which should come at the expected time, or within a week of the expected time. If your period is delayed beyond 1 week, it is possible you may be pregnant. You should get a pregnancy test and follow up with your healthcare professional.

Will I experience any side effects?

- some women may have changes in their period, such as a period that is heavier or lighter or a period that is early or late. **If your period is more than a week late, you may be pregnant.**
- If you have severe abdominal pain, you may have an ectopic pregnancy, and should get immediate medical attention.
- when used as directed, AfterPill is safe and effective. Side effects may include changes in your period, nausea, lower stomach (abdominal) pain, tiredness, headache, dizziness, and breast tenderness.
- if you vomit within 2 hours of taking the medication, call a healthcare professional to

find out if you should repeat the dose.

What if I still have questions about AfterPill?

If you have questions or need more information, call our toll-free number, 1-855-889-3012 or visit our website at www.afterpill.com.

Other Information

Keep this and all medication out of reach of children:

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Do not use if the wallet seal is opened.

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Active Ingredient: levonorgestrel 1.5 mg

Inactive Ingredients: colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, and povidone.

AfterPill® is a registered trademark of Syzygy Healthcare Solutions, LLC.

Manufactured in India for

Syzygy Healthcare Solutions, LLC



Westport, CT 06880

Revised: May 2023

ID#: 273469

If you are sexually active, you should see a healthcare provider for routine checkups. Your healthcare provider will talk to you about and, if necessary, test you for sexually transmitted diseases, teach you about effective methods of routine birth control, and answer any other questions you may have.

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

AfterPill® (levonorgestrel) Tablet, 1.5 mg

NDC 64122-605-01

Wallet Label: 1 Tablet

Drug Facts

Active ingredient
Levonorgestrel USP 1.5 mg

Purpose
Emergency contraceptive

Use
For women to reduce chance of pregnancy after unprotected sex (if a contraceptive failed or if you did not use birth control)

Warnings
Allergy alert: Do not use if you have ever had an allergic reaction to levonorgestrel

Sexually transmitted diseases (STDs) alert: This product does not protect against HIV/AIDS or other STDs

Do not use
■ if you are already pregnant (because it will not work)
■ for regular birth control

Ask a doctor or pharmacist before use if you are taking efavirenz (HIV medication) or ritonavir (tuberculosis treatment) or medication for seizures (epilepsy). These medications may reduce the effectiveness of levonorgestrel.

When using this product you may have
■ menstrual changes ■ dizziness ■ nausea ■ headache ■ breast pain ■ lower stomach/abdominal pain ■ vomiting

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ take as soon as possible within 72 hours (3 days) after unprotected sex. The sooner you take it the better it will work.

Drug Facts (continued)

■ If you vomit within 2 hours after taking the medication, call a healthcare professional to find out if you should repeat the dose.

Other information
■ read the instructions, warnings and enclosed product insert before use.
■ this product works mainly by preventing ovulation (egg release). It may also prevent fertilization of a released egg (joining of sperm and egg) or attachment of a fertilized egg to the uterus (implantation).
■ tablet is enclosed in wallet seal.
Do not use if the wallet seal is broken.
■ store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) (see USP Controlled Room Temperature).

Inactive ingredients colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, and povidone

Questions or comments?
For more information, ask your healthcare professional or call 1-855-889-3012 M-F 8:00 am - 4:00 pm (CST). Visit our website at www.afterpill.com

NDC 64122-605-01

The sooner you take AfterPill, the more effective it is

Take as soon as possible, up to 3 days (72 hours) after unprotected sex

Will not harm an existing pregnancy

Contains One 1.5mg Tablet

Not for regular birth control

Reduces the chance of pregnancy after unprotected sex

Contains the emergency contraceptive medicine doctors recommend most

AfterPill.
Levonorgestrel Tablet, 1.5 mg

Emergency Contraceptive

Manufactured in India with FDA authorization

LOT NO. _____

EXP. _____

273164

W.L. 28/8/2010

PUSH TABLET FROM FRONT

Manufactured for:

Syzygy
HEALTHCARE SOLUTIONS

Westport, CT 06609

AFTERPILL

levonorgestrel tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:64122-605
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
LEVONORGESTREL (UNII: 5W7SIA7YZW) (LEVONORGESTREL - UNII:5W7SIA7YZW)		LEVONORGESTREL	1.5 mg
Inactive Ingredients			
Ingredient Name			Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			

POVIDONE K30 (UNII: U725QWY32X)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	WHITE (white to off white)		Score	no score
Shape	ROUND (round biconvex)		Size	9mm
Flavor			Imprint Code	LU;S25
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64122-605-01	1 in 1 BLISTER PACK; Type 0: Not a Combination Product	09/14/2017	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA201446	09/14/2017	

Labeler - Syzygy Healthcare Solutions, LLC PO Box 588, Westport, CT 06880 (079360097)

Registrant - LUPIN LIMITED (675923163)

Establishment

Name	Address	ID/FEI	Business Operations
LUPIN LIMITED		650582310	MANUFACTURE(64122-605) , PACK(64122-605)

Revised: 5/2025

Syzygy Healthcare Solutions, LLC PO Box 588, Westport, CT 06880