

HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet H E B

Drug Facts

Active ingredients (in each gelta)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives

- facial swelling
- shock
- asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- adults take more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood

- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red and swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours; do not take more than 8 geltabs in 24 hours
- children under 12 years: do not use unless directed by a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

black printing ink, corn starch, D&C yellow #10 aluminum lake, edetate disodium, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hydroxypropyl cellulose, hydroxypropyl cellulose-low substituted, hypromellose, iron oxide black, maltitol, povidone, purified water, stearic acid, talc, titanium dioxide

Questions or comments?

Call toll free: 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Excedrin® Extra Strength active ingredients**

Extra Strength

Headache Relief

Acetaminophen, 250 mg

Aspirin (NSAID), 250 mg

Caffeine, 65 mg

Pain Reliever/Pain Reliever Aid

GELTABS

**This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

MADE WITH PRIDE AND CARE FOR H-E-B® SAN ANTONIO, TX 78204

Package Label

Compare to Excedrin® Extra Strength active ingredients™

NDC 37808-586-20

H-E-B
Extra Strength
Headache Relief
Acetaminophen, 250 mg
Aspirin (NSAID), 250 mg
Caffeine, 65 mg

Pain Reliever/Pain Reliever Aid

actual size

20 GELTABS

Drug Facts

Active Ingredients (in each geltaab)

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID).....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to
headache a cold arthritis muscular aches
toothache premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness blisters rash. If a skin reaction occurs, stop use and seek medical help right away. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: hives facial swelling shock asthma (wheezing). **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

Drug Facts (continued)

are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more for a longer time than directed. **Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you have asthma you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis taking any other drug or are under a doctor's care for any serious condition.

Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. you experience

Drug Facts (continued)

hydroxypropyl cellulose-low substituted, hypromellose, iron oxide black, malitol, povidone, purified water, stearic acid, talc, titanium dioxide

Questions or comments? Call toll free: 1-877-753-3838 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

MADE WITH PAPER AND CARBON FROM H-E-B® SAN ANTONIO, TX 78204

LOT No.: 4122024295
Exp. Date:

PLD-0314C FC008738 39694 - 2305

100% GUARANTEE
If you are not satisfied with your purchase, please return it to the store for a full refund. No time limit on return.

Drug Facts (continued)

any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better ringing in the ears or loss of hearing occurs painful area is red or swollen pain gets worse or lasts for more than 10 days fever gets worse or lasts for more than 3 days any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions do not use more than directed drink a full glass of water with each dose adults and children 12 years and over: take 2 geltaabs every 6 hours; do not take more than 8 geltaabs in 24 hours children under 12 years: do not use unless directed by a doctor

Other information

store between 20°-25°C (68°-77°F)

Inactive ingredients black printing ink, corn starch, D&C yellow #10 aluminum lake, edetate disodium, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hydroxypropyl cellulose,

HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-586
--------------	----------------	--------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MALTITOL (UNII: D65DG142WK)	
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYDROXYPROPYL CELLULOSE, LOW SUBSTITUTED (UNII: 2165RE0K14)	

Product Characteristics

Color	green, white	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	P
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-586-80	80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/30/2021	
2	NDC:37808-586-20	1 in 1 BOX	07/30/2021	
2		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/30/2021	

Labeler - H E B (007924756)

Revised: 4/2024

H E B