

**DAYTIME AND NIGHTTIME COLD AND FLU- acetaminophen,
dextromethorphan hydrobromide, phenylephrine hydrochloride, doxylamine
succinate
Strive Pharmaceuticals Inc.**

***RIGHT REMEDIES* DayTime and NightTime Cold & Flu**

***RIGHT REMEDIES* DayTime Cold & Flu**

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine oxidase

inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless
• pain, nasal congestion or cough get worse or last more than 7 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present
• new symptoms occur
• cough comes back, or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away.
Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see **overdose warning**
- do not exceed 4 doses per 24 hrs

adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use

Other information

- store at room temperature between 20-25°C (68-77°F)

Inactive ingredients

edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments

1-888-577-8033 Monday - Friday 8am - 4pm EST

RIGHT REMEDIES NightTime Cold & Flu

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 15 mg
Doxylamine succinate 6.25 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms: • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever • runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

• taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see **overdose warning**
- do not exceed 4 doses per 24 hrs

adults and children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use

Other information

- store at room temperature between 20-25°C (68-77°F)

Inactive ingredients

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments

1-888-577-8033 Monday - Friday 8am - 4pm EST

Compare to the active ingredients of **Vicks® DayQuil™** Cold & Flu Relief LiquiCaps™*

Compare to the active ingredients of **Vicks® NyQuil™** Cold & Flu Relief LiquiCaps™*

RIGHT REMEDIES

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil™, NyQuil™ and LiquiCaps™.


Distributed by: Strive Pharmaceuticals Inc.,
East Brunswick, NJ 08816

Product of UAE

Packaged and Quality Assured in the USA

REV.00- 072022

Packaging

<p>Questions or comments 1-888-577-8033 Monday - Friday 8am - 4pm EST</p> <p>Active ingredients D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution</p> <p>Other information ■ store at room temperature between 20-25°C (68-77°F)</p> <p>Directions ■ take only as directed - see overdose warning ■ do not exceed 4 doses per 24 hrs adults and children 12 yrs & over: 2 softgels with water every 4 hrs children 4 to under 12: ask a doctor children under 4 yrs: do not use</p> <p>Drug Facts (continued) (Daytime)</p>	<p>Questions or comments 1-888-577-8033 Monday - Friday 8am - 4pm EST</p> <p>Active ingredients D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution</p> <p>Other information ■ store at room temperature between 20-25°C (68-77°F)</p> <p>Directions ■ take only as directed - see overdose warning ■ do not exceed 4 doses per 24 hrs adults and children 12 yrs & over: 2 softgels with water every 6 hrs children 4 to under 12: ask a doctor children under 4 yrs: do not use</p> <p>Drug Facts (continued) (Nighttime)</p>
<p>or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough get worse or last more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>	<p>pharmacist before taking this product. ■ to make a child sleep know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema ■ trouble urinating due to enlarged prostate gland</p> <p>Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin</p> <p>When using this product, ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, & tranquilizers may increase drowsiness</p> <p>Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts.</p> <p>If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>
<p>BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING Manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® and LiquiCaps™ Strive Pharmaceuticals Inc., East Brunswick, NJ 08816 Product of the USA</p> <p>3 70692 00058 0</p>  	

DO NOT USE IF BLIS
 *This product is not manufactured by Vicks® DayQuil™, NyQuil™
 Distributed by: Strive Pharm
 East Brunswick, NJ 08816
 Product of UAE
 Packaged and Quality Assured

CT 7069280974
 REV 03-072022

Drug Facts (continued) (Nighttime)
 Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's

Drug Facts (continued) (Daytime)
 Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions,

Drug Facts (Nighttime)
Active ingredients (in each softgel)
 Acetaminophen 325 mg.....Pain reliever/Fever reducer
 Dextromethorphan HBr 15 mg.....Cough suppressant
 Doxylamine succinate 6.25 mg.....Antihistamine
Uses temporarily relieves common cold/flu symptoms: ■ cough due to minor throat & bronchial irritation ■ sore throat & pain ■ fever ■ runny nose and sneezing
Warnings: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product. Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (Daytime)
Active ingredients (in each softgel)
 Acetaminophen 325 mg.....Pain reliever/Fever reducer
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Phenylephrine HCl 5 mg.....Nasal decongestant
Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever
Warnings: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product. Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION



Compare to the active ingredients of Vicks® DayQuil™ Cold & Flu Relief LiquiCaps™*

Compare to the active ingredients of Vicks® NyQuil™ Cold & Flu Relief LiquiCaps™*

NDC 70692-809-74

DayTime

Cold & Flu

- Acetaminophen 325 mg
Pain reliever/ Fever reducer
- Dextromethorphan HBr 10 mg
Cough Suppressant
- Phenylephrine HCL 5 mg
Nasal Decongestant



temporarily relieves common cold/flu symptoms: headache, fever, sore throat, minor aches and pains, nasal congestion, and cough

32
softgels



NightTime

Cold & Flu

- Acetaminophen 325 mg
Pain reliever/ Fever reducer
- Dextromethorphan HBr 15 mg
Cough Suppressant
- Doxylamine Succinate 6.25 mg
Antihistamine



temporarily relieves common cold/flu symptoms: headache, fever, sore throat, minor aches and pains, runny nose, sneezing, and cough

16
softgels



DRUG FACTS TABLE

Drug Facts (Nighttime)	Drug Facts (Daytime)
<p>Active ingredients (in each softgel) <i>Purpose</i></p> <p>Acetaminophen 325 mg.....Pain reliever/Fever reducer Dextromethorphan HBr 15 mg.....Cough suppressant Doxylamine succinate 6.25 mg.....Antihistamine</p>	<p>Active ingredients (in each softgel) <i>Purpose</i></p> <p>Acetaminophen 325 mg.....Pain reliever/Fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant</p>
<p>Uses temporarily relieves common cold/flu symptoms: ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever ■ runny nose and sneezing</p>	<p>Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever</p>
<p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	<p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>

Drug Facts (continued) (Nighttime)	Drug Facts (continued) (Daytime)
<p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ to make a child sleep</p> <p>Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema ■ trouble urinating due to enlarged prostate gland</p> <p>Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin</p> <p>When using this product ■do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, & tranquilizers may increase drowsiness</p> <p>Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>	<p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough get worse or last more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>

Drug Facts (continued) (Nighttime)	Drug Facts (continued) (Daytime)												
<p>Directions</p> <p>■ take only as directed - see overdose warning ■ do not exceed 4 doses per 24 hrs</p> <table border="1"> <tr> <td>adults and children 12 yrs & over</td> <td>2 softgels with water every 6 hrs</td> </tr> <tr> <td>children 4 to under 12</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>	adults and children 12 yrs & over	2 softgels with water every 6 hrs	children 4 to under 12	ask a doctor	children under 4 yrs	do not use	<p>Directions</p> <p>■ take only as directed - see overdose warning ■ do not exceed 4 doses per 24 hrs</p> <table border="1"> <tr> <td>adults and children 12 yrs & over</td> <td>2 softgels with water every 4 hrs</td> </tr> <tr> <td>children 4 to under 12</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>	adults and children 12 yrs & over	2 softgels with water every 4 hrs	children 4 to under 12	ask a doctor	children under 4 yrs	do not use
adults and children 12 yrs & over	2 softgels with water every 6 hrs												
children 4 to under 12	ask a doctor												
children under 4 yrs	do not use												
adults and children 12 yrs & over	2 softgels with water every 4 hrs												
children 4 to under 12	ask a doctor												
children under 4 yrs	do not use												
<p>Other information</p> <p>■ store at room temperature between 20-25°C (68-77°F)</p>	<p>Other information</p> <p>■ store at room temperature between 20-25°C (68-77°F)</p>												
<p>Inactive ingredients</p> <p>D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution</p>	<p>Inactive ingredients</p> <p>edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution</p>												
<p>Questions or comments</p> <p>1-888-577-8033 Monday - Friday 8am - 4pm EST</p>	<p>Questions or comments</p> <p>1-888-577-8033 Monday - Friday 8am - 4pm EST</p>												

DAYTIME AND NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-809
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-809-74	1 in 1 CARTON; Type 0: Not a Combination Product	04/05/2023	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	4 BLISTER PACK	32
Part 2	4 BLISTER PACK	16

Part 1 of 2

DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Item Code (Source)	NDC:70692-807
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	

GLYCERIN (UNII: PDC6A3C0OX)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
WATER (UNII: 059QF0KO0R)
SORBITOL (UNII: 506T60A25R)
SORBITAN (UNII: 6O92ICV9RU)

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	807
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/05/2023	

Part 2 of 2

NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:70692-808
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	808
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/05/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/05/2023	

Labeler - Strive Pharmaceuticals Inc. (080028013)