# IBUPROFEN JUNIOR- ibuprofen tablet, chewable Physicians Total Care, Inc.

-----

## Amerisource Bergen Junior Strength Ibuprofen Tablets, 100 mg Drug Facts

### **Active ingredient (in each tablet)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever/fever reducer

#### Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

## **Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

#### Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

## Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

## When using this product

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

## Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within the first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- find the right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart				
Weight (lb)	Age (yr)	Tablets		
under 24	under 24 under 2			
24-35	2-3	1		
36-47	4-5	1 ½		
48-59	6-8	2		
60-71	9-10	2 ½		
72-95	11	3		

#### Other information

- phenylketonurics: contains phenylalanine 6 mg per tablet
- do not use if printed seal under cap is broken or missing
- store at 20-25°C (68-77°F)
- see end panel for lot number and expiration date

### **Inactive ingredients**

acesulfame potassium, ammonium glycerrhizin, aspartame, carnauba wax, croscarmellose sodium, FD&C yellow no. 6 aluminum lake, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, prosweet, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

#### Questions or comments?


1-800-719-9260

#### **Principal Display Panel**

junior strength Ibuprofen tablets, 100 mg

Pain Reliever/Fever Reducer (NSAID)

Lasts up to 8 hours NDC 54868-5623-0 24 chewable tablets



Compare to Motrin® Junior Strength active ingredient See New Warnings Information FOR AGES 2 TO 11

## Additional barcode labeling by:

Physicians Total Care, Inc. Tulsa, Oklahoma 74146

## **IBUPROFEN JUNIOR**

ibuprofen tablet, chewable

Droduct	Information
Product	IIIIOTIIIauoii

Product Type HUMAN OTC DRUG Item Code (Source) NDC:54868-5623(NDC:24385-546)

Route of Administration ORAL

## **Active Ingredient/Active Moiety**

,				
	Ingredient Name	Basis of Strength	Strength	
	IBUPRO FEN (UNII: WK2XYI10 QM) (IBUPRO FEN - UNII: WK2XYI10 QM)	IBUPROFEN	100 mg	

## **Inactive Ingredients**

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29 V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SOYBEAN OIL (UNII: 241ATL177A)	
SUCCINIC ACID (UNII: AB6MNQ6J6L)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	

## **Product Characteristics**

Color	ORANGE	Score	2 pieces
Shape	ROUND	Size	13mm
Flavor	ORANGE	Imprint Code	L461
Contains			

## **Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54868-5623-0	1 in 1 CARTON		
1		24 in 1 BOTTLE		

## **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076359	10/16/2007	

## Labeler - Physicians Total Care, Inc. (194123980)

Establishment			
Name	Address	ID/FEI	Business Operations
Physicians Total Care, Inc.		194123980	relabel

Revised: 5/2012 Physicians Total Care, Inc.