

**BISACODY ENTERIC COATED- bisacodyl tablet, coated
MEIJER, INC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Bisacodyl USP 5 mg

Purpose

Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Dulcolax® Laxative

LAXATIVE

Bisacodyl | 5 mg

Stimulant laxative

Gentle, Predictable, Overnight Relief

ENTERIC COATED

Tablets

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals Inc., distributor of Dulcolax® Laxative.

DIST. BY MEIJER DISTRIBUTION, INC.

GRAND RAPID, MI 49544

www.meijer.com

Product Label



NDC 41250-191-25

*Compare to the active ingredient in Dulcolax® Laxative

laxative

Bisacodyl | 5 mg Stimulant laxative

Gentle, Predictable, Overnight Relief

ENTERIC
COATED

25 Tablets



Actual Size

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



WWW.MEIJER.COM/SATISFACTION

PID 428392



PLD-F23K
FC005694

Questions or comments? Call
1-877-753-3935 Monday-Friday 9AM-5PM EST

Inactive ingredients acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

Other information
■ store between 20-25°C (68-77°F)

- a sudden change in bowel habits that lasts more than 2 weeks
 - do not chew or crush tablet(s)
 - do not use within 1 hour after taking an antacid or milk
 - it may cause stomach discomfort, faintness and cramps
- When using this product**
- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
 - you need to use a laxative for more than 1 week
- Stop use and ask a doctor if**

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Bisacodyl USP 5 mg.....Stimulant laxative</p> <p>Purpose</p>		<p>Drug Facts</p> <p>Use</p> <ul style="list-style-type: none"> for temporary relief of occasional constipation and irregularly this product generally produces a bowel movement in 6 to 12 hours <p>Warnings</p> <p>Do not use if you cannot swallow without chewing.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> stomach pain, nausea or vomiting 	
<p>Directions</p> <p>take with a glass of water</p> <p>adults and children 12 years of age and over take 1 to 3 tablets in a single daily dose</p> <p>children 6 to under 12 years of age take 1 tablet in a single daily dose</p> <p>children under 6 years of age ask a doctor</p>		<p>Use</p> <p>if pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	

MEIJER Laxative

BISACODYL ENTERIC COATED			
bisacodyl tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-191
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)		BISACODYL	5 mg
Inactive Ingredients			
Ingredient Name			Strength
ACACIA (UNII: 5C5403N26O)			
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)			
ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
GELATIN (UNII: 2G86QN327L)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
SHELLAC (UNII: 46N107B71O)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)
SUCROSE (UNII: C151H8M554)
POVIDONE (UNII: FZ989GH94E)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)
ALUMINUM OXIDE (UNII: LMI26O6933)

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	TCL003
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-191-25	25 in 1 CARTON	05/31/2016	05/31/2025
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	05/31/2016	05/31/2025

Labeler - MEIJER, INC. (006959555)