

# **POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution A-S Medication Solutions**

-----

## **Polyethylene Glycol 3350 Drug Facts**

### **Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 g (cap filled to line)

### **Purpose**

Osmotic Laxative

### **Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### **Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

### **Do not use**

if you have kidney disease, except under the advice and supervision of a doctor

### **Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug

### **When using this product**

you may have loose, watery, more frequent stools

### **Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - use once a day
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - do not combine with starch-based thickeners used for difficulty swallowing
  - ensure that the powder is fully dissolved before drinking
  - do not drink if there are any clumps
  - do not use more than 7 days
- children 16 years of age or under: ask a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

### **Inactive ingredients**

none

### **Questions or comments?**

**1-800-719-9260**

### **Polyethylene Glycol 3350**

NDC 50090-2430-0  
A-S Medication Solutions, LLC  
Product No. 6524-0  
LOT

POLYETHYLENE  
GLYCOL 3350  
POWDER FOR SOLUTION

14 ONCE - DAILY DOSES  
ACTIVE INGREDIENT (ON EACH DOSE)  
POLYETHYLENE GLYCOL 3350, 17 G  
(CAP FILLED TO LINE)  
PURPOSE: OSMOTIC LAXATIVE

STORE AT 68 TO 77 DEGREES F

8.3 OZ (238 G)



GTIN: 00350090243001

DISTRIBUTED BY:  
A-S Medication Solutions  
Libertyville, IL 60048

SOURCE NDC: 45802-868-02



## POLYETHYLENE GLYCOL 3350

polyethylene glycol 3350 powder, for solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50090-2430(NDC:45802-868)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50090-2430-0	238 g in 1 BOTTLE; Type 0: Not a Combination Product	07/05/2016	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	10/13/2009	

**Labeler** - A-S Medication Solutions (830016429)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
A-S Medication Solutions		830016429	RELABEL(50090-2430)

Revised: 12/2023

A-S Medication Solutions