BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN FAIR-titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN LIGHT- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MEDIUM- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MEDIUM-TAN- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN TANtitanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN TAN DEEP- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN DEEP- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN RICHtitanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MAHOGANY- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN FAIR-LIGHT- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN LIGHT-MEDIUM- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN DEEP-SAND- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN RICH HONEY- titanium dioxide and zinc oxide liquid

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MAHOGANY HONEY- titanium dioxide and zinc oxide liquid BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN

ESPRESSO- titanium dioxide and zinc oxide liquid Tarte, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

BB tinted treatment primer Broad Spectrum SPF 30 Sunscreen

Drug Facts

Active Ingredients	Purpose
Titanium Dioxide 4.55%	Sunscreen
Zinc Oxide 3.5%	Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, mica, polysilicone-11, polymethylsilsesquioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Dist. by tarte, Inc. New York, NY 10018.

PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - fair

tarte high-performance naturals™

BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, mica, polysilicone-11, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina. triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

www.tarte.com

high-performance naturals

- triple-B complex: diamond powder, plants & lipoamino
- smoothing & brightening
- dermatologist tested
- wear alone for a sheer tint or under

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%..

 Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.Apply liberally 15 minutes before sun exposure.

- use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
 Sun Protection Measures.
 Spending time in the sun increases your risk of skin capper and early. your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15

or higher and other sun protection measures including: -Limit time in the sun, especially

from 10 a.m.-2 p.m.

-Wear long-sleeved shirts, pants, hats and sunglasses


- Other information
 Protect the product in this container from excessive heat and
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



FG01187-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - light

tarte high-performance naturals™ BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

light 12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, mica, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JO, UK.

@tartecosmetics

www.tarte.com

tarte

high-performance naturals

- triple-B complex: diamond powder, plants & lipoamino
- smoothing & brightening
- 12-hour primer & dermatologist
- wear alone for a sheer tint or under

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%..

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
 Apply liberally 15 minutes before

- Apply liberally 10 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Chidren under 6 months: Ask a doctor.
 Sun Protection Measures.
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection
- measures including:

 -Limit time in the sun, especially
- from 10 a.m.-2 p.m.

 -Wear long-sleeved shirts, pants, hats and sunglasses

Other information


- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



FG01188-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - medium

tarte high-performance naturals™ BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

medium 12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, mica, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

www.tarte.com

high-performance naturals:

- dermatologist tested

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%... .Sunscreen

Uses

 Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center

Directions

- Shake before use.
 Apply liberally 15 minutes before sun exposure.

 • Use a water resistant sunscreen if
- swimming or sweating.

 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
- Sun Protection Measures.
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection

measures including:

-Limit time in the sun, especially

from 10 a.m.-2 p.m.

-Wear long-sleeved shirts, pants, hats and sunglasses

Other information

Protect the product in this

ontainer from excessive heat and direct sunlight.


You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



FG01189-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - medium-tan

tarte
high-performance naturals™
BB tinted treatment primer
Broad Spectrum
SPF 30 Sunscreen

medium-tan 12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, mica, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina. triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

www.tarte.com

BB tinted treatment primer **Broad Spectrum**



- high-performance naturals
- smoothing & brightening
- dermatologist tested
- wear alone for a sheer tint or under

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%.. .Sunscreen

Uses

 Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center

Directions

right away.

- Shake before use.Apply liberally 15 minutes before sun exposure.

- sun exposure.

 Use a water resistant sunscreen if swimming or sweating.

 Reapply at least every 2 hours.

 Children under 6 months: Ask a doctor.

 Sun Protection Measures.

 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a regularly use a sunscreen with a Broad Spectrum SPF value of 15
- or higher and other sun protection measures including: -Limit time in the sun, especially
- from 10 a.m.-2 p.m.

 -Wear long-sleeved shirts, pants, hats and sunglasses

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



FG01570-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - tan

tarte high-performance naturals™ BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

tan 12-hour primer

high-performance naturals

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, mica, polymethylsilsesqui-oxane, betyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina. triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

@tartecosmetics

www.tarte.com

BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



dermatologist tested

sheer tint or under

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%.. .Sunscreen

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center

Directions

- Shake before use.
 Apply liberally 15 minutes before

- Apply liberally 13 minutes
 Apply liberally 13 minutes
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
 Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early.
- your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection
- measures including:

 _Limit time in the sun, especially
 from 10 a.m.-2 p.m.


 _Wear long-sleeved shirts, pants,
 hats and sunglasses

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



FG08096-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - tan-deep

tarte
high-performance naturals™

BB tinted treatment primer

Broad Spectrum

SPF 30 Sunscreen

tan-deep 12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, mica, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

@tartecosmetics

www.tarte.com

high-performance naturals

- dermatologist tested

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%.... Sunscreen

Uses

 Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center

Directions

- Shake before use.Apply liberally 15 minutes before

- Apply liberally 15 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
 Sun Protection Measures.
 Spending time in the sun increases your risk of skin capaca and eastly. your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection

measures including:

_Limit time in the sun, especially
from 10 a.m.-2 p.m.

_Wear long-sleeved shirts, pants,
hats and sunglasses

Other information


- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse-reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



BB tinted treatment primer Broad Spectrum



FG08129-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - deep

tarte
high-performance naturals™

BB tinted treatment primer

Broad Spectrum

SPF 30 Sunscreen

deep 12-hour primer

high-performance naturals

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polysilicone-11, polymethylsilsesqui-oxane, mica, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid cetyl acid, cetyl ' PEG/PPG-10/1 dimethicone, triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JO, UK.

#claypowered @tartecosmetics

www.tarte.com

Broad Spectrum SPF 30 Sunscreen

BB tinted treatment primer



triple-B complex: Drug Facts

diamond powder, plants & lipoamino

smoothing & brightening

12-hour primer & dermatologist

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%..

Uses

 Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
 Apply liberally 15 minutes before sun exposure.

- sun exposure.

 Use a water resistant sunscreen if swimming or sweating.

 Reapply at least every 2 hours.

 Children under 6 moriths: Ask a doctor.

 Sun Protection Measures.

 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a regularly use a sunscreen with a Broad Spectrum SPF value of 15 Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

 -Limit time in the sun, especially from 10 a.m.-2 p.m.

 -Wear long-sleeved shirts, pants, hats and sunglasses

- Other information
 Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



FG07880-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - rich

tarte high-performance naturals™ BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

rich

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, mica, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

#claypowered @tartecosmetics

www.tarte.com

BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



high-performance naturals

triple-B complex:

- smoothing & brightening
- dermatologist tested
- wear alone for a sheer tint or under

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%... .Sunscreen

Uses

 Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.Apply liberally 15 minutes before sun exposure.

- sun exposure.

 Use a water resistant sunscreen if swimming or sweating.

 Reapply at least every 2 hours.

 Children under 6 morths: Ask a doctor.

 Sun Protection Measures.

 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a regularly use a sunscreen with a Broad Spectrum SPF value of 15

or higher and other sun protection measures including:

-Limit time in the sun, especially from 10 a.m.-2 p.m.


-Wear long-sleeved shirts, pants, hats and sunglasses

Other information

- · Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, N. 08536-0222.



FG08099-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - mahogany

tarte high-performance naturals $^{\text{\tiny M}}$

BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

mahogany 12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, mica, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsi-lane, dipalmitoyl hydroxyproline, diamond powder, titanium dioxide, iron OX111FZ000131

USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

@tartecosmetics

www.tarte.com

tarte

high-performance naturals

- triple-B complex: diamond powder, plants & lipoamino
- 12-hour primer & dermatologist
- sheer tint or under

Drug Facts

Active Ingredients

Purpose

Titanium Dioxide 4.55%....Sunscreen Zinc Oxide 3.5%..

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If

swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
 Apply liberally 15 minutes before

- Apply liberally 13 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Chidren under 6 months: Ask a doctor.
 Sun Protection Measures.
 Spending time in the sun increases. your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection
- measures including:

 _Limit time in the sun, especially
 from 10 a.m.-2 p.m.

 _Wear long-sleeved shirts, pants,
 hats and sunglasses

- Other information
 Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



SPF 30 Sunscreen

BB tinted treatment primer **Broad Spectrum**

crueltyfree andvegan

FG08265-01



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - fair-light

tarte
high-performance naturals™
BB tinted treatment primer
Broad Spectrum

SPF 30 Sunscreen

fair-light 12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, diamond powder, stearic acid, cetyl PEG/PPG-10/1 cetyl PEG/PPG-10/1 dimethicone, alumina, dipalmitoyl hydroxyproline, triethoxycaprylylsi-lane, mica, titanium dioxide (Cl 77891), iron oxides (Cl 77499, Cl 77492, Cl 77499. OX100FP000371 OX100FP000371.

net 30 mL / 1 Fl. Oz. Made in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS, Wavre, BE / London WC2H 9JQ, UK.

@tartecosmetics www.tarte.com

tarte

high-performance naturals

- triple-B complex:
- smoothing & brightening
- dermatologist tested
- wear alone for a sheer tint or under foundation for

Drug Facts

Active Ingredients

Titanium Dioxide 4.55%...Sunscreen Zinc Oxide 3.5%.....SunscreenSunscreen

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs Do not use on darnaged or broken skin. When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
 Apply liberally 15 minutes before

- Apply liberally 15 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
 Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection. or higher and other sun protection measures including:
- measures including.

 -Limit time in the sun, especially from 10 a.m.-2 p.m.

 -Wear long-sleeved shirts, pants, hats and sunglasses

- Other information
 Protect the product in this container from excessive heat and direct sunlight.
- direct sunlight.


 You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ



BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



FG11684-02



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - light-medium

tarte high-performance naturals™ BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

light-medium

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesquioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, diamond powder, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, dipalmitoyl hydroxyproline, riydroxyprome, triethoxycaprylylsi-lane, mica, titanium dioxide (Cl 77891), iron oxides (Cl 77491, Cl 77492, Cl 77499. OX100FP000371.

in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS, Wavre, BE/London WC2H 9JQ, UK.

@tartecosmetics www.tarte.com

tarte

high-performance naturals

- triple-B complex:
- dermatologist tested
- sheer tint or under foundation for

Drug Facts

Active Ingredients

Titanium Dioxide 4.55%...Sunscreen Zinc Oxide 3.5%......Sunscreen

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs Do not use on darnaged or broken skin. When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center

Directions

- Shake before use.Apply liberally 15 minutes before

- Apply liberally 15 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
 Sun Protection Measures.
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
- -Limit time in the sun, especially from 10 a.m.-2 p.m.
 -Wear long-sleeved shirts, pants, hats and sunglasses

- Other information
 Protect the product in this container from excessive heat and direct sunlight.
- direct sunlight.


 You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ





FG11685-02

BB tinted treatment primer Broad Spectrum SPF 30 Sunscreen



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - deep sand

tarte high-performance naturals $^{\text{™}}$

BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

deep sand

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, diamond powder, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, dipalmitoyl hydroxyproline, triethoxycaprylylsi-lane, mica, titanium dioxide (CI 77891), iron oxides (CI 77491, CI 77492, CI 77499. OX100FP000371.

in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS, Wavre, BE/London WC2H 9JQ, UK.

@tartecosmetics www.tarte.com

tarte

high-performance naturals

- triple-B complex:
- dermatologist tested
- sheer tint or under foundation for

Drug Facts

Active Ingredients

Titanium Dioxide 4.55%...Sunscreen Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs Do not use on damaged or broken skin. When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.Apply liberally 15 minutes before

- Apply liberally 15 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
 Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection. or higher and other sun protection measures including:
- -Limit time in the sun, especially from 10 a.m.-2 p.m. -Wear long-sleeved shirts, pants, hats and sunglasses

Other information

- Protect the product in this
- ontainer from excessive heat and direct sunlight.


 You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



FG11686-02



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - rich honey

tarte high-performance naturals™ BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

rich honey 12-hour primer

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilesequioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 sostearate, diamond powder stearic acid powder, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, dipalmitoyl alumina, dipalimitoyi hydroxyproline, triethoxycaprylylsi-lane, mica, titanium dioxide (Cl 77891), iron oxides (Cl 77491, Cl 77492, Cl 77499. OX100FP000371.

in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS, Wavre, BE/London WC2H 9JQ, UK.

@tartecosmetics www.tarte.com

tarte

high-performance naturals

- dermatologist tested
- sheer tint or under foundation for

Drug Facts

Active Ingredients

Titanium Dioxide 4.55%...Sunscreen Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs Do not use on damaged or broken skin. When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.Apply liberally 15 minutes before

- Apply liberally 15 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 months: Ask a doctor.
 Sun Protection Measures.
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
- -Limit time in the sun, especially from 10 a.m.-2 p.m. -Wear long-sleeved shirts, pants, hats and sunglasses

- Other information
 Protect the product in this container from excessive heat and direct sunlight.
- direct sunlight.


 You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ





FG11687-02

BB tinted treatment primer Broad Spectrum SPF 30 Sunscreen



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - mahogany honey

tarte high-performance naturals $^{\text{\tiny M}}$

BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

mahogany honey

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, diamond powder, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, dipalmitoyl aiumna, dipaimitoyi hydroxyproline, triethoxycaprylylsi-lane, mica, titanium dioxide (Cl 77891), iron oxides (Cl 77491, Cl 77492, Cl 77499, OX100FP000371.

in USA. Dist. by starte, Inc. New York, NY 10018. All rights reserved. BIORIUS, Wavre, BE/London WC2H 9JQ, UK.

@tartecosmetics www.tarte.com

tarte

high-performance naturals

- 12-hour primer &
- wear alone for a sheer tint or under

Drug Facts

Active Ingredients

Titanium Dioxide 4.55%... Zinc Oxide 3.5%....SunscreenSunscreen

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs. Do not use on damaged or broken skin. When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

- Shake before use.
 Apply liberally 15 minutes before

- Apply liberally 15 minutes before sun exposure.
 Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Children under 6 morths: Ask a doctor.
 Sun Protection Measures.
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection or higher and other sun protection measures including:
- -Limit time in the sun, especially from 10 a.m.-2 p.m. -Wear long-sleeved shirts, pants, hats and sunglasses

Other information


- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction LLC, P.O. Box 22, Plainsboro, N. 08536-0222



BB tinted treatment primer **Broad Spectrum** SPF 30 Sunscreen



FG11688-02



PRINCIPAL DISPLAY PANEL - 30 mL Tube Carton - espresso

tarte high-performance naturals $^{\text{™}}$

BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen

espresso

Drug Facts (continued)

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesqui-oxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, diamond powder, stearic acid, cetyl PEG/PPG-10/1 cetyl PEG/PPG-10/1 dimethicone, alumina, dipalmitoyl hydroxyproline, triethoxycaprylylsi-lane, mica, titanium dioxide (Cl 77891), iron oxides (Cl 77491, Cl 77492, Cl 77499. OX100FP000371.

in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS, Wavre, BE/London WC2H 9JQ, UK.

@tartecosmetics www.tarte.com

tarte

high-performance naturals

- · wear alone for a sheer tint or under

Drug Facts

Active Ingredients

Titanium Dioxide 4.55%... Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
 If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs Do not use on darnaged or broken skin. When using this product keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center

Directions

right away.

- Shake before use.
 Apply liberally 15 minutes before

- Use a water resistant sunscreen if swimming or sweating.
 Reapply at least every 2 hours.
 Chidren under 6 months: Ask a doctor.
 Sun Protection Measures. Spending time in the sun increases. your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
- -Limit time in the sun, especially from 10 a.m.-2 p.m. -Wear long-sleeved shirts, pants, hats and sunglasses

Other information

- · Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, N.






FG11689-02

BB tinted treatment primer

Broad Spectrum SPF 30 Sunscreen



BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN FAIR

titanium dioxide and zinc oxide liquid

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-243	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15I	FIX9V2JP) TITANIUM DIOXIDE	45.5 mg in 1 mL	
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL	

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
MICA (UNII: V8A1AW0880)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
BENZIMIDAZOLE (UNII: E24GX49LD8)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-243- 01	1 in 1 CARTON	09/01/2012	
1		30 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
OTC monograph not final	part352	09/01/2012		

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN LIGHT

titanium dioxide and zinc oxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-244
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL		
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL		

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
MICA (UNII: V8A1AW0880)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
BENZIMIDAZOLE (UNII: E24GX49LD8)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51060-244- 01	1 in 1 CARTON	09/01/2012		
1		30 mL in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
OTC monograph not final	part352	09/01/2012		

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MEDIUM

titanium dioxide and zinc oxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-245
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL	
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL	

Inactive Ingredients		
Ingredient Name	Strength	
CYCLOMETHICONE 5 (UNII: OTHT5PCIOR)		
ISODODECANE (UNII: A8289P68Y2)		
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)		
MICA (UNII: V8A1AW0880)		
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)		
HEXYL LAURATE (UNII: 4CG9F9W01Q)		
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)		
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)		
ALUMINUM OXIDE (UNII: LMI2606933)		
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)		

DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
DIAMOND (UNII: 6GRV67N0U2)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-245- 01	1 in 1 CARTON	09/01/2012	
1		30 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	09/01/2012	

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MEDIUM-TAN

titanium dioxide and zinc oxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-246
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL	
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL	

Inactive Ingredients		
Ingredient Name	Strength	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)		
ISODODECANE (UNII: A8289P68Y2)		
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)		
MICA (UNII: V8A1AW0880)		
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)		
HEXYL LAURATE (UNII: 4CG9F9W01Q)		
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)		
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)		

STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
DIAMOND (UNII: 6GRV67N0U2)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-246- 01	1 in 1 CARTON	09/01/2012	
1		30 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	09/01/2012	

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN TAN

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-247
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: OTHT5PCIOR)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	

HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
CYCLOMETHICONE 6 (UNII: XHK3U310BA)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
DIAMOND (UNII: 6GRV67N0U2)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

ı	Packaging				
7	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
:	NDC:51060-247-	1 in 1 CARTON	09/01/2012		
:	L	30 mL in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Marketing End Date		
OTC monograph not final	part352	09/01/2012	

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN TAN DEEP

titanium dioxide and zinc oxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-248
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: OTHT5PCIOR)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
MICA (UNII: V8A1AW0880)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI2606933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
DIAMOND (UNII: 6GRV67N0U2)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

ı	Packaging				
4	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51060-248- 01	1 in 1 CARTON	09/01/2012		
1	L	30 mL in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date			
OTC monograph not final	part352	09/01/2012	

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN DEEP

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-249	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL	

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

F	Packaging					
#	tem Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:51060-249- 01	1 in 1 CARTON	09/01/2012			
1		30 mL in 1 TUBE; Type 0: Not a Combination Product				

Marketing Information				
Marketing Application Number or Monograph Marketing Start Mark Category Citation Date				
OTC monograph not final	part352	09/01/2012		

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN RICH

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-250
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL	
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL	

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
CYCLOMETHICONE 6 (UNII: XHK3U310BA)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
DIAMOND (UNII: 6GRV67N0U2)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51060-250- 01	1 in 1 CARTON	09/01/2012		
1		30 mL in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
OTC monograph not final	part352	09/01/2012		

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MAHOGANY

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-251
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients		
Ingredient Name	Strength	
CYCLOMETHICONE 5 (UNII: 0THT5PCIOR)		
ISODODECANE (UNII: A8289P68Y2)		
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)		
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)		
HEXYL LAURATE (UNII: 4CG9F9W01Q)		
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)		
MICA (UNII: V8A1AW0880)		
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)		
ALUMINUM OXIDE (UNII: LMI2606933)		
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)		
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)		
DIAMOND (UNII: 6GRV67N0U2)		
FERRIC OXIDE RED (UNII: 1K09F3G675)		
FERRIC OXIDE YELLOW (UNII: EX43802MRT)		
FERROSOFERRIC OXIDE (UNII: XM0M87F357)		

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-251- 01	1 in 1 CARTON	09/01/2012	
1		30 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	09/01/2012	

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN FAIR-LIGHT

titanium dioxide and zinc oxide liquid

Product	Inform	ation
Product	11110111	lation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:51060-364

Route of Administration TOPICAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP) ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z) ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z) 35 mg in 1 mL

Inactive Ingredients		
Ingredient Name	Strength	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)		
ISODODECANE (UNII: A8289P68Y2)		
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)		
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)		
HEXYL LAURATE (UNII: 4CG9F9W01Q)		
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)		
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)		
DIAMOND (UNII: 6GRV67N0U2)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)		
ALUMINUM OXIDE (UNII: LMI2606933)		
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)		
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)		
MICA (UNII: V8A1AW0880)		
FERRIC OXIDE YELLOW (UNII: EX43802MRT)		
FERRIC OXIDE RED (UNII: 1K09F3G675)		
FERROSOFERRIC OXIDE (UNII: XM0M87F357)		

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-364- 01	1 in 1 CARTON	09/01/2012	
1		30 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Info	ormation		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN LIGHT-MEDIUM

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-359
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCIOR)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI2606933)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

# Item Code Package Des 1 NDC:51060-359- 1 in 1 CARTON	Date Date
I IN LUBRIUM	00/01/2012
01	09/01/2012
30 mL in 1 TUBE; Type 0: N Product	Not a Combination

Marketing Information				
Marketing Category Application Number or Monog Citation	raph Marketing Start Date	Marketing End Date		
OTC MONOGRAPH NOT part352	09/01/2012			

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN DEEP-SAND

titanium dioxide and zinc oxide liquid

Product	Intorm	ation
FIUMULL		ation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:51060-360

Route of Administration TOPICAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCIOR)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1			

1	01 01-300-300-	1 in 1 CARTON	09/01/2012	1
1		30 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH NOT FINAL	part352	09/01/2012		

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN RICH HONEY

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-361	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCIOR)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51060-361- 01	1 in 1 CARTON	09/01/2012		
1		30 mL in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category Application Number or Monograph Citation Date Marketing Start Date				
OTC MONOGRAPH NOT FINAL	part352	09/01/2012		

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MAHOGANY HONEY

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-362
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL		
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL		

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	

FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51060-362- 01	1 in 1 CARTON	09/01/2012		
1		30 mL in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

BB TINTED TREATMENT PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN ESPRESSO

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-363	
Route of Administration	TOPICAL			

Ingredient Name Basis of	of Strength St	rength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP) TITANIUM	1 DIOXIDE 45.5	mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z) ZINC OXI	IDE 35 mg	g in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: OTHT5PCIOR)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
MICA (UNII: V8A1AW0880)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51060-363- 01	1 in 1 CARTON	09/01/2012		
1		30 mL in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

Labeler - Tarte, Inc. (027905186)

Revised: 8/2022 Tarte, Inc.