

PAIN RELIEF ASPIRIN LOW DOSE- aspirin tablet, delayed release
Rite Aid Corporation

Rite Aid 44-600A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor.
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

Compare to the active ingredient in
Bayer® Low Dose Aspirin**

NDC 11822-0600-7

**Talk to your doctor or other healthcare
provider before using this product for
your heart.**

LOW DOSE PAIN RELIEF

ASPIRIN

ASPIRIN 81 mg

PAIN RELIEVER (**NSAID**)

Aspirin regimen

Safety coated

ACTUAL

SIZE

300

ENTERIC

COATED TABLETS

NOT FOR

HOUSEHOLDS

WITH YOUNG

CHILDREN

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin. 50844 REV0122B60017

DISTRIBUTED BY: RITE AID, 200 NEWBERRY COMMONS, ETTERS, PA 17319

www.riteaid.com

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Compare to the active ingredient in Bayer® Low Dose Aspirin**

NDC 11822-0600-7
Talk to your doctor or other healthcare provider before using this product for your heart.

LOW DOSE PAIN RELIEF
ASPIRIN
ASPIRIN 81 mg
PAIN RELIEVER (NSAID)
Aspirin regimen
Safety coated

ACTUAL SIZE
300
ENTERIC COATED TABLETS

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN



**This product is not manufactured or distributed by Bayer AG, owner of the registered trademark, Bayer® Low Dose Aspirin. 50844 REV0122860017 DISTRIBUTED BY: RITE AID, 200 NEUBERRY COMMONS, ETTES, PA 17319 www.riteaid.com SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money. TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID) Pain reliever
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed ■ take a blood thinning (anticoagulant) or steroid drug



0 11822 12986 2
PEEL HERE FOR MORE DRUG FACTS

STOP PEELING

Questions or comments? 1-800-426-9391

Inactive ingredients corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Other information ■ store at 25° C (77° F); excursions permitted between 15° - 30° C (59° - 86° F) ■ use by expiration date on package

Directions ■ do not take more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ feel faint or ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur These could be signs of a serious condition.

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Rite Aid 44-600A

PAIN RELIEF ASPIRIN LOW DOSE			
aspirin tablet, delayed release			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0600
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg	
Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3S)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)
METHACRYLIC ACID (UNII: 1CS02G8656)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
POLYDEXTROSE (UNII: VH2XOU12IE)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
SHELLAC (UNII: 46N107B71O)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
DIMETHICONE (UNII: 92RU3N3Y1O)
WATER (UNII: 059QF0KO0R)
SODIUM BICARBONATE (UNII: 8MDF5V39QO)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIACETIN (UNII: XHX3C3X673)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0600-2	1 in 1 CARTON	03/01/2023	
1		32 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:11822-0600-3	1 in 1 CARTON	02/12/2008	
2		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:11822-0600-6	1 in 1 CARTON	02/12/2008	
3		200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:11822-0600-7	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/12/2008	
5	NDC:11822-0600-4	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/12/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/12/2008	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0600)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-0600)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(11822-0600) , pack(11822-0600)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0600)

Revised: 4/2024

Rite Aid Corporation