

PAIN RELIEF ASPIRIN LOW DOSE- aspirin tablet, coated

Rite Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid 44-600A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease

- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

**RITE
AID®
PHARMACY**

****Compare to the active ingredient in Bayer®**

LOW DOSE

pain relief

aspirin

aspirin 81 mg

pain reliever (NSAID)

aspirin

regimen

safety coated

ACTUAL
SIZE

120 ENTERIC COATED TABLETS

**This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer® Low Dose Aspirin.

50844 REV0613A60032

**DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011**

**IF YOU'RE NOT SATISFIED, WE'LL
HAPPILY REFUND YOUR MONEY.**

**TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING**

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients corn starch, D&G yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

RITE AID PHARMACY

LOW DOSE pain relief aspirin

aspirin 81 mg

pain reliever (NSAID)

120 ENTERIC COATED TABLETS

aspirin regimen safety coated ACTUAL SIZE

****Compare to the active ingredient in Bayer®**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Low Dose Aspirin. 50844 REV0613A60032

DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011

IF YOU'RE NOT SATISFIED, WE'LL HAPPILY REFUND YOUR MONEY.

Drug Facts (continued)

Questions or comments? 1-800-425-9391

ITEM 0398876

Drug Facts KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active Ingredient (in each tablet) Aspirin 81 mg (NSAID)*. *nonsteroidal anti-inflammatory drug Pain reliever

Purpose For the temporary relief of minor aches and pains or as recommended by your doctor

Warnings **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or the-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed

B-1702-600A-32-R
REV0613A60032

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

These could be signs of a serious condition.

Rite Aid 44-600A

PAIN RELIEF ASPIRIN LOW DOSE			
aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0600
Route of Administration	ORAL		
Active Ingredient/Active Moiety			

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	ROUND	Size	6 mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0600-3	1 in 1 CARTON	02/12/2008	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-0600-4	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/12/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	02/12/2008	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(11822-0600)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(11822-0600)

Revised: 5/2017

Rite Aid Corporation