

MOTION SICKNESS RELIEF- dimenhydrinate tablet
Topco Associates, LLC

TopCare 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-888-423-0139

Principal display panel

TopCare®
health

NDC 36800-930-12

**COMPARE TO DRAMAMINE® ORIGINAL
FORMULA ACTIVE INGREDIENT***

ORIGINAL FORMULA

Motion Sickness Relief

DIMENHYDRINATE, 50 mg - ANTIEMETIC

Prevents Nausea, Dizziness & Vomiting

12 TABLETS

For Children & Adults


actual size

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula. 50844 REV0518C19802

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ITASCA, IL 60143

©TOPCO LNKA0923 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula. 50844 REV0518C19802
 DISTRIBUTED BY TOPCO ASSOCIATES LLC
 ITASCA, IL 60143
 ©TOPCO LNKA0923 QUESTIONS? 1-888-423-0139
 topcare@topco.com www.topcarebrand.com



QUALITY
 GUARANTEED

B-1910-198-02-RR
 REV0518C19802

Scan here for more
 information or call 1-888-423-0139

No Print/No Varnish Area
 Lot & Exp. date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
 OPENED OR IF BLISTER UNIT IS TORN, BROKEN
 OR SHOWS ANY SIGNS OF TAMPERING

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet) Purpose
 Dimenhydrinate 50 mgAntiemetic

Uses for prevention and treatment of these symptoms associated with motion sickness: ■ nausea ■ vomiting ■ dizziness


Warnings
 Do not use for children under 2 years of age unless directed by a doctor.
 Ask a doctor before use if you have
 ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma
 ■ difficulty in urination due to enlargement of the prostate gland
 Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
 When using this product ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information
 ■ each tablet contains calcium 35 mg
 ■ see end flap for expiration date and lot number
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGING IS OPENED OR BLISTER IS TORN OR BROKEN
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments? 1-888-423-0139

Directions
 ■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity
 adults and children 12 years and over 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
 children 6 to under 12 years ½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
 children 2 to under 6 years ½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor



TopCare 44-198

MOTION SICKNESS RELIEF

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-930
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-930-12	2 in 1 CARTON	12/01/1992	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/01/1992	

Labeler - Topco Associates, LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-930)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(36800-930) , pack(36800-930)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(36800-930)