

BALSALAZIDE DISODIUM- balsalazide disodium capsule AvPAK

HIGHLIGHTS OF PRESCRIBING INFORMATION Balsalazide Disodium Capsules, USP

These highlights do not include all the information needed to use BALSALAZIDE DISODIUM CAPSULES safely and effectively. See full prescribing information for BALSALAZIDE DISODIUM CAPSULES.

BALSALAZIDE DISODIUM capsules, for oral administration Initial U.S. Approval: 2000

INDICATIONS AND USAGE

- Balsalazide is a locally acting aminosalicylate indicated for the treatment of mildly to moderately active ulcerative colitis in patients 5 years of age and older. (1)
- Safety and effectiveness of balsalazide beyond 8 weeks in children (ages 5 to 17 years) and 12 weeks in adults have not been established. (1)

DOSAGE AND ADMINISTRATION

- Adult dose is three 750 mg balsalazide disodium capsules 3 times a day (6.75 g/day) with or without food for 8 weeks. Some adult patients required treatment for up to 12 weeks. (2.1)
- Pediatric dose is EITHER: (2.2, 8.4)

1. Three 750 mg balsalazide disodium capsules 3 times a day (6.75 g/day) with or without food for 8 weeks. OR:

2. One 750 mg balsalazide disodium capsule 3 times a day (2.25 g/day) with or without food for up to 8 weeks. Capsules may be swallowed whole or may be opened and sprinkled on applesauce, then chewed or swallowed immediately. (2.3, 12.3)

DOSAGE FORMS AND STRENGTHS

Capsules: 750 mg (3)

CONTRAINDICATIONS

Patients with hypersensitivity to salicylates or to any of the components of balsalazide disodium capsules or balsalazide metabolites. Hypersensitivity reactions may include, but are not limited to the following: anaphylaxis, bronchospasm, and skin reaction. (4)

WARNINGS AND PRECAUTIONS

- Exacerbation of the symptoms of ulcerative colitis was reported in both adult and pediatric patients. Observe patients closely for worsening of these symptoms while on treatment. (5.1)
- Prolonged gastric retention of balsalazide may occur in patients with pyloric stenosis. (5.2)

ADVERSE REACTIONS

Most common adverse reactions (incidence $\geq 3\%$) are headache, abdominal pain, diarrhea, nausea, vomiting, respiratory infection, and arthralgia. Adverse reactions in children were similar. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact AvKARE at 1-855-361-3993 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

In an *in vitro* study using human liver microsomes, balsalazide and its metabolites were not shown to inhibit the major CYP enzymes evaluated (CYP1A2, CYP2C9, CYP2C19, CYP2D6, and CYP3A4/5). (7)

USE IN SPECIFIC POPULATIONS

Renal impairment: Use balsalazide with caution in patients with a history of renal disease. (5.3)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 9/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Adult Dose

2.2 Pediatric Dose

2.3 Administration Alternatives

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Exacerbations of Ulcerative Colitis

5.2 Pyloric Stenosis

5.3 Renal

6 ADVERSE REACTIONS

6.1 Clinical Studies Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology

14 CLINICAL STUDIES

14.1 Adult Studies

14.2 Pediatric Studies

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

17.1 Important Precautions Regarding Balsalazide

17.2 What Patients Should Know About Adverse Reactions

17.3 What Patients Should Know About Taking Balsalazide with Other Medication

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Balsalazide disodium capsules are indicated for the treatment of mildly to moderately active ulcerative colitis in patients 5 years of age and older. Safety and effectiveness of balsalazide beyond 8 weeks in children (ages 5 to 17 years) and 12 weeks in adults have not been established.

2 DOSAGE AND ADMINISTRATION

2.1 Adult Dose

For treatment of active ulcerative colitis in adult patients, the usual dose is three 750 mg balsalazide disodium capsules to be taken 3 times a day (6.75 g per day) for up to 8 weeks. Some patients in the adult clinical trials required treatment for up to 12 weeks.

2.2 Pediatric Dose

For treatment of active ulcerative colitis in pediatric patients, aged 5 to 17 years, the usual dose is EITHER:

- three 750 mg balsalazide disodium capsules 3 times a day (6.75 g per day) for up to 8 weeks; OR:
- one 750 mg balsalazide disodium capsule 3 times a day (2.25 g per day) for up to 8 weeks.

Use of balsalazide in the pediatric population for more than 8 weeks has not been evaluated in clinical trials. [See *Clinical Studies Section (14)*]

2.3 Administration Alternatives

Balsalazide disodium capsules may also be administered by carefully opening the capsule

and sprinkling the capsule contents on applesauce. The entire drug/applesauce mixture should be swallowed immediately; the contents may be chewed, if necessary, since contents of balsalazide disodium capsules are NOT coated beads/granules. Patients should be instructed not to store any drug/applesauce mixture for future use.

If the capsules are opened for sprinkling, color variation of the powder inside the capsules ranges from orange to yellow and is expected due to color variation of the active pharmaceutical ingredient.

Teeth and/or tongue staining may occur in some patients who use balsalazide disodium capsules in sprinkle form with food.

3 DOSAGE FORMS AND STRENGTHS

Balsalazide disodium capsules, USP are available as opaque white capsules containing 750 mg balsalazide disodium, imprinted with "APO B750" in red ink.

4 CONTRAINDICATIONS

Patients with hypersensitivity to salicylates or to any of the components of balsalazide disodium capsules or balsalazide metabolites. Hypersensitivity reactions may include, but are not limited to the following: anaphylaxis, bronchospasm, and skin reaction.

5 WARNINGS AND PRECAUTIONS

5.1 Exacerbations of Ulcerative Colitis

In the adult clinical trials, 3 out of 259 patients reported exacerbation of the symptoms of ulcerative colitis. In the pediatric clinical trials, 4 out of 68 patients reported exacerbation of the symptoms of ulcerative colitis.

Observe patients closely for worsening of these symptoms while on treatment.

5.2 Pyloric Stenosis

Patients with pyloric stenosis may have prolonged gastric retention of balsalazide disodium capsules.

5.3 Renal

Renal toxicity has been observed in animals and patients given other mesalamine products. Therefore, caution should be exercised when administering balsalazide capsules to patients with known renal dysfunction or a history of renal disease. [See *Nonclinical Toxicology (13.2)*]

6 ADVERSE REACTIONS

6.1 Clinical Studies Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

Adult Ulcerative Colitis

During clinical development, 259 adult patients with active ulcerative colitis were exposed to 6.75 g/day balsalazide in 4 controlled trials.

In the 4 controlled clinical trials patients receiving a balsalazide dose of 6.75 g/day most frequently reported the following adverse reactions: headache (8%), abdominal pain (6%), diarrhea (5%), nausea (5%), vomiting (4%), respiratory infection (4%), and arthralgia (4%). Withdrawal from therapy due to adverse reactions was comparable among patients on balsalazide and placebo.

Adverse reactions reported by 1% or more of patients who participated in the 4 well-controlled, Phase 3 trials are presented by treatment group (Table 1).

The number of placebo patients (35), however, is too small for valid comparisons. Some adverse reactions, such as abdominal pain, fatigue, and nausea were reported more frequently in women than in men. Abdominal pain, rectal bleeding, and anemia can be part of the clinical presentation of ulcerative colitis.

Table 1: Adverse Reactions Occurring in ≥ 1 % of Adult Balsalazide Patients in Controlled Trials*

Adverse Reaction	Balsalazide Capsules 6.75 g/day [N=259]	Placebo [N=35]
Abdominal pain	16 (6%)	1 (3%)
Diarrhea	14 (5%)	1 (3%)
Arthralgia	9 (4%)	0%
Rhinitis	6 (2%)	0%
Insomnia	6 (2%)	0%
Fatigue	6 (2%)	0%
Flatulence	5 (2%)	0%
Fever	5 (2%)	0%
Dyspepsia	5 (2%)	0%
Pharyngitis	4 (2%)	0%
Coughing	4 (2%)	0%
Anorexia	4 (2%)	0%
Urinary tract infection	3 (1%)	0%
Myalgia	3 (1%)	0%
Flu-like disorder	3 (1%)	0%
Dry mouth	3 (1%)	0%
Cramps	3 (1%)	0%
Constipation		

*Adverse reactions occurring in at least 1 % of balsalazide Patients which were less frequent than placebo for the same event were not included in the table.

Pediatric Ulcerative Colitis

In a clinical trial in 68 pediatric patients aged 5 to 17 years with mildly to moderately active ulcerative colitis who received 6.75 g/day or 2.25 g/day balsalazide disodium for 8

weeks, the most frequently reported adverse reactions were headache (15%), abdominal pain upper (13%), abdominal pain (12%), vomiting (10%), diarrhea (9%), colitis ulcerative (6%), nasopharyngitis (6%), and pyrexia (6%). [see Table 2]

One patient who received balsalazide disodium 6.75 g/day and 3 patients who received balsalazide disodium 2.25 g/day discontinued treatment because of adverse reactions. In addition, 2 patients in each dose group discontinued because of a lack of efficacy.

Adverse reactions reported by 3% or more of pediatric patients within either treatment group in the Phase 3 trial are presented in Table 2.

Table 2: Treatment-Emergent Adverse Reactions Reported by ≥3% of Patients in Either Treatment Group in a Controlled Study of 68 Pediatric Patients

Adverse Reaction	balsalazide disodium		
	6.75 g/day [N=33]	2.25 g/day [N=35]	Total [N=68]
Headache	5 (15%)	5 (14%)	10 (15%)
Abdominal pain upper	3 (9%)	6 (17%)	9 (13%)
Abdominal pain	4 (12%)	4 (11%)	8 (12%)
Vomiting	1 (3%)	6 (17%)	7 (10%)
Diarrhea	2 (6%)	4 (11%)	6 (9%)
Colitis ulcerative	2 (6%)	2 (6%)	4 (6%)
Nasopharyngitis	3 (9%)	1 (3%)	4 (6%)
Pyrexia	0 (0%)	4 (11%)	4 (6%)
Hematochezia	0 (0%)	3 (9%)	3 (4%)
Nausea	0 (0%)	3 (9%)	3 (4%)
Influenza	1 (3%)	2 (6%)	3 (4%)
Fatigue	2 (6%)	1 (3%)	3 (4%)
Stomatitis	0 (0%)	2 (6%)	2 (3%)
Cough	0 (0%)	2 (6%)	2 (3%)
Pharyngolaryngeal pain	2 (6%)	0 (0%)	2 (3%)
Dysmenorrhea	2 (6%)	0 (0%)	2 (3%)

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of balsalazide in clinical practice:

myocarditis, pericarditis, vasculitis, pruritus, pleural effusion, pneumonia (with and without eosinophilia), alveolitis, renal failure, interstitial nephritis, pancreatitis, and alopecia.

Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. These adverse reactions have been chosen for inclusion due to a combination of seriousness, frequency of reporting, or potential causal connection to balsalazide.

Hepatic

Postmarketing adverse reactions of hepatotoxicity have been reported for products which contain (or are metabolized to) mesalamine, including elevated liver function tests (SGOT/AST, SGPT/ALT, GGT, LDH, alkaline phosphatase, bilirubin), jaundice, cholestatic jaundice, cirrhosis, hepatocellular damage including liver necrosis and liver failure. Some of these cases were fatal; however, no fatalities associated with these adverse reactions were reported in balsalazide clinical trials. One case of Kawasaki-like syndrome which

included hepatic function changes was also reported, however, this adverse reaction was not reported in balsalazide clinical trials.

7 DRUG INTERACTIONS

In an in vitro study using human liver microsomes, balsalazide and its metabolites [5-aminosalicylic acid (5-ASA), N-acetyl-5-aminosalicylic acid (N-Ac-5-ASA), 4-aminobenzoyl- β -alanine (4-ABA) and N-acetyl-4-aminobenzoyl- β -alanine (N-Ac-4-ABA)] were not shown to inhibit the major CYP enzymes evaluated (CYP1A2, CYP2C9, CYP2C19, CYP2D6, and CYP3A4/5). Therefore, balsalazide and its metabolites are not expected to inhibit the metabolism of other drugs which are substrates of CYP1A2, CYP2C9, CYP2C19, CYP2D6, or CYP3A4/5.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B. Reproduction studies were performed in rats and rabbits at oral doses up to 2 g/kg/day, 2.4 and 4.7 times the recommended human dose based on body surface area for the rat and rabbit, respectively, and revealed no evidence of impaired fertility or harm to the fetus due to balsalazide disodium. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

8.3 Nursing Mothers

It is not known whether balsalazide disodium is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when balsalazide is administered to a nursing woman.

8.4 Pediatric Use

Use of balsalazide disodium in pediatric and adolescent patients 5 to 17 years of age for the treatment of mildly to moderately active ulcerative colitis is supported by:

- extrapolation of results from clinical studies that supported the approval of balsalazide disodium for adults.
- a clinical trial of 68 patients ages 5 to 17 years comparing two doses of balsalazide disodium (6.75 g/day and 2.25 g/day), and
- a pharmacokinetic study performed on a subset of the pediatric study population. [See *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14)*].

Based on the limited data available, dosing can be initiated at either 6.75 or 2.25 g/day.

Safety and efficacy of balsalazide disodium in pediatric patients below the age of 5 years have not been established.

10 OVERDOSAGE


No case of overdose has occurred with balsalazide. A 3-year-old boy is reported to have ingested 2 g of another mesalamine product. He was treated with ipecac and activated charcoal with no adverse reactions.

If an overdose occurs with balsalazide, treatment should be supportive, with particular attention to correction of electrolyte abnormalities.

11 DESCRIPTION

Each balsalazide disodium capsule, USP contains 750 mg of balsalazide disodium, a prodrug that is enzymatically cleaved in the colon to produce mesalamine (5-aminosalicylic acid or 5-ASA), an anti-inflammatory drug. Each capsule of balsalazide (750 mg) is equivalent to 267 mg of mesalamine. Balsalazide disodium has the chemical

name (E)-5-[[4-[[2-(2-carboxyethyl)amino]carbonyl]phenyl]azo]-2-hydroxybenzoic acid, disodium salt, dihydrate. Its structural formula is:



Molecular Weight: 437.32

Molecular Formula: C₁₇H₁₃N₃O₆Na₂•2H₂O

Balsalazide disodium is a stable, odorless orange to yellow microcrystalline powder. It is freely soluble in water and isotonic saline, sparingly soluble in methanol and ethanol, and practically insoluble in all other organic solvents.

Inactive Ingredients: Each hard gelatin capsule contains colloidal silicon dioxide, magnesium stearate and titanium dioxide. The capsule imprinting ink contains propylene glycol, strong ammonia solution, shellac and red iron oxide. The sodium content of each capsule is approximately 86 mg.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Balsalazide disodium is delivered intact to the colon where it is cleaved by bacterial azoreduction to release equimolar quantities of mesalamine, which is the therapeutically active portion of the molecule, and the 4-aminobenzoyl-β-alanine carrier moiety. The carrier moiety released when balsalazide disodium is cleaved is only minimally absorbed and is largely inert.

The mechanism of action of 5-ASA is unknown, but appears to be local to the colonic mucosa rather than systemic. Mucosal production of arachidonic acid metabolites, both through the cyclooxygenase pathways, i.e., prostanooids, and through the lipoxygenase pathways, i.e., leukotrienes and hydroxyeicosatetraenoic acids, is increased in patients with chronic inflammatory bowel disease, and it is possible that 5-ASA diminishes inflammation by blocking production of arachidonic acid metabolites in the colon.

12.3 Pharmacokinetics

Balsalazide disodium capsules contain a powder of balsalazide disodium that is insoluble in acid and designed to be delivered to the colon as the intact prodrug. Upon reaching the colon, bacterial azoreductases cleave the compound to release 5-ASA, the therapeutically active portion of the molecule, and 4-aminobenzoyl-β-alanine. The 5-ASA is further metabolized to yield Nacetyl-5-aminosalicylic acid (N-Ac-5-ASA), a second key metabolite.

Absorption

The plasma pharmacokinetics of balsalazide and its key metabolites from a crossover study in healthy volunteers are summarized in Table 3. In this study, a single oral dose of balsalazide 2.25 g was administered to healthy volunteers as intact capsules (3 x 750 mg) under fasting conditions, as intact capsules (3 x 750 mg) after a high-fat meal, and unencapsulated (3 x 750 mg) as sprinkles on applesauce.

Table 3: Plasma Pharmacokinetics for Balsalazide and Key Metabolites (5-ASA and N-Ac-5-ASA) with Administration of Balsalazide Following a Fast, a High-Fat Meal, and Drug Contents Sprinkled on Applesauce (Mean ± SD)

	Fasting n = 17	High-Fat Meal n = 17	Sprinkled n = 17
C _{max} (mcg/mL)			
Balsalazide	0.51 ± 0.32	0.45 ± 0.39	0.21 ± 0.12
5-ASA	0.22 ± 0.12	0.11 ± 0.136	0.29 ± 0.17

N-Ac-5-ASA AUC _{last} mcg·hr/mL)	0.88 ± 0.39	0.64 ± 0.534	1.04 ± 0.57
Balsalazide	1.35 ± 0.73	1.52 ± 1.01	0.87 ± 0.48
5-ASA	2.59 ± 1.46	2.10 ± 2.58	2.99 ± 1.70
N-Ac-5-ASA T _{max} (h)	17.8 ± 8.14	17.7 ± 13.7	20.0 ± 11.4
Balsalazide	0.8 ± 0.85	1.2 ± 1.11	1.6 ± 0.44
5-ASA	8.2 ± 1.98	22.0 ± 8.23	8.7 ± 1.99
N-Ac-5-ASA	9.9 ± 2.49	20.2 ± 8.94	10.8 ± 5.39

A relatively low systemic exposure was observed under all three administered conditions (fasting, fed with high-fat meal, sprinkled on applesauce), which reflects the variable, but minimal absorption of balsalazide disodium and its metabolites. The data indicate that both C_{max} and AUC_{last} were lower, while t_{max} was markedly prolonged, under fed (high-fat meal) compared to fasted conditions. Moreover, the data suggest that dosing balsalazide disodium as a sprinkle or as a capsule provides highly variable, but relatively similar mean pharmacokinetic parameter values. No inference can be made as to how the systemic exposure differences of balsalazide and its metabolites in this study might predict the clinical efficacy under different dosing conditions (i.e., fasted, fed with high-fat meal, or sprinkled on applesauce) since clinical efficacy after balsalazide disodium administration is presumed to be primarily due to the local effects of 5-ASA on the colonic mucosa.

In a separate study of adult patients with ulcerative colitis, who received balsalazide, 1.5 g twice daily, for over 1 year, systemic drug exposure, based on mean AUC values, was up to 60 times greater (0.008 mcg·hr/mL to 0.480 mcg·hr/mL) when compared to that obtained in healthy subjects who received the same dose.

Distribution

The binding of balsalazide to human plasma proteins was ≥99%.

Metabolism

The products of the azoreduction of this compound, 5-ASA and 4-aminobenzoyl-β-alanine, and their N-acetylated metabolites have been identified in plasma, urine and feces.

Elimination

Following single-dose administration of 2.25 g balsalazide (three 750 mg capsules) under fasting conditions in healthy subjects, mean urinary recovery of balsalazide, 5-ASA, and N-Ac-5-ASA was 0.20%, 0.22% and 10.2%, respectively.

In a multiple-dose study in healthy subjects receiving a balsalazide dose of two 750 mg capsules twice daily (3 g/day) for 10 days, mean urinary recovery of balsalazide, 5-ASA, and N-Ac-5-ASA was 0.1%, 0%, and 11.3%, respectively. During this study, subjects received their morning dose 0.5 hours after being fed a standard meal, and subjects received their evening dose 2 hours after being fed a standard meal.

In a study with 10 healthy volunteers, 65% of a single 2.25-gram dose of balsalazide was recovered as 5-ASA, 4-aminobenzoyl-β-alanine, and the N-acetylated metabolites in feces, while <1% of the dose was recovered as parent compound.

In a study that examined the disposition of balsalazide in patients who were taking 3 to 6 g of balsalazide daily for more than 1 year and who were in remission from ulcerative colitis, less than 1% of an oral dose was recovered as intact balsalazide in the urine. Less than 4% of the dose was recovered as 5-ASA, while virtually no 4-aminobenzoyl-β-alanine was detected in urine. The mean urinary recovery of N-Ac-5-ASA and N-acetyl-4-aminobenzoyl-β-alanine comprised <16% and <12% of the balsalazide dose, respectively. No fecal recovery studies were performed in this population.

Pediatric Population

In studies of pediatric patients with mild-to-moderate active ulcerative colitis receiving three 750 mg balsalazide disodium capsules 3 times daily (6.75 g/day) for 8 weeks,

steady state was reached within 2 weeks, as observed in adult patients. Likewise, the pharmacokinetics of balsalazide, 5-ASA, and N-Ac-5-ASA were characterized by very large inter-patient variability, which is also similar to that seen in adult patients.

The pro-drug moiety, balsalazide, appeared to exhibit dose-independent (i.e., doselinear) kinetics in children, and the systemic exposure parameters (C_{max} and $AUC_{0\text{ to }8}$) increased in an almost dose-proportional fashion after the 6.75 g/day versus the 2.25 g/day doses. However, the absolute magnitude of these exposure parameters was greater relative to adults. The C and AUC observed in max 0 to 8 pediatric patients were 26% and 102% greater than those observed in adult patients at the 6.75 g/day dosage level. In contrast, the systemic exposure parameters for the active metabolites, 5-ASA and N-Ac-5-ASA, in pediatric patients increased in a less than dose-proportional manner after the 6.75 g/day dose versus the 2.25 g/day dose. Additionally, the magnitude of these exposure parameters was decreased for both metabolites relative to adults. For the metabolite of key safety concern from a systemic exposure perspective, 5-ASA, the C and AUC observed in pediatric patients were max 0 to 8 67% and 64% lower than those observed in adult patients at the 6.75 g/day dosage level. Likewise, for N-Ac-5-ASA, the C and AUC observed in pediatric patients were 68% and 55% lower than max 0 to 8 those observed in adult patients at the 6.75 g/day dosage level.

All pharmacokinetic studies with balsalazide are characterized by large variability in the plasma concentration versus time profiles for balsalazide and its metabolites, thus half-life estimates of these analytes are indeterminate.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 24-month rat (Sprague Dawley) carcinogenicity study, oral (dietary) balsalazide disodium at doses up to 2 g/kg/day was not tumorigenic. For a 50-kg person of average height this dose represents 2.4 times the recommended human dose on a body surface area basis. Balsalazide disodium was not genotoxic in the following *in vitro* or *in vivo* tests: Ames test, human lymphocyte chromosomal aberration test, and mouse lymphoma cell (L5178Y/TK+/-) forward mutation test, or mouse micronucleus test. However, it was genotoxic in the *in vitro* Chinese hamster lung cell (CH V79/HGPRT) forward mutation test.

4-aminobenzoyl- β -alanine, a metabolite of balsalazide disodium, was not genotoxic in the Ames test and the mouse lymphoma cell (L5178Y/TK+/-) forward mutation test but was positive in the human lymphocyte chromosomal aberration test. N-acetyl-4-aminobenzoyl- β -alanine, a conjugated metabolite of balsalazide disodium, was not genotoxic in Ames test, the mouse lymphoma cell (L5178Y/TK+/-) forward mutation test, or the human lymphocyte chromosomal aberration test. Balsalazide disodium at oral doses up to 2 g/kg/day, 2.4 times the recommended human dose based on body surface area, was found to have no effect on fertility and reproductive performance in rats.

13.2 Animal Toxicology

Renal Toxicity

In animal studies conducted at doses up to 2000 mg/kg (approximately 21 times the recommended 6.75 g/day dose on a mg/kg basis for a 70 kg person), balsalazide demonstrated no nephrotoxic effects in rats or dogs.

Overdosage


A single oral dose of balsalazide disodium at 5 g/kg or 4-aminobenzoyl- β -alanine, a metabolite of balsalazide disodium, at 1 g/kg was non-lethal in mice and rats. No symptoms of acute toxicity were seen at these doses.

14 CLINICAL STUDIES

14.1 Adult Studies

Two randomized, double-blind studies were conducted in adults. In the first trial, 103 patients with active mild-to-moderate ulcerative colitis with sigmoidoscopy findings of friable or spontaneously bleeding mucosa were randomized and treated with balsalazide 6.75 g/day or balsalazide 2.25 g/day. The primary efficacy endpoint was reduction of rectal bleeding and improvement of at least one of the other assessed symptoms (stool frequency, patient functional assessment, abdominal pain, sigmoidoscopic grade, and physician's global assessment [PGA]). Outcome assessment for rectal bleeding at each interim period (week 2, 4, and 8) encompassed a 4-day period (96 hours). Results demonstrated a statistically significant difference between high and low doses of balsalazide (Figure 1).

Figure 1: Percentage of Patients Improved at 8 weeks



A second study, conducted in Europe, confirmed findings of symptomatic improvement.

14.2 Pediatric Studies

A clinical trial was conducted comparing two doses (6.75 g/day and 2.25 g/day) of balsalazide disodium in 68 pediatric patients (age 5 to 17, 23 males and 45 females) with mildly to moderately active ulcerative colitis. 28/33 (85%) patients randomized to 6.75 g/day and 25/35 (71%) patients randomized to 2.25 g/day completed the study. The primary endpoint for this study was the proportion of subjects with clinical improvement (defined as a reduction of at least 3 points in the Modified Sutherland Ulcerative Colitis Activity Index [MUCAI] from baseline to 8 weeks). Fifteen (45%) patients in the balsalazide disodium 6.75 g/day group and 13 (37%) patients in the balsalazide disodium 2.25 g/day group showed this clinical improvement. In both groups, patients with higher MUCAI total scores at baseline were likely to experience greater improvement.

Rectal bleeding improved in 64% of patients treated with balsalazide disodium 6.75 g/day and 54% of patients treated with balsalazide disodium 2.25 g/day. Colonic mucosal appearance upon endoscopy improved in 61% of patients treated with balsalazide disodium 6.75 g/day and 46% of patients treated with balsalazide disodium 2.25 g/day.

16 HOW SUPPLIED/STORAGE AND HANDLING

Balsalazide Disodium Capsules, USP are available as white, opaque capsules imprinted "APO B750" in red ink.

Balsalazide Disodium Capsules, USP are supplied as follows:

NDC 50268-102-13 (10 capsules per car, 3 cards per carton)

Dispensed in Unit Dose package. For Institutional Use Only.

Storage

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° and 30°C (59° and 86°F).

See USP Controlled Room Temperature.

17 PATIENT COUNSELING INFORMATION

17.1 Important Precautions Regarding Balsalazide

- Instruct patients not to take balsalazide if they have a hypersensitivity to salicylates (e.g., aspirin).
- Patients should be instructed to contact their health care provider under the following circumstances:
 - If they experience a worsening of their ulcerative colitis symptoms.
 - If they are diagnosed with pyloric stenosis, because balsalazide disodium capsules may be slow to pass through their digestive tract.
 - If they are diagnosed with renal dysfunction. Damage to the kidney has been observed in people given medications similar to balsalazide.

17.2 What Patients Should Know About Adverse Reactions

- In adult clinical trials the most common adverse reactions were headache, abdominal pain, diarrhea, nausea, vomiting, respiratory infection, and arthralgia.
- In the pediatric clinical trial the most common adverse reactions were headache, abdominal pain, vomiting, diarrhea, ulcerative colitis, nasopharyngitis, and pyrexia.
- Inform patients that this listing of adverse reactions is not complete and not all adverse reactions can be anticipated. If appropriate, a more comprehensive list of adverse reactions can be discussed with patients.

17.3 What Patients Should Know About Taking Balsalazide with Other Medication

- Based upon limited studies conducted in a test tube, balsalazide is not believed to interfere with other drugs by preventing how the liver functions. However, as the studies were limited in scope, you should always consult your doctor and discuss potential interactions prior to initiating any new drug.

BALSALAZIDE DISODIUM CAPSULES, USP 750 mg

Manufactured for:
AvKARE, Inc.
Pulaski, TN 38478

Mfg. Rev. 03/16

AV Rev. 09/17 (P)

AvPAK

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 50268-102-13

Balsalazide Disodium Capsules, USP

750 mg

Rx Only

30 Capsules (3 X 10) Unit Dose



5026810213

NDC 50268-102-13

Balsalazide Disodium Capsules, USP

750 mg

Rx Only

30 Capsules (3 X 10) Unit Dose



5026810213

Each capsule contains 750 mg of balsalazide disodium.

Usual Dosage: See package insert.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Keep this and all medications out of the reach of children.

Manufactured for:
AvKARE, Inc.
Pulaski, TN 38478



Mfg. Rev. 03/16

AV Rev. 09/17 (P)

BALSALAZIDE DISODIUM

balsalazide disodium capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:50268-102(NDC:60505-2575)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
BALSALAZIDE DISODIUM (UNII: 1XL6BJI034) (BALSALAZIDE - UNII:P80AL8J7ZP)		BALSALAZIDE DISODIUM	750 mg	
Inactive Ingredients				
Ingredient Name		Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
AMMONIA (UNII: 5138Q19F1X)				
SHELLAC (UNII: 46N107B71O)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
Product Characteristics				
Color	white	Score	no score	
Shape	CAPSULE	Size	23mm	
Flavor		Imprint Code	APO;B750	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50268-102-13	30 in 1 BOX	11/06/2014	
1	NDC:50268-102-11	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA077883	11/06/2014		

Labeler - AvPAK (832926666)

Revised: 1/2024

AvPAK